



## Erratum

Erratum to “Retrieval of the aerosol size distribution in the complex anomalous diffraction approximation” [Atmospheric Environment 35 (30) 5099–5104] <sup>☆</sup>

Ghislain R. Franssens

*Belgian Institute for Space Aeronomy, Ringlaan 3, B-1180 Brussels, Belgium*

The Publisher regrets that in the printed version of the above paper, formulas 2, 9, and the formula on page 5101 in section 3, (line 4) were printed incorrectly. The correct versions of these formulas are printed below.

$$Q_{\text{ext}}(k, r) \triangleq \frac{C(k, r)}{\pi r^2} = \frac{2}{(kr)^2} \text{Re} \left[ \sum_{l=1}^{+\infty} (2l+1)(a_l(k, r) + b_l(k, r)) \right] \quad (2)$$

$$\kappa(k) \triangleq 2[(n_r(k) - 1) - in_i(k)]k \quad (\text{page 5101, section 3, line 4})$$

$$\tau(\kappa) = N_{\text{tot}} \left\{ 2\pi(a+2)(a+1)r_0^2 + 4\pi \text{Re} \left[ \frac{1}{(i2\kappa)^2} \left( -1 + \frac{1 + (a+2)i2\kappa r_0}{(1 + i2\kappa r_0)^{a+2}} \right) \right] \right\} \quad (9)$$

<sup>☆</sup>PII of original article: S1352-2310(01)00331-4.