

SIMULTANEOUS MEASUREMENTS OF NO AND NO₂ IN THE STRATOSPHERE

M. ACKERMAN,* J. C. FONTANELLA,† D. FRIMOUT,* A. GIRARD,†
N. LOUISNARD† and C. MULLER*

(Received 30 July 1974)

Abstract—Simultaneous measurements of NO and NO₂ in the stratosphere leading to an NO_x determination have been performed by means of i.r. absorption spectrometry using the Sun as a source in the 5.2 μm band of NO and in the 6.2 μm band of NO₂. The observed abundance of NO_x peaks at 26 km where it is equal to $(4.2 \pm 1) \times 10^9 \text{ cm}^{-3}$. The volume mixing ratio of NO_x was observed to vary from 1.3×10^{-9} at 20 km to 1.3×10^{-8} at 34 km.

1. INTRODUCTION

Since the first detection of nitric oxide in the stratosphere (Toth *et al.*, 1973, Girard *et al.*, 1973 and Ridley *et al.*, 1973) useful information about the vertical distribution of this constituent have been derived by absorption spectrometry using the Sun as a source (Ackerman *et al.*, 1973) and by chemiluminescence (Ridley *et al.*, 1974). More recently the variation of NO vs time at 28 km after sunrise has been observed by spin flit Raman laser spectrometry by Patel *et al.* (1974) and observations from a very high flying aircraft have been performed by Loewenstein *et al.* (1974).

Separate determinations of the abundance of NO₂ have been derived from absorption spectra (Ackerman and Muller, 1972, 1973; Farmer, 1974; Fontanella *et al.*, 1974) from emission spectra (Harries 1973) and from photometric observations in the visible (Brewer *et al.*, 1973).

As it is known that NO and NO₂ are photochemically linked through the reaction of nitric oxide with ozone and the photodissociation of NO₂ by the solar radiation with a time constant that is very short compared with day length, it is most useful to determine both constituents simultaneously. This appears even more crucial when it is realized that the rate of conversion of NO into NO₂ at sunset and of NO₂ into NO at sunrise is altitude dependent due to the variation with height of the attenuation of solar radiation by Rayleigh scattering and by ozone and NO₂ absorptions and due to the variation with height of the abundance of ozone as an NO scavenger.

A first step towards the determination of the natural variability of NO_x requires the simultaneous measurement of NO and NO₂. Such a measurement is presented here. The method involves absorption spectrometry, using the Sun as a source, in two narrow spectral ranges: 1913–1923 cm⁻¹ in the 5.2 μm band of NO and 1593–1603 cm⁻¹ in the 6.2 μm band of NO₂.


2. EXPERIMENTAL DETAILS

A grille spectrometer (Girard, 1963) associated with a telescope and a Sun pointer has been used to record the spectra.

* Institut d'Aéronomie Spatiale de Belgique, 3, avenue Circulaire, B-1180 Bruxelles, Belgique.

† Office National d'Etudes et de Recherches Aérospatiales, 29, avenue de la Division Leclerc, F-92320 Chatillon, Belgique.

The spectrometer uses a Littrow mounting with an off-axis parabolic mirror whose focal length is 60 cm. Its Jobin-Yvon grating ruled at 60 grooves per mm over an area of 130×68.6 mm is blazed at 70° . It is used in the 5th and in the 6th orders associated with two band pass interference filters. At every inversion of the grating drive the filter wheel switches from one filter to the other in order to scan the two wavelength ranges shown in Fig. 1. The grille is deposited on an 18×18 mm calcium fluoride window and the minimum step is 0.2 mm. The resolution approaches the theoretical limit. The signal is chopped at 360 Hz by vibrating the collimating mirror at 180 Hz, which is the resonance frequency of the mechanical mounting included in the servo system stabilizing the vibration amplitude. The spectral calibrations are performed by means of H_2O and CO_2 absorption lines and by the simultaneous recording of the spectra and of position grating marks. To maintain the instrumental performance during flight the optical path inside the spectrometer is servo controlled to balance the mechanical deformations due to temperature and attitude alterations. Two auxiliary grilles located alongside of the main grille were used for this purpose. The first was illuminated by a filament lamp, with ray paths close to the path of the solar radiation to be measured. The light flux was received on a germanium photodiode after reflection on the second grille. The parabolic mirror vibrating at a frequency f induces a photodiode signal at frequencies f and $2f$. The f component called error signal is used to keep the instrument properly tuned. A gold doped germanium detector was used at solid nitrogen temperature. The detector signals were preamplified and telemetered to the ground



through three channels having different sensitivities. The linearity of the system was controlled in flight by means of calibration voltages as shown in Fig. 1.

A 4.2 m focal length modified Cassegrain telescope of 32 cm diameter was used to image the Sun on the grille.

The Sun pointing system controls the attitude of the spectrometer telescope assembly. The whole gondola is uncoupled from the balloon on the three axis. In a first mode the gondola is oriented along a fixed axis referred to the Earth. The couples are provided by inertia wheels desaturated by means of moving masses. At the end of the acquisition mode a rendez-vous is realized with the Sun. A timer switches then to the tracking mode involving optical Sun sensors. The pointing accuracy is better than 30 arc sec.

Temperatures and ambient pressure were also telemetered as described in more detail by Ackerman *et al.* (1973).

3. THE FLIGHT

The gondola was flown on 13 May 1974 from the CNES range in Aire sur l'Adour (France). A Winzen 11.6 million cubic ft balloon carried the payload up to 38.5 km after 3 hr of ascent. As usual at that period of the winds inversion, the balloon remained practically at the same geographic coordinates during the measurements namely 0°25' East and 43°35' North. The recording of spectra started at a solar zenith angle equal to 86° showing the solar continuum and small absorption lines due to H₂O. Solar absorption lines were also present in the 1900 cm⁻¹ range (Kunde, 1967). Sample spectra obtained at solar zenith angles larger than 90° are shown in Fig. 1. The two spectral ranges scanned in sequence show telluric absorption features due to NO₂, NO, H₂O and CO₂.

4. INTERPRETATION OF THE DATA

The identification of the lines has been based on the laboratory measurements of Hurlock *et al.* (1974) and of Dana and Fontanella (1973) for NO₂, on the compilation made by Cieslik and Muller (1974) for NO and on the compilation of McClatchey *et al.* (1973) for H₂O and for CO₂. The quantitative evaluation of the amount of NO and NO₂ on the optical path has been based on the measured equivalent widths and on the absorption cross sections whose values are listed in Table 1 and have been adapted for the stratospheric temperatures. For NO the line of 1914.993 cm⁻¹ has been chosen for the sake of comparison of the results of this flight with those of a previous flight for which preliminary data have already been published (Ackerman *et al.*, 1973).

Figures 2(a) and (b) show the evolution of the absorption due to this NO line while the minimum altitude reached by the solar rays decreases in the cases of the May 1973 and May 1974 flights respectively. Figure 3 shows the evolution of NO₂ absorption lines during the increase of solar depression angles corresponding to lower and lower minimum altitudes

TABLE 1. INTEGRATED ABSORPTION CROSS SECTIONS, s , USED IN THE INTERPRETATION OF THE DATA

Constituent	Wavenumber (cm ⁻¹)	s (300 K) (cm ² cm ⁻¹ mole ⁻¹)
NO ₂	1914.993	9.56×10^{-20}
NO ₂	1600.411	3.18×10^{-19}
NO ₂	1600.058	1.70×10^{-19}
NO ₂	1596.946	2.11×10^{-19}




FIG. 2(a). INTENSITY OF SOLAR RADIATION RECORDED VS WAVENUMBER DURING THE EXPERIMENT PERFORMED ON 14 MAY 1973 (ACKERMAN *et al.*, 1973) AT 40 km ALTITUDE.

The spectral region from 1913.5 cm^{-1} to 1915.3 cm^{-1} recorded during sunset is shown. The upper spectrum corresponds to a solar zenith angle, χ , equal to 89° . For each of the other spectra χ was larger than 90° and the minimum altitude reached by the solar radiation in the atmosphere is indicated. The NO line at 1914.993 cm^{-1} appears clearly, well separated from the solar absorption line at 1914.73 cm^{-1} . It is well separated from the closest CO_2 lines due to the $(12^202)-(01^101)$ transitions that occur at 1914.541 cm^{-1} and 1915.328 cm^{-1} .

of the solar rays. The equivalent width is shown in Fig. 4 for NO and NO_2 vs minimum altitude reached by the solar radiation in the atmosphere. These values are used in the inversion process involving the division of the atmosphere in layers of 2 km thickness. The correct amount of absorber is introduced in the successive layers in order to reproduce the observed integrated absorptions by means of an iterative Newton-Raphson process (Hartree, 1958). The Voigt line profile is represented by the approximation derived by Whiting (1968).




FIG. 2(b) INTENSITY OF SOLAR RADIATION RECORDED VS WAVENUMBER DURING THE EXPERIMENT PERFORMED ON 13 MAY 1974 AT 38.5 km ALTITUDE.

An intensity change corresponding to 10 per cent of absorption at the position of the NO line at 1914.993 cm^{-1} is indicated. The other main absorptions are identified in Fig. 1. The rate of growth is obviously different for the NO line and for the CO₂ lines while the minimum altitude reached by the solar radiation in the atmosphere decreases. The altitude is indicated in km for each spectrum excepted for the upper one obtained at a solar zenith angle χ equal to 90° .

5. RESULTS AND DISCUSSION

The deduced vertical distributions are shown in Figs. 5 and 6 for NO and NO₂ respectively. Numerical values are listed in Table 2 for NO, NO₂, NO + NO₂ and for the NO₂ over NO ratio. The sum of NO and NO₂ is plotted versus altitude in Fig. 7. The limits of uncertainties indicated in Table 2 and in Fig. 5–7 are given to take into account the uncertainty in determining the equivalent widths from the spectra, amplified by the inversion process leading to the vertical distributions of number densities. They do not include the effects on the inversion process that could be due to the conversion of NO into NO₂ by reaction with ozone when the photodissociation of the latter decreases. Theoretical considerations (Brasseur and Nicolet, 1973) indicate that for an average ozone profile the NO₂ over NO ratio should exhibit a broad maximum of its day-time value close to one in the 20–30 km altitude range. The observed values listed versus altitude in Table 2 indicate that the measurements still correspond to day time ratios down to at least 25 km. The sunset conversion process has no significant effect on the values given here for (NO + NO₂) over the whole altitude range considered within the given limits of uncertainties. The range of uncertainty is larger for the NO measurements of May 1973 than for those of May 1974. This is due to the smaller equivalent widths observed in the first case for which no value is given for altitudes lower than 24 km where the uncertainty appears to be too large.




FIG. 3. INTENSITY OF SOLAR RADIATION RECORDED VS WAVENUMBER DURING THE EXPERIMENT PERFORMED ON 13 MAY 1974.

The spectral region shown is in the range of the $6.2 \mu\text{m}$ band of NO_2 . The lines are identified in Fig. 1. The upper spectrum corresponds to a solar zenith angle, χ , equal to 89.5° . For each of the other spectra χ was larger than 90° and the minimum altitude reached by the solar radiation in the atmosphere is indicated in km. Two of the lines selected for the quantitative interpretation are indicated. An intensity change corresponding to 10 per cent of absorption is shown.




FIG. 4. INTEGRATED ABSORPTION EXPRESSED IN cm^{-1} DEDUCED FROM THE EQUIVALENT WIDTHS MEASURED THAT HAVE BEEN USED IN THE INVERSION PROCESS TO DETERMINE THE NUMBER DENSITIES. Each value corresponds to a minimum altitude expressed in km reached by the solar radiation in the atmosphere. The NO curves for May 1973 and May 1974 are very different even if their overall shapes are similar. The overall shape of the NO_2 curve differs much from the ones observed for NO.




FIG. 5. NUMBER DENSITY OF NO VS ALTITUDE OBTAINED BY THE INVERSION PROCESS. The differences between the values obtained in May 1973 and in May 1974 is outside the uncertainty limit discussed in the text. They relate not only to the abundances but also to the shape of the vertical distributions.




FIG. 6. NUMBER DENSITY OF NO₂ VS ALTITUDE OBTAINED BY THE INVERSION PROCESS. The results are in good agreement with the values reported by Ackerman and Muller (1972, 1973) and by Fontanella *et al.* (1974). They are lower than those obtained by Brewer *et al.* (1973).




FIG. 7. SUM OF THE NUMBER DENSITIES OF NO AND NO₂ DEDUCED FROM THE EXPERIMENT OF 13 MAY 1974, VS ALTITUDE.

The vertical distribution shown here corresponds to a decrease by a factor of 10 of the NO₂ volume mixing ratio from 34 to 20 km.

It is unambiguous that the NO densities derived from the May 1974 experiments are larger than those measured in May 1973 by means of the same instrumentation used under the same conditions. No precise deduction can however be drawn at present from this point, since the NO₂ over NO ratio depends on the ozone profile, which is not available to the authors for comparison at the two periods. For this reason it is not possible to attribute the observed difference to a variation in time of the NO₂ over NO ratio or to an increase of the sum of both components that could for instance, be related to transport processes or to other phenomena. It should however be mentioned that the tropopause altitude over the Gulf of Biscay was higher (≈ 13 km) in the case of the May 1973 experiment than in the case of the May 1974 one (≈ 9 km). The NO₂ vertical distribution is in rather good agreement with the values deduced by Ackerman and Muller in 1972 and by Fontanella *et al.* (1974). All of them seem to be within the limits of the possible natural variability.

The vertical distribution of the sum of NO and NO₂ shown in Fig. 7 exhibits a well marked maximum centered at about 27 km. The total amount in a vertical column between 20 and 37 km is equal to $(5.3 \pm 1.5) \times 10^{15}$ molecules per cm². This amount would correspond to an apparent volume mixing ratio above 20 km equal to $(4.9 \pm 1.5) \times 10^{-9}$.

TABLE 2. NO AND NO₂ RESULTS

Z(km)	NO May 1973 (10 ⁸ cm ⁻³)	NO, May 1974 (10 ⁸ cm ⁻³)	NO ₂ , May 1974 (10 ⁸ cm ⁻³)	NO + NO ₂ , May 1974 (10 ⁸ cm ⁻³)	NO ₂ /NO, May 1974
36	3.5 ± 2	11 ± 6	6 ± 2	17 ± 8	0.55
34	6.0 ± 3	16 ± 6	8 ± 2.5	24 ± 8.5	0.50
30	10 ± 4	20 ± 5	19 ± 4	39 ± 9	0.95
26	6 ± 3	21 ± 6	21 ± 4	42 ± 10	1.0
24	4 ± 2	18 ± 7	20 ± 4	38 ± 11	1.1
20	—	5.5 ± 3	13.5 ± 4	19 ± 7	3

TABLE 3. VOLUME MIXING RATIOS

Z(km)	NO + NO ₂ (M)	NO + NO ₂ + HNO ₃ * (M)	NO + NO ₂ + HNO ₃ † (M)
35	1.2×10^{-8}	1.3×10^{-8}	1.3×10^{-8}
34	1.3	1.4	1.4
30	1.1	1.3	1.4
26	6.4×10^{-9}	1.0	1.2
24	4.2	7.9×10^{-9}	1.0
20	1.3	3.3	1.0

* HNO₃ values taken from Lazrus and Gandrud (1974).

† HNO₃ values taken from Murcray *et al.* (1974) and Fontanella *et al.* (1974).

In fact this number can be a factor of 3 too small compared with the real maximum mixing ratio observed and listed in Table 3 where two sets of mixing ratios of the sum of three odd nitrogen molecules, NO₂, NO and HNO₃ are also given. The numbers for HNO₃ are from *in situ* measurements (Lazrus and Gandrud, 1974) in one set and from spectroscopic measurements (Fontanella *et al.*, 1974; Murcray *et al.*, 1974) in the other. In the first case the volume mixing ratio of the odd nitrogen molecules decreases by a factor of 4 from 30 to 20 km while in the second case it remains practically constant and equal to $(1.2 \pm 0.2) \times 10^8$ over the whole altitude range considered. Similar values, slightly varying or constant with height in the middle stratosphere have been derived or used in various models (Brasseur and Nicolet 1973; Dütch, 1973; MacElroy *et al.*, 1974; Crutzen, 1973; Isaksen, 1973). This emphasizes the need for resolving the discrepancy between the *in situ* and spectroscopic measurements of nitric acid that is considered in all the models above mentioned as the main loss agent for stratospheric oxides of nitrogen.

6. CONCLUSIONS

The variation with altitude of the concentration of NO_x appears to be firmly established by these first measurements. A considerable variation of NO at the same local time and season is observed in comparison with the results of those of a previous experiment one year earlier.

The main role played by nitric acid in the balance of odd nitrogen molecules in the lower stratosphere has been experimentally demonstrated. This should obviously be taken into account in models dedicated to realistic practical predictions of future trends of atmospheric constituents. Such models ought particularly to represent among many other aspects, the observed natural variations of various kinds from *ab initio* computations.

The measurements presented here can be considered as a first order attempt to determine the stratospheric abundance of odd nitrogen molecules. A more complete picture could be drawn if other constituents including at least O₃ and HNO₃ were also measured simultaneously with NO and NO₂.

It should be noted that possible uncertainties on the molecular absorption cross sections have not been taken into account in the absolute values given here and that more laboratory data are needed in this field.

Acknowledgements—This work has been partly supported by the French "Comité sur les Conséquences des Vols Stratosphériques" and by the Climatic Impact Assessment Program Office of the U.S. Department of Transportation through the Office of Naval Research under contract N00014-73-C-0076.

REFERENCES

- ACKERMAN, M., FRIMOUT, D., MULLER, C., NEVEJANS, D., FONTANELLA, J. C., GIRARD, A. and LOUISNARD, N. (1973). Stratospheric nitric oxide from infrared spectra. *Nature, Lond.* **245**, 205.
- ACKERMAN, M., FRIMOUT, D., FONTANELLA, J. C., GIRARD, A., GOBIN, R., GRAMONT, L. and LOUISNARD, N. (1973). Airborne and balloonborne spectroscopy for the study of atmospheric gas pollutants, ONERA T.P. 1309, paper presented at the *2nd Conf. Sensing of Environmental Pollutants*. Washington, 10–12 December 1973.
- ACKERMAN, M. and MULLER, C. (1972). Stratospheric nitrogen dioxide from infrared spectra. *Nature, Lond.* **240**, 300.
- ACKERMAN, M. and MULLER, C. (1973). Stratospheric methane and nitrogen dioxide from infrared spectra. *Pageoph* **106–108**, 1325.
- BRASSEUR, G. and NICOLET, M. (1973). Chemospheric processes of nitric oxide in the mesosphere and stratosphere. *Planet. Space Sci.* **21**, 939.
- BREWER, A. W., MCELROY, C. T. and KERR, J. B. (1973). Nitrogen dioxide concentrations in the atmosphere. *Nature, Lond.* **246**, 129.
- CIESLIK, S. and MULLER, C. (1973). Absorption raie par raie dans la bande fondamentale infrarouge du monoxyde d'azote. *Bull. Cl. Sci. Acad. Roy. Belg.* **59**, 869.
- CRUTZEN, P. (1973). A discussion of the chemistry of some minor constituents in the stratosphere and troposphere. *Pageoph* **106–108**, 1385.
- DANA, V. and FONTANELLA, J. C., (1973). Etude de l'absorption du dioxyde d'azote dans la région de 6, 18 μ . *Nouv. Rev. Optique* **4**, 237.
- DUTSCH, H. U. (1973). Recent developments in photochemistry of atmospheric ozone, *Pageoph* **106–108**, 1361.
- FONTANELLA, J. C., GIRARD, A., GRAMONT, L. and LOUISNARD, N. (1974). Vertical distribution of NO, NO₂ and HNO₃ as derived from stratospheric absorption spectra. *3rd CIAP Symp.*, Department of Transportation, Cambridge, Mass. and *Appl. Opt.* In press.
- GIRARD, A. (1963). Spectromètre à grilles. *Appl. Opt.* **2**, 79.
- GIRARD, A., FONTANELLA, J. C. and GRAMONT, L. (1973). Détections de l'oxyde azotique dans la stratosphère, *C.r. hebdomadaire Acad. Sci., Paris* **276**, 845–846.
- HARRIES, J. E. (1973). Measurements of some hydrogen–oxygen–nitrogen compounds in the stratosphere from Concorde 002. *Nature, Lond.* **241**, 515.
- HARTREE, D. R. (1958). *Numerical Analysis*. Clarendon, Oxford.
- HURLOCK, S. C., LAFFERTY, W. J. and RAO, K. N. (1974). Analysis of the ν_3 band of ¹⁴N¹⁶O₂, *J. molec. Spectrosc.* **50**, 246.
- ISAISEN, I. A. (1973). The production and distribution of nitrogen oxides in the lower stratosphere. *Pageoph.* **106–108**, 1438.
- KUNDE, V. G. (1967). Tables of theoretical line positions and intensities for the $v = 1$, $v = 2$ and $v = 3$ vibration–rotation bands of C¹²O¹⁶ and C¹³O¹⁶. NASA, Goddard Space Flight Center, X-622-67-248.
- LAZARUS, A. L. and GANDRUD, B. W. (1974). Distribution of stratospheric nitric acid vapor. *J. atmos. Sci.* **31**, 1102.
- LOEWENSTEIN, M., PADDOCK, J. P., POPPOFF, I. G. and SAVAGE, H. F. (1974). NO and O₃ measurements in the lower stratosphere from a U-2 aircraft. *Nature, Lond.* **249**, 817.
- MCCLATCHY, R. A., BENEDICT, W. S., CLOUGH, S. A., BURCH, D. E., CALFEE, R. F., FOX, K., ROTHMAN L. S. and GARING, J. S. (1973). AFCRL atmospheric absorption line parameters compilation. AFCRL-TR-73-0096, Environmental Research Papers No. 434, L. G. Hanscom Field, Bedford, Mass. 01730.
- MCELROY, M. B., WOFSEY, C., PENNER, J. E. and MCCONNELL, J. C. (1974). Atmospheric ozone: Possible impact of stratospheric aviation. *J. atmos. Sci.* **31**, 287.
- MURCRAY, D. G., GOLDMAN, A., CSOKE-POECKH, A., MURCRAY, F. H., WILLIAMS, W. J. and STOCKER, R. N. (1973). Nitric acid distribution in the stratosphere. *J. geophys. Res.* **78**, 7033.
- PATEL, C. K. N., BURKHART, E. G. and LAMBERT, C. A. (1974). Spectroscopic Measurements of stratospheric nitric oxide and water vapor. *Science*, **184**, 1173.
- RIDLEY, B. A., SCHIFF, H. I., SHAW, A. W., MEGILL, L. R., BATES, L., HOWLETT, C., LEVAUX, H. and ASHENFELTER, T. E. Measurements *in situ* of nitric oxide in the stratosphere between 17.4 km and 22.9 km. *Nature, Lond.* **245**, 310.
- RIDLEY, B. A., SCHIFF, H. I., SHAW, A. and MEGILL, L. R. (1974). *In-situ* measurements of NO in the stratosphere using chemiluminescence, Proc. 3rd Conf. Climatic Impact Assessment Prog. U.S. Department of Transportation, Cambridge, Mass., U.S.A.
- TOTH, R. A., FARMER, C. B., SCHINDLER, R. A., RAPER, O. F. and SHAPER, P. W. (1973). Detection of nitric oxide in the lower atmosphere. *Nature Phys. Sci.* **244**, 7.
- WHITING, E. E. (1968). An empirical approximation to the Voigt profile. *J. Quant. Spectr. Rad. Transfer* **8**, 1379.