

The Action of Chlorine on the Ozone Layer as Given by a Zonally Averaged Two-Dimensional Model

G. BRASSEUR¹⁾ and M. BERTIN²⁾

Abstract – In order to study the behavior of stratospheric minor constituents related to aeronomic processes and atmospheric transport in the meridional plane, a numerical two-dimensional model is established.

This model is applied to the study of chlorine compounds in the stratosphere. A special attention is devoted to the effect in the ozonosphere of an increase of ClX due to anthropogenic activities.

Key words: Chlorofluoromethanes; Ozone photochemistry; Two dimensional model.

1. Introduction

The stability of the atmospheric ozone layer has become an important problem since it has been recognized that the O₃ molecule could be destroyed by different stratospheric trace species [5], [11], [42]. In particular, it has been emphasized that the injection into the stratosphere of gases which are of anthropogenic origin, essentially nitrogen oxides [23] and halogen compounds [37], could lead to an ozone depletion and subsequently to an increase of the anaerobic UV radiation at ground level.

In order to study the potential effects of natural and artificial trace species on the ozonosphere, numerical models have been established extensively. Aeronomical models take into account the most important chemical and photochemical reactions and simulate the transport mechanisms using a very crude representation of the motions. The input data needed by these models are the chemical reaction rates and the absorption cross sections spectra, both obtained from laboratory investigations. Furthermore, the solar intensity spectrum in the UV range, the distribution of the mean temperature and the averaged transport parameters, which are supplied by observational studies, have to be specified. Finally a special attention has to be given to the boundary conditions. Output of these models are, besides the concentration profile of the minor constituents which are taken into account, several parameters,

¹⁾ Aspirant au Fonds National de la Recherche Scientifique, Institut d'Aéronomie Spatiale Bruxelles, Belgique.

²⁾ Société Nationale Industrielle Aérospatiale Paris, France.

such as the production and destruction rates, the particle flux and its divergence. It should be remembered that the results provided by mathematical models are always related to the input conditions and should be interpreted as such.

The purpose of this paper is to present very concisely some results obtained by the steady state version of our model which has already been described in detail elsewhere [8], [9]. This work deals mainly with the action of chlorine compounds and its relations with ozone. However, the model also takes into account the influence of hydrogen and nitrogen species. With respect to our previous work, the chemical reaction rates and the solar flux spectrum have been updated according to the most recent and most reliable laboratory determination and observation.

2. Short description of the model

The version of our 2-D model described hereafter considers steady state conditions. In this case, the continuity equation of the i th constituent is written

$$\nabla \cdot \phi_i = P_i - L_i \quad (1)$$

if ϕ_i is the particle flux, P_i and L_i respectively the production and the destruction rate. In order to relate the flux to the volume mixing ratio $f_i = n_i/n(M)$, where n_i is the concentration of the i th constituent and $n(M)$ the atmospheric concentration, a transport representation has to be specified. Since the detail of the atmospheric dynamics is not considered in this work, a simple parametrization of the large scale motions is adopted. According to the K theory, one writes the horizontal (y) and vertical (z) components of the flux as follows:

$$\phi_i^y = -n(M) \left[K_{yy} \frac{\partial f_i}{\partial y} + K_{yz} \frac{\partial f_i}{\partial z} \right] \quad (2a)$$

$$\phi_i^z = -n(M) \left[K_{zy} \frac{\partial f_i}{\partial y} + K_{zz} \frac{\partial f_i}{\partial z} \right] \quad (2b)$$

where K_{yy} , K_{yz} , K_{zy} and K_{zz} are the components of the tensor $\overline{\overline{K}}$ which is assumed to be symmetrical ($K_{yz} = K_{zy}$). These parameters have to be considered as purely phenomenological. Different authors [35], [18], [27], have derived stratospheric and tropospheric values of these coefficients. However, since the determination of the K 's is not unique, there is no general agreement about the values that should be adopted. Therefore, it is necessary to scale the coefficients so that the meridional distribution of species, which are sensitive to the transport, is close to the mean observation. In order to oversimplify the conditions, we have adopted a constant value of $K_{yy} = 10^{10} \text{ cm}^2 \text{ s}^{-1}$ and $K_{zz} = 10^4 \text{ cm}^2 \text{ s}^{-1}$ at all latitudes. K_{yz} has been adjusted in order to fit the observation of ozone in the meridional plane [26]. Figure 1 represents the latitudinal variation of K_{yz} resulting from the calibration.




Figure 1

Latitudinal profile of K_{yz} adopted below 25 km. Above this altitude, the values have been multiplied by the factor $\exp(-0.184(z - 25))$ where z is the altitude expressed in km. (The negative y -axis points to the winter-(Northern) hemisphere.)

The chemical and photochemical scheme adopted in the model results from the analysis of the aeronomical processes as made by Nicolet [29], [30], [31], [32]. Since a special attention is devoted to the chlorine chemistry, Fig. 2 represents a schematic graph indicating the most important reactions paths related to these compounds. It can be seen that odd chlorine (Cl , ClO , HCl and ClONO_2) is produced by dissociation of various halocarbons (CH_3Cl , CFCl_3 , CF_2Cl_2 , $\text{CCl}_4 \dots$) which are injected at ground level either by natural or by industrial processes, and disappears by wash-out mechanisms in the tropospheric clouds. Table 1 gives the complete list




Figure 2

Schematic graph of the most important chlorine reactions in the stratosphere.

Table I
The reactions and their rates used in this work.

Reaction	Rate constant (cm ³ s ⁻¹)	Reference
O(³ P) + O ₂ + M → O ₃ + M	$k_2 = 1.1 \times 10^{-34} e^{510/T} n(M)$	HUIE <i>et al.</i> [21]
O(³ P) + O ₃ → 2O ₂	$k_3 = 1.05 \times 10^{-11} e^{-2150/T}$	DAVIS <i>et al.</i> [12]
O(¹ D) + M → O(³ P) + M	$k_4 = 5 \times 10^{-11}$	NICOLET [29]
H + O ₂ + M → HO ₂ + M	$a_1 = 2.1 \times 10^{-32} e^{290/T} n(M)$	WONG and DAVIS [45]
H + O ₃ → OH + O ₂	$a_2 = 2.6 \times 10^{-11}$	PHILLIPS and SCHIFF [34]
OH + O → H + O ₂	$a_5 = 4 \times 10^{-11}$	NICOLET [32]
OH + O ₃ → HO ₂ + O ₂	$a_6 = 1.6 \times 10^{-12} e^{-1000/T}$	ANDERSON and KAUFMAN [3]
HO ₂ + O ₃ → OH + 2O ₂	$a_{6b} = 1.0 \times 10^{-13} e^{-1250/T}$	NICOLET [32]
HO ₂ + O → OH + O ₂	$a_7 = a_5$	Working value
OH + OH → H ₂ O + O	$a_{16} = 2 \times 10^{-12}$	WESTENBERG and DE HAAS [44]
		CLYNE and DOWN [10]
OH + HO ₂ → H ₂ O + O ₂	$a_{17} = 3 \times 10^{-11}$	HUDSON [20]
HO ₂ + NO → NO ₂ + OH	$a_{26} = 8 \times 10^{-12}$	HOWARD and EVENSON [19]
OH + CO → CO ₂ + H	$a_{36} = 1.25 \times 10^{-13}$	GREINER [17]
		DAVIS <i>et al.</i> [13]
NO ₂ + O(³ P) → NO + O ₂	$b_3 = 9.2 \times 10^{-12}$	DAVIS <i>et al.</i> [14]
NO + O ₃ → NO ₂ + O ₂	$b_4 = 2.1 \times 10^{-12} e^{-1450/T}$	HUDSON [20]
NO ₂ + O ₃ → NO ₃ + O ₂	$b_9 = 1.3 \times 10^{-13} e^{-2450/T}$	NICOLET [32]
NO ₂ + NO ₃ + M → N ₂ O ₅ + M	$b_{12} = 2.8 \times 10^{-30} n(M)$	BALAUCH <i>et al.</i> [6]
	$b_{12\infty} = 3.8 \times 10^{-12}$	Limiting value
NO ₂ + OH + M → HNO ₃ + M	$b_{22} = 2 \times 10^{-30} n(M)$	ANDERSON and KAUFMAN [2]
	$b_{22\infty} = 5 \times 10^{-12}$	Limiting value
HNO ₃ + OH → NO ₃ + H ₂ O	$b_{27} = 9 \times 10^{-14}$	SMITH and ZELLNER [41]
N ₂ O ₅ + M → NO ₂ + NO ₃	$b_{32} = 2.2 \times 10^{-5} e^{-9700/T}$	BAULCH <i>et al.</i> [6]
		NIKI [33]
N ₂ O + O(¹ D) → N ₂ + O ₂	$b_{38} = 6 \times 10^{-11}$	SCHIFF [39]
N ₂ O + O(¹ D) → 2NO	$b_{39} = 9 \times 10^{-11}$	SCHIFF [39]
H ₂ O + O(¹ D) → OH + OH	$a_{1a}^* = 2.1 \times 10^{-10}$	SCHIFF [39]
H ₂ + O(¹ D) → OH + H	$a_{1b}^* = 1.3 \times 10^{-10}$	SCHIFF [39]
CH ₄ + O(¹ D) → CH ₃ + OH	$a_{1c}^* = 1.3 \times 10^{-10}$	SCHIFF [39]
CH ₄ + OH → CH ₃ + H ₂ O	$c_2 = 3.5 \times 10^{-12} e^{-1800/T}$	GREINER [17]
		DAVIS <i>et al.</i> [13]
CH ₃ O ₂ + NO → CH ₃ O + NO ₂	$c_5 = 5 \times 10^{-13}$	Working value
CH ₃ Cl + OH → CH ₂ Cl + H ₂ O	$d_1 = 2.2 \times 10^{-12} e^{-1142/T}$	WATSON [43]
Cl + O ₃ → ClO + O ₂	$d_2 = 2.7 \times 10^{-11} e^{-257/T}$	WATSON [43]
ClO + O → Cl + O ₂	$d_3 = 7.7 \times 10^{-11} e^{-130/T}$	WATSON [43]
ClO + NO → Cl + NO ₂	$d_4 = 1.0 \times 10^{-11} e^{200/T}$	HUDSON [20]
Cl + CH ₄ → CH ₃ + HCl	$d_5 = 7.3 \times 10^{-12} e^{-1260/T}$	WATSON [43]
Cl + HO ₂ → O ₂ + HCl	$d_7 = 3 \times 10^{-11}$	WATSON [43]
HCl + OH → Cl + H ₂ O	$d_{11} = 3 \times 10^{-12} e^{-425/T}$	WATSON [43]
ClO + NO ₂ + M → ClONO ₂ + M	$d_{22} = \frac{3.3 \times 10^{-23} T^{-3.34} n(M)}{1 + 8.7 \times 10^{-9} T^{-0.6} [n(M)]^{0.5}}$	HUDSON [20]
ClONO ₂ + O → products	$d_{32} = 3.0 \times 10^{-12} e^{-808/T}$	WATSON [43]

Table 1.—*contd.*

Reaction	Photodissociation frequency (s^{-1})	Reference of absorption cross sections
$O_2 + hv \rightarrow O + O$	$J_{O_2}(\lambda < 242.4 \text{ nm})$	ACKERMAN [1]
$O_3 + hv \rightarrow O(^3P) + O_2$	$J_{O_3}(\lambda > 310 \text{ nm})$	ACKERMAN [1]
$O_3 + hv \rightarrow O(^1D) + O_2$	$J_{O_3}^*(\lambda < 310 \text{ nm})$	ACKERMAN [1]
$NO_2 + hv \rightarrow NO + O$	$J_{NO_2}(\lambda < 405 \text{ nm})$	BASS and LAUFER [4]
$HNO_3 + hv \rightarrow NO_2 + OH$	$J_{HNO_3}(\lambda < 320 \text{ nm})$	BIAUMÉ [7]
$N_2O + hv \rightarrow N_2 + O$	$J_{N_2O}(\lambda < 7420 \text{ nm})$	JOHNSTON and SELWYN [24]
$N_2O_5 + hv \rightarrow NO_2 + NO_3$	$J_{N_2O_5}(\lambda < 380 \text{ nm})$	GRAHAM [15]
$CH_3Cl + hv \rightarrow CH_3 + Cl$	J_{CH_3Cl}	ROBBINS [36]
$CFCl_3 + hv \rightarrow CFCl_2 + Cl$	$J_{CFCl_3}(\lambda < 226 \text{ nm})$	ROWLAND and MOLINA [37]
$CF_2Cl_2 + hv \rightarrow CF_2Cl + Cl$	$J_{CF_2Cl_2}(\lambda < 215 \text{ nm})$	ROWLAND and MOLINA [37]
$CCl_4 + hv \rightarrow CCl_3 + Cl$	J_{CCl_4}	ROWLAND and MOLINA [37]
$HCl + hv \rightarrow H + Cl$	$J_{HCl}(\lambda < 220 \text{ nm})$	INN [22]
$ClONO_2 + hv \rightarrow ClO + NO_2$	J_{ClONO_2}	ROWLAND <i>et al.</i> [38]

of the aeronomical reactions which are considered and the corresponding rate constants. The distribution of the mean temperature adopted in this work is the same as the one used by BRASSEUR [8]. References concerning the adopted absorption cross sections of these molecules are also given in Table 1. However, the attenuation of the UV flux due to molecular oxygen absorption in the Schumann–Runge bands is performed according to a numerical procedure established by KOCKARTS [25]. The spectrum of the solar radiation at the top of the atmosphere is taken from SIMON [40].

Finally, it should be added that a constant mixing ratio of 5×10^{-7} is assumed for H_2 , the corresponding value for CO being 5×10^{-8} in the stratosphere and 10^{-7} in the troposphere. In the case of H_2O , it is assumed at all latitudes that the volume mixing ratio decreases from 10^{-2} at ground level to 3×10^{-6} at 15 km and increases slightly to reach 5×10^{-6} at the stratopause. A working value of 1×10^{-12} is attributed to CH_3O_2 .

The model ranges from the North Pole to the South Pole and from the ground to the stratopause level. It attempts to represent for mean solar illumination conditions the distribution of minor constituents in the steady state. The seasonal conditions are chosen in such manner that the winter hemisphere corresponds to the Northern hemisphere. In order to treat the problem numerically, the derivatives are approximated by finite-differences. The distance between the gridpoints in the meridional plane has been fixed to 1 km vertically and 5 degrees latitude horizontally. An alternating direction method is applied to compute the solutions. However, in order to avoid divergent oscillations due to the stiffness of the equations system and related to the large differences between the lifetime of the various constituents, some of the chemical species are grouped so that isolated or quasi isolated systems of constituents are formed. This method is used to determine the global behavior of

$O_x = O_3 + O(^3P) + O(^1D)$, $HO_x = OH + HO_2$, $NO_y = NO + NO_2 + HNO_3 + ClONO_2$ and $CIX = Cl + ClO + HCl + ClONO_2$. The concentration of the individual constituents is derived by assuming equilibrium conditions between the various species belonging to a specified family. Other constituents, namely CH_4 , N_2O , CH_3Cl , $CFCl_3$, CF_2Cl_2 and CCl_4 , whose lifetime is rather long, are treated individually.

In order to specify the boundary conditions, one assumes no horizontal flux of any trace species at the North and the South Pole, so that the symmetry around the terrestrial axis is respected. At the ground and stratopause levels, a mixing ratio is specified as indicated in Table 2. The lower condition related to the chlorofluoromethanes is representative of a 1977 situation. Most of the conditions on the upper boundary are based on the results provided by 1-D models which extend to higher altitudes.

Table 2
Boundary conditions (volume mixing ratio).

	0 km	50 km
O_3	2.6×10^{-8}	photochemical equilibrium
N_2O	3×10^{-7}	1×10^{-10}
NO_y	1×10^{-9}	6×10^{-10}
CH_4	1.5×10^{-6}	2×10^{-8}
CH_3Cl	8×10^{-10}	2×10^{-14}
$CFCl_3$	1×10^{-10}	0
CF_2Cl_2	2×10^{-10}	5×10^{-13}
CCl_4	1×10^{-10}	0
CIX	1×10^{-9}	1.5×10^{-9}

3. Results and discussion

3.1. Chlorine in the atmosphere of 1977

The formation of odd chlorine in the stratosphere is related to the destruction of different chlorinated molecules which are released at ground level. The most important contribution [32] is due on the one hand to methylchlorides (CH_3Cl), whose origin is almost completely natural, and on the other hand to carbon tetrachloride (CCl_4), to trichlorofluoromethane ($CFCl_3$) and to dichlorofluoromethane (CF_2Cl_2) which are industrially produced. All these constituents whose mixing ratio is of the order of 10^{-10} to 10^{-9} in the troposphere [28] are injected into the stratosphere where they are photodissociated by UV radiation. Moreover, CH_3Cl reacts rather rapidly with hydroxyl radicals and is therefore partly destroyed in the troposphere. Figure 3 shows the vertical distribution of these species as given by our 2-D model at $30^\circ N$ latitude (winter conditions). It appears that the mixing ratio of those constituents decreases rapidly in the stratosphere so that, for example, CCl_4 and $CFCl_3$ have almost completely disappeared above the altitude of 35 km. Figure 4




Figure 3
Vertical distribution of CH_3Cl , CCl_4 , CFCl_3 and CF_2Cl_2 computed at 30°N for winter conditions.

represents the production rate of CIX in the meridional plane assuming that all the chlorine atoms present in the halocarbons are converted into odd chlorine. The production rate reaches values of the order of 10 to $1000 \text{ cm}^{-3} \text{ s}^{-1}$ in the troposphere due to the destruction of CH_3Cl by OH but these chlorine atoms are quickly washed out. In the stratosphere the production rate reaches a maximum of about $10 \text{ cm}^{-3} \text{ s}^{-1}$ in the equatorial region at about 22 km. This formation is due to the photodissociation of the different halocarbons.




Figure 4
Production rate of odd chlorine in the meridional plane assuming that all chlorine atoms present in the halocarbons are converted into CIX.

The vertical distribution at 30°N (winter) of Cl, ClO, ClONO₂ and HCl is shown on Fig. 5. According to the model, hydrogen chloride is the most abundant constituent at all altitudes but its concentration is very close to the concentration of chlorine nitrate between 20 and 30 km and is not far from the concentration of ClO between 40 and 46 km. Further theoretical and observational investigations are needed in order to establish the atmospheric value of the ClO/HCl ratio which is of particular importance to determine the effective loss of ozone by chlorine species. As a matter of




Figure 5
Vertical distribution of Cl, ClO, ClONO₂ and HCl computed at 30°N for winter conditions.




Figure 6
Distribution of the ClO mixing ratio in the meridional plane. The dashed line represents the altitude where the mixing ratio reaches its maximum value.

fact, the loss rate of ozone is proportional to the concentration of ClO and therefore the complete distribution of this molecule in the meridional plane is represented on Fig. 6.

3.2. Sensitivity of ozone to an increase of ClX

In order to investigate in a simplified way the effect of an increase of ClX related to anthropogenic activities, the amount of the actual chlorine resulting from the computation described in the previous section has been multiplied at all altitudes and all latitudes by 5 and 10 respectively. The corresponding relative ozone variation as a function of the altitude calculated for winter mid-latitude conditions is represented on Fig. 7. The graph shows that the largest ozone depletion occurs between 35 and 50 km with a maximum at about 43 km. The relative reduction is rather large (60 to 80 percent) at these heights but the absolute ozone concentration at these altitudes contributes only slightly to the total ozone column. On the other hand, the depletion is the smallest where the ozone concentration is the largest. According to our computation, one finds even a slight increase of the local concentration around 30 km. In the lower stratosphere, the reduction is of the order of 10–20 percent. Finally, the computation provides a global ozone depletion of 9 and 17 percent when the two different cases are considered. However, it should be noted, as shown on Fig. 8, that the relative reduction appears to be slightly latitude dependent.




Figure 7

Relative variation of the ozone concentration as a function of the altitude when the actual ClX amount is uniformly multiplied by 5 and 10 respectively.




Figure 8

Relative variation of the ozone column as a function of the latitude when the actual ClX amount is uniformly multiplied by 5.

These results should be considered as numerical experiments more than final answers to the problems of the stratosphere. Many uncertainties in the input data (aeronomical and transport parameters) remain and further investigations are needed before it will be possible to consider such results as representative of reality.

Acknowledgement

The authors would like to thank Professor M. Nicolet for his continual interest during the preparation of this work.

REFERENCES

- [1] ACKERMAN, M., *Ultraviolet solar radiation related to mesospheric processes*, in *Mesospheric Models and Related Experiments*, Reidel, Dordrecht, Holland (1971).
- [2] ANDERSON, J. G. and KAUFMAN, F. (1972), *Kinetics of the reaction* $\text{OH} + \text{NO}_2 + \text{M} \rightarrow \text{HNO}_3 + \text{M}$, *Chem. Phys. Letters* 16, 375.
- [3] ANDERSEN, J. G. and KAUFMAN, F. (1973), *Kinetics of the reaction* $\text{OH}(v=0) + \text{O}_3 \rightarrow \text{HO}_2 + \text{O}_2$, *Chem. Phys. Letters* 19, 483.
- [4] BASS, A. M. and LAUFER, A. H. (1973), *The extinction coefficient of NO₂ between 195 nm and 410 nm*, AIAA/AMS international Conference on the environmental impact of aerospace operations in the high atmosphere, Denver, Colorado.
- [5] BATES, D. R. and NICOLET, M. (1950), *The photochemistry of water vapor*, *J. Geophys. Res.* 55, 301.
- [6] BAULCH, D. L., DRYSDALE, D. D., HORNE, D. G. and LLOYD, A. C., *Evaluated kinetic data for high temperature reactions, Vol. 2, Homogeneous gas phase reactions of the H₂-N₂-O₂ system*, Butterworths, London (1973), 557 pp.
- [7] BIAUME, F. (1973), *Nitric acid vapour absorption cross-section spectrum and its dissociation in the stratosphere*, *J. Photochem.* 2, 139.

- [8] BRASSEUR, G. (1976), *L'action des oxydes d'azote sur l'ozone dans la stratosphère*, Thèse de doctorat, Université Libre de Bruxelles.
- [9] BRASSEUR, G. (1977), *Un modèle bidimensionnel du comportement de l'ozone dans la stratosphère*, Planet. Space. Sci. in press.
- [10] CLYNE, M. A. A. and DOWN, S. (1974), *Kinetic behaviour of OH $X^2\Pi$ and $A^2\Sigma^+$ using molecular resonance fluorescence spectroscopy*, J. Chem. Soc. Faraday Trans. 2, 70, 253.
- [11] CRUTZEN, P. J. (1970), *The influence of nitrogen oxides on the atmospheric ozone content*, Quart. J. Roy. Meteorol. Soc. 96, 320.
- [12] DAVIS, D. D., WONG, W. and LEPHARDT, J. (1973a), *A laser flash photolysis-resonance fluorescence kinetic study: Reaction of O(3P) with O₃*, Chem. Phys. Lett. 22, 273.
- [13] DAVIS, D. D., FISHER, S. and SCHIFF, R. (1973b), *Flash photolysis resonance fluorescence kinetics study, Temperature dependence of the reaction OH + CO \rightarrow CO₂ + H and OH + CH₄ \rightarrow H₂O + CH₃*, J. Chem. Phys. 61, 2213.
- [14] DAVIS, D. D., HERRON, J. H. and HUIE, R. E. (1973c), *Absolute rate constants for the reaction O(3P) + NO₂ \rightarrow NO + O₂ over the temperature range 230–339 °K*, J. Chem. Phys. 58, 530.
- [15] GRAHAM, R. A. (1975), *Photochemistry of NO₃ and the kinetics of the N₂O₅ – O₃ system*, Ph. D. Thesis, University of California, Berkeley.
- [16] GREENBERG, R. I. and HEICKLEN, J. (1970), *Reaction of O(1D) with N₂O*, Int. J. Chem. Phys. 2, 185.
- [17] GREINER, N. R. (1968), *Hydroxyl radical kinetics by kinetic spectroscopy*, J. Phys. Chem. 72, 406 (1968); J. Chem. Phys. 51, 5049 (1969); J. Chem. Phys. 53, 1070 (1970).
- [18] GUDIKSEN, P. H., FAIRHALL, A. W. and REED, R. J. (1968), *Roles of mean meridional circulation and eddy diffusion in the transport of trace substances in the lower stratosphere*, J. Geophys. Res. 73, 14, 4461.
- [19] HOWARD, C. J. and EVENSON, K. M. (1977), *Kinetics of the reaction of HO₂ radicals with NO*, NOAA, ERL-Aeronomy Lab., Boulder, Co, Preprint.
- [20] HUDSON, R. D. (1977), *Chlorofluoromethanes and the stratosphere*, NASA Reference publication 1010.
- [21] HUIE, R. E., HERRON, J. T. and DAVIS, D. D. (1972), *Absolute rate constants for the reaction O + O₂ + M \rightarrow O₃ + M in the temperature range 200–346°K*, J. Phys. Chem. 76, 2653.
- [22] INN, E. C. Y. (1975), *Absorption coefficient of HCl in the region 1400 to 2200 Å*, J. Atm. Sci. 32, 2375.
- [23] JOHNSTON, H. S. (1971), *Reduction of stratospheric ozone by nitrogen oxide catalysts from supersonic transport exhaust*, Science 173, 517.
- [24] JOHNSTON, H. S. and SELWYN, G. S. (1975), *New cross sections for the near ultraviolet radiation by nitrous oxide (N₂O)*, Geophys. Res. Lett. 2, 549.
- [25] KOCKARTS, G. (1976), *Absorption and photodissociation in the Schumann–Runge bands of molecular oxygen in the terrestrial atmosphere*, Planet Space Sci. 24, 589.
- [26] LONDON, J. (1967), *The average distribution and time variation of ozone in the stratosphere and mesosphere*, Space Res. 7, 172.
- [27] LUTHER, F. M. (1973), *Monthly mean values of eddy diffusion coefficients in the lower stratosphere*, AIAA paper No. 73–498.
- [28] NATIONAL ACADEMY OF SCIENCES, *Halocarbons effects on stratospheric ozone*, Washington D.C. (1976).
- [29] NICOLET, M. (1970), *Aeronomic reactions of hydrogen and ozone*, Aeronomica Acta A–79 (1970) and in *Mesospheric Models and Related Experiments*, Reidel Publ. Co., Dordrecht, (1971), pp. 1–51.
- [30] NICOLET, M. (1972), *Aeronomic chemistry of the stratosphere*, Planet. Space Sci. 20, 1671.
- [31] NICOLET, M. (1974), *An overview of aeronomic processes in the stratosphere and mesosphere*, Canad. J. Chem. 52, 1381.
- [32] NICOLET, M. (1975), *Stratospheric ozone: An introduction to its study*, Rev. Geophys. Space Phys. 13, 593.
- [33] NIKI, H. (1974), *Reaction kinetics involving O and N compounds*, Can. J. Chem. 52, 1397.
- [34] PHILLIPS, L. F. and SCHIFF, H. I. (1962), *Reactions in the atomic nitrogen–ozone systems*, J. Chem. Phys. 36, 1509.
- [35] REED, R. J. and GERMAN, K. E. (1965), *A contribution to the problem of stratospheric diffusion by large scale mixing*, Month. Weather Rev. 93, 313.
- [36] ROBBINS, D. E. (1976), *Photodissociation of methyl chloride and methyl bromide in the atmosphere*, Geophys. Res. Lett. 3, 213.

- [37] ROWLAND, F. S. and MOLINA, M. J. (1975), *Chlorofluoromethanes in the environment*, Rev. Geophys. Space Phys. 13, 1.
- [38] ROWLAND, F. S., SPENCER, J. E. and MOLINA, M. J. (1976), *The stratospheric chemistry of chlorine nitrate*, ClONO₂, J. Phys. Chem. 80, 2711.
- [39] SCHIFF, H. (1977), *The role of nitrogen oxides in stratospheric chemistry*, Paper presented at the IAGA/IAMAP Joint Assembly, Seattle.
- [40] SIMON, P. C. (1977), Private communication.
- [41] SMITH, I. W. M. and ZELLNER, R. (1975), *Rate measurements of reactions of OH by resonance absorption*, 4, *Reactions of OH with NH₃ and HNO₃*, Int. J. Chem. Kinet. 7, 341.
- [42] STOLARSKI, R. S. and CICERONE, R. J. (1974), *Stratospheric chlorine: a possible sink for ozone*, Can. J. Chem. 52, 1610.
- [43] WATSON, R. T. (1977), *Review of halogen kinetics*, Position paper for NASA CFM Assessment.
- [44] WESTERNBERG, A. A. and DE HAAS, N. (1973), *Rate of reaction of OH + OH → H₂O + O*, J. Chem. Phys. 58, 4046.
- [45] WONG, W. and DAVIS, D. D. (1974), *A flash photolysis-resonance fluorescence study of the reaction of atomic hydrogen with molecular oxygen: H + O₂ + M → HO₂ + M*, Intern. J. Chem. Kinetics 6, 401.
- [46] YOUNG, R. A., BLACK, G. and SLANGER, T. G. (1968), *Reaction and desactivation of O(¹D)*, J. Chem. Phys. 49, 4758.

(Received 21st November 1977)
