

Density and temperatures of Venus upper atmosphere measured by stellar occultations with SPICAV/Venus Express

A. Piccialli (1), F. Montmessin (1), J. L. Bertaux (1), E. Marcq (1), A. Fedorova (2), O. Korablev (2), A. Mahieux (3), A. C. Vandaele (3)

(1) LATMOS-UVSQ, Guyancourt, France, (2) Space Research Institute (IKI), 84/32 Profsoyuznaya, 117810 Moscow, Russia, (3) Planetary Aeronomy, Belgian Institute for Space Aeronomy, Brussels, Belgium (arianna.piccialli@latmos.ipsl.fr)

Abstract

We will present a preliminary study of the Venus atmospheric structure between 90 and 140 km of altitude using SPICAV-UV/VEx stellar occultation observations. More than 500 density and temperature profiles were retrieved at various latitudes and local time. The upper atmosphere of Venus shows a very large temporal variability; variations both from orbit-to-orbit and with local time are observed.

1. Introduction

Venus upper atmosphere (80-140 km altitude) is one of the most interesting regions on the planet. It is a transition region characterized by a complex dynamics: strong retrograde zonal winds dominate the lower mesosphere while a solar-antisolar circulation driven by a day-to-night temperature gradient can be observed in the upper mesosphere/lower thermosphere. CO₂ density and temperature profiles of Venus upper atmosphere have been measured from both ground-based [1, 2] and spacecraft missions [3, 4]. The thermal structure of Venus mesosphere shows a significant latitudinal variability probably driven by the dynamics. Atmospheric temperatures reach a minimum value of ≈ 170 K at 90–100 km altitudes on the dayside of the planet. More recently, a layer of warm air has been detected at altitudes of 90-120 km on the nightside both by SPICAV/SOIR [5, 6] and by ground-based [7] observations.

2. Observations

The SPICAV (Spectroscopy for the investigation of the characteristics of the atmosphere of Venus) [8] instrument has been operating on board the ESA orbiting platform Venus Express since 2006. It is a

remote sensing spectrometer covering distinct spectral regions in ultraviolet (118–320 nm) and near-infrared (650–1700 nm). In the stellar occultation mode the UV sensor is particularly well suited to measure the vertical profiles of CO₂, temperature, SO₂, SO, clouds and aerosols of the middle and upper atmosphere of Venus. A very large dataset has been collected by SPICAV-UV that consists of more than 500 stellar occultation profiles performed at all latitudes and seasons during nighttime (Fig. 1).




Figure 1: distribution of SPICAV-UV stellar occultations in local time and latitude.

3. Results

CO₂ density and temperature profiles were derived by SPICAV-UV stellar occultations covering the altitude range 90 – 140 km (Fig. 2).




Figure 2: Example of (top) CO₂ density and (bottom) temperature profiles for orbit 95A17.

Fig. 3 displays the local time – altitude cross section of atmospheric temperature. A permanent warm area appears distinctly at the mesopause at about 90-100 km of altitude. Temperature then decreases with altitude reaching a minimum value around 125 km altitude in good agreement with previous observations [6]. Spatial and temporal changes in the thermal structure are analysed.




Figure 3: local time – altitude cross section of atmospheric temperature (K). All SPICAV-UV stellar occultations have been used.

Acknowledgements

AP acknowledges funding from the European Union Seventh Framework Programme (FP7/2007-2013) under grant agreement No. 246556. The research program was supported by the Belgian Federal Science Policy Office and the European Space Agency (PRODEX program). The research was performed as part of the “Interuniversity Attraction Poles” programme financed by the Belgian government (Planet TOPERS).

References

- [1] Clancy, R. T. et al. (2012) *Icarus*, 217, 779-793.
- [2] Lellouch, E. et al. (1991) *Icarus*, 110,315-339.
- [3] Keating, G. et al. (1985) *JGR*, 85, 7941-7956.
- [4] Zasova, L. V. et al. (2006) *Cosmic Res.*, 44, 364-383.
- [5] Bertaux, J. L. et al. (2007) *Nature*, 450,646-649.
- [6] Mahieux, A. et al. (2012) *JGR*, 117, E07001.
- [7] Rengel, M. et al. (2008) *PSS*, 56, 1368–1384.
- [8] Bertaux, J. L. et al. (2007) *PSS*, 55, 1673-1700.