

Scheer Marc

21/01/69

INSTITUT D'AERONOMIE SPATIALE DE BELGIQUE

3, avenue Circulaire, UCCLE - BRUXELLES 18

AERONOMICA ACTA

A - N° 20 - 1963

The ionospheric conditions

by M. NICOLET and W. SWIDER, Jr.

BELGISCH INSTITUUT VOOR RUIMTE-AERONOMIE

3, Ringlaan, UKKEL - BRUSSEL 18

FOREWORD

"The Ionospheric Conditions" has been presented at the COSPAR'S meeting in Warsaw (June 1963) and has been used for a seminar for COPERS in Alpbach (Austria). It will be published in Planetary and Space Science.

M. NICOLET.

AVANT-PROPOS

"The Ionospheric Conditions" a été présenté au Symposium de COSPAR en juin à Varsovie et a été l'objet d'un séminaire au cours d'été de COPERS à Alpbach (Autriche). Il sera publié dans Planetary and Space Science.

M. NICOLET.

VOORWOORD

"The ~~Ionospheric~~ Conditions" werd ten gehore gebracht op het Symposium van COSPAR in juni 1963 te Warchau en is het voorwerp geweest van een seminaire op een zomercursus ingericht door COPERS te Alpbach (Oostenrijk). De tekst zal verschijnen in "Planetary and Space Science".

M. NICOLET.

VORWORT

"The Ionospheric Conditions" wurde während der COSPAR'S Sitzung in Warschau (Juni 1963) vorgestellt und wurde für ein Seminar während des Sommerkursus von COPERS in Alpbach (Österreich) gebraucht. Diese Arbeit wird in Planetary and Space Science herausgegeben werden.

M. NICOLET.

THE IONOSPHERIC CONDITIONS

by

Marcel NICOLET and William SWIDER, Jr.

Résumé.

Une analyse générale des conditions physiques dans l'ionosphère a été effectuée en considérant les diverses réactions auxquelles sont soumis les divers ions atomiques et moléculaires.

Tout d'abord, la détermination de la production et de la recombinaison ioniques indique que la distribution spectrale de l'émission solaire ultraviolette doit être mesurée avec plus de précision et que la recombinaison dissociative doit être fixée en fonction de la température.

Ensuite, le comportement de l'ionosphère dépend de l'ensemble des processus influençant les ions atomiques O^+ et N^+ , c'est-à-dire des réactions avec les molécules N_2 , O_2 et NO . Les ions moléculaires N_2^+ , O_2^+ et NO^+ sont liés entre eux par suite de l'existence de réactions d'interchange entre ions et atomes neutres intervenant en même temps que les échanges de charges.

En outre, la distribution verticale des ions est discutée dans le cadre d'une détermination de l'importance relative des processus au sein des diverses régions ionosphériques. L'ion moléculaire d'azote N_2^+ disparaît par le transfert de charge tandis que l'ion NO^+ ne peut disparaître que par l'intermédiaire du processus de recombinaison radiative.

Au surplus, la production des ions atomiques d'oxygène dépend non seulement de sa production par photo-ionisation directe de l'atome d'oxygène mais aussi de la production de l'ion moléculaire d'azote. Quant au rapport des concentrations des ions NO^+ et O_2^+ , il dépend dans l'ionosphère inférieure de réactions particulières.

Enfin, les ions He^+ et H^+ sont liés à un système de production et de réactions qui les situe à haute altitude où les conditions de diffusion sont essentielles.

INTRODUCTION

The problem of the interpretation of the physics and chemistry of the ionosphere is in a state of confusion. The increasing knowledge of the solar spectrum in the far ultraviolet has, apparently, not as yet lead to a consistent picture of the photoionization processes in the ionosphere. While Hinteregger and Watanabe ^[1] have determined a certain distribution of the number of photons available at the top of the ionosphere, Ivanov-Kholodny ^[2] has adopted another distribution. The data of Table I between 900A and 30A show that the difference is important. It is clear that for $\lambda < 800$ A, the number of photons used by

Table I - Number of photons ($\text{cm}^{-2} \text{sec}^{-1}$) available at the top of the earth's atmosphere according to Hinteregger-Watanabe (H-W) and Ivanov-Kholodny (I-K)

$\Delta\lambda$	910-850	850-800	800-700	700-600	600-500	500-400A	
(H-W)	9.5×10^9	2.5×10^9	5.0×10^9	4.7×10^9	5.9×10^9	2.4×10^9	photons
(I-K)	5.0×10^9	2.1×10^9	4.4×10^{10}	3.2×10^{10}	4.9×10^{10}	3.0×10^{10}	photons
<hr/>							
$\Delta\lambda$	400-300	300-230	230-170	170-110	110-80	80-60	60-30 A
(H-W)	7.2×10^9	3.1×10^9	3.3×10^9	3.5×10^8	2.4×10^8	1.8×10^8	1.6×10^8 photons
(I-K)	9.6×10^{10}	5.3×10^{10}	2.4×10^{10}	5.7×10^9	2.24×10^9	1.1×10^9	8.4×10^8 photons

Ivanov-Kholodny is ten times that obtained by Hinteregger and Watanabe. Nevertheless, Norton, Van Zandt and Denison ^[3], using the spectral distribution given by Hall, Damon and Hinteregger ^[4] and Ivanov-

Kholodny^[2], claim that they have satisfactorily interpreted the ionospheric structure.

It is difficult to understand how it is possible to make such divergent interpretations of optional ionospheric behavior; even allowing for variation of the far ultraviolet spectrum with solar activity. The analysis of the solar spectrum in the region of $\lambda > 800$ by Detwiler, Garrett, Purcell and Tousey^[5] and by Zirin, Hall and Hinteregger^[6] show consistent results. In the spectral region of $\lambda < 800$, Table I indicates a difference of a factor of 10 which will affect the ionization of all thermospheric constituents, namely N_2 , O_2 and O . Such a variation cannot exist in the whole ultra-violet spectrum of wavelengths $\lambda \leq 800$ A, even for the maximum-to-minimum variation of the solar cycle.

On the other hand, Hinteregger^[7], and Norton et al.^[3] consider an atmospheric model with an atomic oxygen concentration which is not less than four times the molecular oxygen concentration at 110 km and not less than the molecular nitrogen concentration at 130 km. Denison and Van Zandt^[3] claim that Nicolet's model^[8] is inconsistent with Hall's et al. data^[4]. This deduction by Hinteregger^[7] and Norton et al.^[3] is based essentially on the behavior of two solar lines, namely $L \gamma$ at 972 A and Si III at 1206 A in the wing of Lyman- α . However, such a deduction can be modified if the laboratory measurements of the absorption cross-sections are not adapted to the atmospheric absorption. As an example, the vertical distribution of solar photons for Lyman- γ at 972.537 A for an absorption cross-section of the order of 1×10^{-16} cm² using Nicolet's model fits the vertical distribution deduced by Hall et al.^[4]. An N_2 band with a head at 972.2 A, which is shaded toward longer wavelengths, has more than 10 rotational lines in a narrow

spectral range with $J_P(15) = 973.98 \text{ \AA}$. The absorption cross-section depends on the distribution of the rotational lines between 972 \AA and 974 \AA and a high resolution spectrum is needed to determine the exact absorption cross-section of $\lambda \ 972.537 \text{ \AA}$. It cannot be claimed that the cross-section is $1 \times 10^{-16} \text{ cm}^2$ or $3 \times 10^{-16} \text{ cm}^2$ without a special analysis of laboratory and atmosphere conditions. It is evident that an observation of solar Ly free from atmospheric absorption is required to make a correct measurement of the exact photon flux at the top of the earth's atmosphere.

Hinteregger's data [7] show several anomalies. The ratio $n(N_2)/n(O_2)$ decreases from a factor of 10 at 120 km to 4 and 2.5 at 130 km and 150 km; respectively. The ratio $n(O)/n(O_2)$ decreases from about 15 between 110 and 120 km to 10 at 130 km and 140 km. The normal behavior for such ratios is to increase with height. Furthermore, it should be noted that Hinteregger gives the ratio $n(N_2)/n(O)$ equal to only 0.4 at 130 km while the ratio $n(O)/n(O_2)$ reaches at least 15 between 110 km and 120 km. On the other hand, Denison and Van Zandt [3] using the same observational data adopt the following ratios: $n(O)/n(O_2) = 4$ at 110 km, 9 at 120 km and about 15 at 130 km where $n(O)$ almost reaches the concentration of $n(N_2)$.

Ratios such as $n(O)/n(N_2) \geq 1$ at 130 km and $n(O)/n(O_2) \geq 10$ at 125 km do not agree with mass spectrometric measurements made in the USA (cf Townsend [9], Schaefer [9a]) and in the USSR (cf Pokhunkov [10]). For example, an observation made by Schaefer shows that $n(O)/n(O_2) = 1$ near 118 km and not more than 3 at 130 km.

The description of the daytime ionospheric regions as given by

Norton et al. [3] or by Ivanov-Kholodny [2] cannot be used for nighttime conditions without the addition of another process. Antonova and Ivanov-Kholodny [11] have introduced the ionization effect of electrons. The flux in the ionosphere should be at least of the order of $1 \text{ erg cm}^{-2} \text{ sec}^{-1}$ for electrons of energy between 100 eV and 10 keV, i. e. at least of the order of $5 \times 10^9 \text{ electrons cm}^{-2} \text{ sec}^{-1}$ for an energy of 200 eV. In other words, Ivanov-Kholodny, after introducing a high ionization rate for daytime conditions, is obliged to consider a high recombination rate to obtain the observed electron concentrations. When such a recombination rate is used for nighttime conditions it would destroy the F_2 layer and a corpuscular flux is postulated to maintain a nocturnal ionosphere. However, such an electron flux would excite the night airglow spectrum in such a way that auroral emissions should occur; in particular the first negative system, and the first positive system of nitrogen. Since such emissions are not observed it can be concluded, as Galperin [12] has shown, that the normal corpuscular flux is very small. It is practically negligible compared with the ultraviolet flux, which is certainly greater than $1 \text{ erg cm}^{-2} \text{ sec}^{-1}$, and is insignificant compared with the 15 ergs claimed by Ivanov-Kholodny as the normal ultraviolet flux of $\lambda < 900 \text{ A}$.

A recent analysis shows how a corpuscular flux must be considered. At mid-latitudes over North America O'Brien [13] has measured, at 1000 km, a flux of precipitated electrons of 10^3 and $10^4 \text{ cm}^{-2} \text{ sec}^{-1}$ for energies $\geq 40 \text{ keV}$. His discussion on the airglow excitation indicates that the normal fluxes are between 10^{-2} and $10^{-1} \text{ ergs cm}^{-2} \text{ sec}^{-1}$ for 1 keV electrons. It is possible that some anomaly could be detected as shown

recently by Gledhill and Van Rooyen [14] using O'Brien's data [13].





The airglow data suggest various limits for the electron flux. An emission of the green line of atomic oxygen above 300 km, equal to the intensity of that of the normal airglow and due to the excitation of electrons of about 10 eV energy, would correspond to a flux of 10^{10} electrons $\text{cm}^{-2} \text{sec}^{-1}$. An electron flux in the energy range 50-500 eV, i. e. less than 1 keV, cannot be more than 10^7 electrons $\text{cm}^{-2} \text{sec}^{-1}$ since it would lead to such an excitation of N_2^+ bands which are not observed in the airglow. Furthermore, the second positive system of nitrogen, which is likewise not observed in the airglow, requires the same limiting conditions for the electron flux. Therefore, the ionization rate coefficient for the corpuscular radiation cannot be greater than 10^{-9}sec^{-1} i. e. is always less than 10^{-2} the photoionization rate coefficient. Since there is no possibility of finding a permanent electron flux in the nighttime ionosphere greater than $0.1 \text{ erg cm}^{-2} \text{sec}^{-1}$ it must be concluded that the total ionization production by corpuscular radiation cannot attain in the whole ionosphere (for normal conditions outside of the auroral zone) one hundredth of that produced by photo-ionization.

Finally, the number of atmospheric models used to explain the upper-atmosphere behavior has added to the confusion and leads to the other difficulties. Consequently, before trying to clarify the general problem, an attempt is made to show the complexity of the problem of ionospheric reactions and the difficulty of deducing a correct explanation of the ionosphere without an exact knowledge of the various physical parameters involved.


I. Positive Ion Reactions

Starting from the observational knowledge of the ionosphere, the following positive ions must be considered: H^+ , He^+ , N^+ , N_2^+ , O^+ , O_2^+ , NO^+ , Mg^+ , Ca^+ , ... etc. Further, it should be noted that metallic ions are not present in the whole ionosphere, H^+ and He^+ occur in the upper part of the ionosphere above the F_2 peak, O_2^+ , NO^+ and O^+ are the essential ionic constituents of the D, E and F regions and N_2^+ and N^+ are minor ions with maximum concentrations in the F region. Thus, O^+ , N^+ , NO^+ , O_2^+ and N_2^+ must be considered as the basic ions which are involved in the various reactions.

The predominant ion in the F region, O^+ , is subject to the following ion-atom interchange reactions:



The following charge transfer process must be added to reactions (1) to (4):



Since there is no practical distinction between (2) and (5), process (5) will be ignored in the analysis. Similarly, a charge transfer process between O^+ and NO is equivalent to reaction (3).

The rate coefficients of the ion-atom interchange reactions (1) to (4) are not well known. The rate of (2) has been measured by Dickinson



and Sayers^[15], Langstroth and Hasted^[16] and Fite, Rutherford, Snow and Van Lint^[17]. The following values have been obtained:

$$\begin{aligned}\gamma_2 &\geq 1 \times 10^{-10} \text{ cm}^3 \text{ sec}^{-1} [17] \\ \gamma_2 &= (2.5 \pm 0.4) \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1} [15] \\ \gamma_2 &= (1.8 \pm 0.2) \times 10^{-12} \text{ cm}^3 \text{ sec}^{-1} [16]\end{aligned}$$

From ionospheric observations, values between $10^{-9} \text{ cm}^3 \text{ sec}^{-1}$ and $10^{-13} \text{ cm}^3 \text{ sec}^{-1}$ have been used to explain recombination and ionic composition; Hertzberg^[18], $\gamma = 10^{-9} \text{ cm}^3 \text{ sec}^{-1}$; Danilov^[19], $\gamma = 10^{-10} \text{ cm}^3 \text{ sec}^{-1}$; Norton, Van Zandt and Denison^[3], $\gamma_2 = 5 \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$; Bates and Nicolet^[20], $\gamma = 10^{-13} \text{ cm}^3 \text{ sec}^{-1}$.





Langstroth and Hasted^[16] have measured, for reaction (1), $\gamma_1 = (4.7 \pm 0.5) \times 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$. But before this recent experimental determination a high value of the order of $10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ was given by Potter^[16a] and a rate coefficient of the order of $10^{-10} \text{ cm}^3 \text{ sec}^{-1}$ has been used in the study of the ionosphere by Krassovsky^[16b]. The recent experimental analysis of Talrose, Markin and Larin^[16c] seems, however, to agree with a lower rate coefficient i.e. $\gamma_1 \leq 6.75 \times 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$. Various values have been used in the analysis of the ionospheric observations such as $10^{-9} \text{ cm}^3 \text{ sec}^{-1}$ by Hertzberg^[18], $10^{-10} \text{ cm}^3 \text{ sec}^{-1}$ by Danilov^[19b], $1 \times 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$ by Norton, Van Zandt and Denison^[3] and of the order of $10^{-13} \text{ cm}^3 \text{ sec}^{-1}$ by Bates and Nicolet^[20].

Several reactions involving O_2^+ lead to NO^+ , and particularly,







This last reaction, according to Fite et al.^[17], is very slow compared with the charge transfer process or ion-atom interchange reaction $O_2^+ + NO \rightarrow O_2 + NO^+$. However, in the atmosphere, since $n(N_2) \gg n(NO)$, (7) cannot be neglected even if the rate coefficient of the transfer process with NO is large. Reaction (6) cannot be rejected since its rate coefficient, γ_6 , may be much greater than γ_7 and also because in the 100 km region, where $n(N)$ is the most important fraction of the $n(N_2)$, its effect could modify the ratio $n^+(O_2)/n^+(O)$.

The processes affecting N_2^+ involve charge transfer processes and ion-atom interchange reactions:



Finally, ion-atom interchange reactions in which N^+ and ON^+ are involved must be considered by taking into account that the lower state of ON^+ is the excited state $^3\Pi$ of the normal ion NO^+ .



Reactions which may occur in the ionosphere require that the excitation energy $E(^3\Pi)$ is less than the energy balance. Thus, reaction

(14) must be rejected since $E(^3\Pi)$ is more than 3eV, for example $E(^3\Pi) = 4.6\text{eV}^{[21]}$. However, Norton, Van Zandt and Denison^[3] have considered reaction (14) as very important in the ionosphere since they have adopted for the rate coefficient $\gamma_{14} = 2 \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$ i.e., more than the value of γ_1 for which they take $1 \times 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$. This is based on the erroneous assumption that the reaction (14) does not involve an excited electronic level of NO^+ . It should be pointed out that the charge transfer ($\text{N}^+ + \text{NO} \rightarrow \text{NO}^+ + \text{N}$) is equivalent in the ion produced to reaction (13), but is different from reaction (11).

The reaction energies derived in (1) to (15) correspond to the following dissociation and ionization potentials.

Dissociation of ionized molecules:

O_2^+ , 6.66 eV	N_2^+ , 8.73 eV	NO^+ , 10.88 eV
--------------------------	--------------------------	--------------------------

Dissociation of neutral molecules:

O_2 , 5.12 eV	N_2 , 9.76 eV	NO , 6.51 eV
------------------------	------------------------	-----------------------

Ionization of neutral molecules:






O_2 , 12.08 eV	N_2 , 15.58 eV	NO , 9.25 eV
-------------------------	-------------------------	-----------------------

Ionization of neutral atoms:


O, 13.62 eV	N, 14.55 eV	H, 13.60 eV
-------------	-------------	-------------

II. Photo-ionization Processes

The reactions (1) to (13) must be used with the photo-ionization and recombination rates to write equations for electron and ion concentrations. The following ionization processes are considered:



The photo-ionization of NO is essentially due to the Lyman- α radiation at 1216 A:



The absorption cross-sections and atmospheric densities are known with sufficient precision as far as the order of magnitude is concerned. Consequently it is possible to state that the following ionization processes occur:

(i) D region

- (a) ionization of nitric oxide by Lyman- α , since O_2 , N_2 and O cannot be photo-ionized by λ 1216.7 A. Other constituents such as Na ($\lambda < 2412 \text{ A}$), Ca ($\lambda < 2071 \text{ A}$), CH_3 ($\lambda < 1260 \text{ A}$), C ($\lambda < 1100 \text{ A}$) can be photo-ionized by radiations which are unable to ionize O_2 .
- (b) ionization by X rays of $\lambda < 10 \text{ A}$, i.e. with cross-sections less than 10^{-19} cm^2 . It is known from rocket and satellite measurements that the Lyman- α flux is between 3 and 6 ergs $\text{cm}^{-2} \text{ sec}^{-1}$ while the X-ray flux is extremely variable with solar conditions. From 2-8 A the minimum-to-maximum variation is a factor of several hundred (see a recent review

by Friedman [22]).

- (c) ionization by cosmic rays which is effective in the lower D region where Lyman- α cannot penetrate and where X rays of $\lambda < 2 \text{ A}$ are not important; particularly during solar quiet conditions.

An exact analysis of the ionization problem in the D region requires the determination of the separate effects of the three ionizing sources. In fact, it is necessary to compare the ionizing effect of the stable sources Lyman- α and cosmic rays as compared with the extremely varying source represented by X-rays. Such an analysis has been made by Nicolet and Aikin [23] who considered various X-ray fluxes according to data obtained by the Naval Research Laboratory over an interval of several years with Lyman- α and cosmic ray ionization processes introduced by Nicolet [23a, 23b].

The ionization rate coefficient of cosmic rays is between (Φ being the geomagnetic latitude)

$$I(\Phi \geq 0^\circ) \geq 10^{-18} \text{ sec}^{-1} \text{ and } I(\Phi \leq 50^\circ) \leq 10^{-17} \text{ sec}^{-1},$$

i. e. with the following values: $\Phi = 0^\circ$, $1.25 \times 10^{-18} \text{ sec}^{-1}$, $\Phi = 30^\circ$, $2 \times 10^{-18} \text{ sec}^{-1}$, $\Phi = 40^\circ$, $4.5 \times 10^{-18} \text{ sec}^{-1}$, $\Phi = 50^\circ$, $1 \times 10^{-17} \text{ sec}^{-1}$ and $\Phi = 60^\circ$, $1.25 \times 10^{-17} \text{ sec}^{-1}$.

It is clear that X-rays of $\lambda \leq 2 \text{ A}$ emitted during completely quiet conditions cannot lead to an ionization rate coefficient of the order of $10^{-18} \text{ sec}^{-1}$. On the other hand, since the ionization rate coefficient of nitric oxide by Lyman- α reaches values of the order of

$$I(\text{NO}) = 5 \times 10^{-7} \text{ sec}^{-1},$$

a small ratio $n(\text{NO})/n(\text{O}_2) = 10^{-10}$ leads to an electron production of the same order as the X-ray production under quiet solar conditions. Thus, the disturbed solar conditions, and particularly solar flare conditions, produce a D region due to X-ray radiation.

(ii) E Region

- (a) ionization of molecular oxygen by ultraviolet radiation of $\lambda > 910 \text{ \AA}$, particularly by monochromatic radiations such as Lyman- β at 1025 \AA , i. e. with absorption cross-sections between 10^{-19} cm^{-2} and 10^{-18} cm^{-2} . In addition, ionization of atomic oxygen by radiation of $\lambda < 910 \text{ \AA}$ and, particularly, by the chromospheric Lyman continuum which can penetrate into the E layer via the windows of molecular nitrogen absorption. The variation with the solar cycle of these ultraviolet radiations should be less than a factor of two.
- (b) ionization by X-rays of $\lambda > 10 \text{ \AA}$ and particularly by the radiation in the spectral range $30 < \lambda < 100 \text{ \AA}$. The minimum-to-maximum solar cycle variation is approximately sevenfold [22].
- (c) ionization of meteor atoms with low ionization potentials, introduced by Nicolet [24], to explain nocturnal conditions in the E layer, and recently observed by Istomin [25].

An exact analysis of the ionization production problem in the E region still requires the determination of the separate effects of the three ionizing sources. The stable source represented by the ultraviolet radiations will have its maximum ionization effect under quiet solar

conditions and, particularly, during the minimum of the solar cycle. The X-ray action will vary with solar activity and will become predominant during disturbed solar conditions.

Considering that the ionization rate coefficient of O_2 at zero optical depth is not less than 10^{-8} sec^{-1} , it is clear that an X-ray flux varying from 0.15 to 1 $\text{erg cm}^{-2} \text{ sec}^{-1}$ for an average wavelength of 50 Å leads to an electron production from less than to greater than the ultraviolet production of O_2^+ . To be consistent with the solar data the ratio of the electron production by ultraviolet radiations (O_2^+ , O^+) and by X-rays (N_2^+ , O_2^+ , O^+) must decrease with increasing solar activity, but at no time may one ionization process be neglected in comparison with the other. In any case, the X-ray action must explain the behavior of the E layer during the entire solar cycle.

(iii) F region

- (a) ionization of molecular nitrogen for $\lambda < 796 \text{ Å}$ with absorption cross-section greater than 10^{-17} cm^2 .
- (b) ionization of atomic oxygen for $\lambda < 796 \text{ Å}$ subject to the absorption of molecular nitrogen.
- (c) ionization of atomic oxygen for $\lambda > 800 \text{ Å}$ with different absorption cross-sections for its different ionization potentials at 910 Å, 732 Å and 665 Å.

An exact analysis of the ionization problem in the F region requires a simultaneous knowledge of the energy of solar emissions and of the absorption cross-sections of O, N_2 and O_2 . In particular, the penetration of monochromatic solar radiations between the E and F_1 peaks must be known in order to determine the exact behavior of the electron


production.

Considering that variations of solar activity will affect chromospheric and coronal lines differently, it is difficult to predict the exact height-distribution of the electron production; for example, the variation of line intensity in the spectrum should be more important for the shortest wavelengths than for the Lyman continuum. However, it appears that N_2 and O ionizations will be subject to almost the same variations, and that the minimum-to-maximum variation during a solar cycle should not be greater than fourfold.

III. Electron Recombination



The various processes (1) to (15) represent the most important ionospheric reactions in which nitrogen and oxygen positive ions are involved. The processes (16) to (21) correspond to photo-ionization of these constituents. It should be kept in mind that similar ionization processes produce metallic ions and also light ions such as He^+ and H^+ . Consequently, the first recombination process to be considered is radiative recombination, such as X^+ (or XY^+) + $e \rightarrow X$ (or XY) + $h\nu$. Since the recombination coefficient is always between 1×10^{-12} and $5 \times 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$ in the ionosphere (see Bates and Dalgarno^[26]), the radiative recombination is neglected compared with other processes affecting ions. Three-body electronic or ionic recombination processes are also neglected since they cannot play a role in the normal ionosphere.

The essential recombination process in the ionosphere is Bates' process of neutralization due to the dissociation of a molecular ion as follows:



(the * indicating possible excitation). Dissociative recombination occurs (see Bates and Dalgarno [26]) as "a result of a radiationless transition to some state of the molecule in which the constituents move apart and gain kinetic energy under the action of their mutual repulsion so that the neutralization is rendered permanent by virtue of the Franck-Condon principle". It is clear from the theoretical analysis that it is practically impossible to predict the temperature variation of the dissociative recombination coefficient, α . Furthermore, since the dissociation depends on the presence of a repulsive curve, it may be assumed that the absolute values of α should be different for different molecules. In other words, if values of α between 10^{-6} and $10^{-9} \text{ cm}^3 \text{ sec}^{-1}$ and variations with temperatures such as $T^{-1} \pm 0.5$ are assumed, they do not contradict theoretical considerations. The problem is quite complicated since there is no simple law establishing the variation of the repulsive curves of molecules. Consequently, the absolute value of α and its dependence on T must be found experimentally.

The principal dissociative recombination processes which occur in the ionosphere are:



Various experimental determinations of the recombination coefficients of these three ions exist, but it is difficult to adopt a correct value for

ionospheric purposes. The determinations of $\alpha(\text{N}_2)$ at room temperature, after 1958, give $4 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ (Faire and Champion^[27]) and $(5.9 \pm 1) \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ (Kasner, Rogers and Biondi^[28]), corrected more recently to $2 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$. At high temperatures, there is an indication given by Sayers^[29] for 3200° K , $\alpha(\text{N}_2) = 1.1 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$, and by Mentzoni, Montgomery and Row^[30] for 725° K , $\alpha(\text{N}_2) = 6.1 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$. Consequently, $\alpha(\text{N}_2)$ is not known from laboratory measurements with sufficient precision for ionospheric purposes. A rough conclusion should be that $\alpha(\text{N}_2) > 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ for the lower ionosphere.

No value can be obtained for the upper ionosphere since the temperature variation is not known. The recent determination of $\alpha(\text{O}_2)$ by Kasner, Rogers and Biondi^[28] leading to $3.8 \pm 1 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ has been corrected to $1.7 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ (private communication) indicating that there seems to be no practical difference between $\alpha(\text{O}_2)$ and $\alpha(\text{N}_2)$ at room temperature. At high temperatures the value obtained by Sayers^[29], $\alpha(\text{O}_2) = 4 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ at 2500° K , seems to indicate a rapid decrease of $\alpha(\text{O}_2)$ with temperature.

Laboratory data on $\alpha(\text{NO})$ follow a pattern similar to those for $\alpha(\text{N}_2)$ and $\alpha(\text{O}_2)$. Extremely high values such as $\alpha(\text{NO}) = 1.3 \times 10^{-6}$ were recently obtained by Gunton and Inn^[31], as were obtained in the 1950's (before 1958) for N_2 , and values between 4×10^{-7} and $2 \times 10^{-6} \text{ cm}^3 \text{ sec}^{-1}$ by Doering and Mahan^[32]. For a high temperature of about 5000° K , Lin^[33] gave $\alpha(\text{NO}) = 10^{-9} \text{ cm}^3 \text{ sec}^{-1}$. Syverson et al.^[34] published a value of $\alpha(\text{NO}) = 1.3 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ at 3000° K . Thus, the temperature dependence of the coefficient is unknown.

In such circumstances it is clear that it is not possible to argue

for or against any value adopted for ionospheric conditions. This points to the complexity of the determination of an ionospheric recombination coefficient depending on the recombination of several ions. Nicolet and Aikin^[23] adopted $\alpha(N_2) = 5 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$, $\alpha(O_2) = 3 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ and $\alpha(NO) = 3 \times 10^{-9} \text{ cm}^3 \text{ sec}^{-1}$ for their D region analysis to show how the ionization is affected when various constituents are involved. However, while extreme values for N_2 and NO still remain a possibility, any value of the order of $10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ would not change the situation since ionic recombination and negative ions are involved.

On the other hand, Whitten and Poppoff^[35] have attempted to interpret the behavior of the D region during ionospheric disturbances by an ionospheric coefficient between $4 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ and $4 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$. However, they claimed^[36] that the D-region ionization can be interpreted, even during quiescent solar conditions by, X-ray radiations. Such an evaluation of the ionospheric behavior in the D region indicates that an uncertainty of a factor of ten in the recombination coefficient still remains a possibility when X-ray energies are not properly related to solar conditions. Thus, any value of α from 10^{-7} to $10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ used for D region conditions cannot be contradicted. In fact Ivanov-Kholodny^[2] adopted $\alpha = 3 \times 10^{-7} \sqrt{300/T} \text{ cm}^3 \text{ sec}^{-1}$ for all recombination coefficients. Norton et al.^[3] have adopted different values: $\alpha(N_2) = 4 \times 10^{-7} (300/T) \text{ cm}^3 \text{ sec}^{-1}$, $\alpha(O_2) = 2 \times 10^{-7} (300/T) \text{ cm}^3 \text{ sec}^{-1}$ and $\alpha(NO) = 6.7 \times 10^{-8} (300/T) \text{ cm}^3 \text{ sec}^{-1}$, i.e. a variation of a factor of 6 between $\alpha(N_2)$ and $\alpha(NO)$. The values adopted by Ivanov-Kholodny and Norton et al. demonstrate that one may

estimate the absolute value of a recombination and its variation with temperature. Obviously temperature dependence is introduced in order to follow ionospheric observations, but a coefficient varying with $T^{-1} \pm 0.5$ leads to very different values in the F layer.

An analysis of ionospheric observations, such as diurnal variations or solar eclipse measurements, lead to different conclusions with recombination coefficients as low as $10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ and greater than $10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ (see Ratcliffe [37]). However, it is clear from eclipse determinations that a high value for the ionospheric recombination coefficient must be taken since there is solar emission during totality.

The nighttime conditions should give a value of the recombination coefficient in the E and F_1 regions. If values of the order of $2 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$, such as reported by Titheridge [38], are accepted for nighttime conditions, they should represent the recombination coefficient of the predominant ion. However, it is not yet clear what the exact vertical distribution of electrons is in the nighttime E and F_1 layers. If there is a peak in the E layer due to meteor ions, the analysis of the normal recombination is perturbed. If the electron concentration at ionosphere sunset is $n_{e,0}$ its value $(n_e)_t$ after t seconds is given by

$$(n_e)_t = \frac{1}{\alpha t + 1/n_{e,0}}$$


Thus, after 4×10^4 seconds, the following values are obtained:

$\alpha (\text{cm}^3 \text{ sec}^{-1})$	$= 2.5 \times 10^{-7}$	2.5×10^{-8}	2.5×10^{-9}	2.5×10^{-10}
$(n_e)_t (\text{cm}^{-3})$	$\leq 10^2$	10^3	10^4	10^5
$n_{e,0} (\text{cm}^{-3})$	$> 10^3$	10^4	10^5	10^6

Electron concentrations of the order of 10^3 cm^{-3} below the F_2 region require an electron recombination coefficient of the order of $2.5 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ for a predominant ion such as NO^+ , or the presence of meteor ions with radiative recombination coefficients. A value of α reaching $10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ cannot result in more than 2.5×10^2 electrons at the end of the night. Nevertheless, types of E_S ionization could modify the analysis since the ion behavior is different according to the nature of the ions. For example, meteor ions and nitric oxide are not subject to resonance charge transfer as is the case for O_2^+ , O^+ and N_2^+ , and their redistributions in the presence of the earth's magnetic field are different. There is also the problem of the recombination coefficient changing with time due to its dependence on electronic and ionic temperatures. In any event, it must be remembered that the electron-ion collision frequency decreases with temperature ($\propto T^{-3/2}$) and that the normal tendency of the temperature dependence of α_D , therefore, should be, in a complicated way, a decrease with increasing temperatures.

IV. Negative Ions

The reactions in which negative ions are involved are important in the D region and must be added to the preceding reactions. The mutual neutralization reaction between positive and negative molecular ions such as




will add to the dissociative recombination $\alpha(\text{XY})$ an additional term $\alpha_i \lambda$ where λ is the ratio of negative ion and electron concentrations

for equilibrium conditions. In the same way




could be considered in a certain ionospheric region. The rate coefficients of reactions such as (25) and (26) are not accurately known (cf. Dalgarno^[39]); values between 10^{-7} and $10^{-8} \text{ cm}^3 \text{ sec}^{-1}$ are possible.

The introduction of negative ions in the D region certainly depends on the three-body attachment



for which Chanin, Phelps and Biondi^[40] have obtained an experimental value of the order of $2.3 \times 10^{-30} \text{ cm}^6 \text{ sec}^{-1}$ for low atmospheric temperatures. For atomic oxygen, radiative attachment is an essential process



for which an attachment coefficient of the order of $1.3 \times 10^{-15} \text{ cm}^3 \text{ sec}^{-1}$ has been deduced from the photodetachment cross-section measured by Branscomb, Burch, Smith and Geltman^[41].

For such negative ions, the photodetachment by solar radiation leads to a photodetachment

$$d(O^-) = 1.4 \text{ sec}^{-1} \quad (29)$$

using experimental data obtained by Branscomb et al.^[41] while $d(O_2^-)$ is about 10 times smaller than (29). Photodetachment is the principal loss process for negative ions in a sunlit atmosphere while mutual neutralization is important in the nighttime D region. However,

it should be noted that ions such as O^+ do not exist at low altitudes due to ion-atom interchange reactions. Furthermore, O_2^+ is involved in an ion-atom interchange reaction leading to NO^+ . Thus, the positive ions in (25) must be the molecular ions O_2^+ and NO^+ .

In the lower nighttime D region, atomic oxygen is excluded from collisional detachment of a negative ion because it disappears via ozone formation. Negative ions such as OH^- , O_3^- and NO_2^- , for which higher affinities exist, depend on the behavior of O_2^- and may be considered as playing a role during twilight conditions. In other words, their concentrations are related to $n(O_2^-)$ for nighttime conditions through reactions which generally involve atoms. For example, $n(O_3^-)/n(O_2^-)$ depends on $n(O)$, $n(NO_2^-)/n(O_2^-)$ depends on $n(N)$; i. e. on atoms which are not important in the lower D region after sunset.

However, charge transfer processes should be considered. For example, the OH ion under the same conditions should increase since a charge transfer such as $OH + O_2^- \rightarrow OH^- + O_2$ will not be compensated for collisional detachment with atomic hydrogen, H , which is unimportant where OH can play a role. Finally, all reactions leading to NO_2^- , an ion having a high affinity (greater than 3.8 eV according to Curran^[42]), should be studied.

For other conditions, the processes of associative detachment discussed by Dalgarno^[39], i. e.



are certainly the most important processes for collisional detachment.

Thus, even for nighttime conditions, negative ions are important only in the D region, and it is difficult to find adequate processes to introduce negative ions in the E layer.

V. Analysis of Ionospheric Reactions


1 - General conditions

All processes (1) to (24) (after neglecting the effect of negative ions) must be considered in order to determine the essential reactions in the ionosphere. The scheme of Figure 1 gives a general idea of the relationships between the various processes. From this, it is clear that only NO^+ can be considered as disappearing by dissociative recombination alone, O_2^+ and N_2^+ concentrations are affected by ion-atom interchange reactions (γ_6 and γ_7) and by charge transfer processes (γ_8 and γ_9); respectively. O^+ is transformed into molecular ions by ion-atom interchange reactions (γ_1 to γ_4). Its transformation into NO^+ is due to reactions with N_2 and NO (γ_1 and γ_3) and into O_2^+ by O_2 and NO (γ_2 and γ_4). However, O^+ production is increased by the charge transfer process between atomic oxygen and molecular nitrogen ions (γ_8). The ratio $n^+(\text{O}_2)/n^+(\text{NO})$ is not independent of reactions leading to O_2^+ or NO^+ since O_2^+ may transform into NO^+ by ion-interchange reactions (γ_7 and γ_6).

2 - Equilibrium conditions

In order to determine the relative importance of reactions (1) to (24), ionization equations are first considered for steady state conditions. They may be written as follows

I O N O S P H E R I C R E A C T I O N S



IONOSPHERIC REACTIONS. PHOTOIONIZATION WITH RATE COEFFICIENT I , DISSOCIATIVE RECOMBINATION WITH COEFFICIENT α , CHARGE TRANSFER OR ION-ATOM INTERCHANGE WITH RATE COEFFICIENT γ .

FIGURE 1

$$n^+(N_2) = \frac{n(N_2) I(N_2) + n^+(N) \gamma_{11} n(NO)}{\alpha(N_2)n_e + \gamma_8 n(O) + \gamma_9 n(O_2) + \gamma_{10} n(N)} \quad (32)$$

$$n^+(N) = \frac{n(N_2) I_2(N) + n(N) [I(N) + \gamma_{10} n^+(N_2)]}{\gamma_{12} n(O_2) + [\gamma_{11} + \gamma_{13}] n(NO)} \quad (33)$$

$$n^+(O) = \frac{n(O) [I(O) + \gamma_8 n^+(N_2)]}{\gamma_1 n(N_2) + \gamma_2 n(O_2) + [\gamma_3 + \gamma_4] n(NO)} \quad (34)$$

$$n^+(O_2) = \frac{n(O_2) [I(O_2) + \gamma_2 n^+(O) + \gamma_9 n^+(N_2)] + n(NO) \gamma_4 n^+(O)}{\alpha(O_2) n_e + \gamma_6 n(N) + \gamma_7 n(N_2)} \quad (35)$$

$$n^+(NO) = \frac{1}{\alpha(NO)n_e} \left\{ n^+(O) [\gamma_1 n(N_2) + \gamma_3 n(NO)] + n^+(O_2) \right. \quad (36)$$

$$\left. [\gamma_6 n(N) + \gamma_7 n(N_2)] + n^+(N) [\gamma_{12} n(O_2) + \gamma_{13} n(NO)] + n(NO) I(NO) \right\}$$

In the denominator of (32) the process $N_2^+ + NO \rightarrow NO^+ + N_2$ has not been introduced, since the concentration of nitric oxide is only a fraction of that of molecular oxygen. According to Nicolet's formula [43], adapted to recent rate coefficients,

$$n(NO) = n(O_2) \times 10^{-1} e^{-3000/T} \quad (37)$$

which shows that $n(NO)$ is always a small fraction of $n(O_2)$. With diffusion conditions for nitric oxide, $n(NO)$ and $n(O_2)$ decrease with height at almost at the same rate since $M(NO) = 30$ and $M(O_2) = 32$, and the ratio $n(NO)/n(O_2)$ almost corresponds to the ratio fixed by photo-equilibrium conditions at the lower boundary of the diffusion region.

On the other hand, reactions (14) and (15), which would lead directly to NO^+ , are not considered since (8) and (9) are more effective processes. However, a special remark must be made here on the problem of nitric oxide, since Danilov claimed that the relation (37) cannot be accepted. His arguments are based on the assumption that molecular oxygen is strongly dissociated at 150-160 km altitude [19c]. In fact, he assumed the following equilibrium:

$$\alpha_D n^+(\text{O}_2) n_e = \gamma n^+(\text{O}) n(\text{O})$$

where the rate coefficient α_D of the dissociative recombination is $10^{-6} \text{ cm}^3 \text{ sec}^{-1}$, while the rate coefficient γ for a radiative association of O_2^+ is about $5 \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$. Such a value is much too great for a radiative association; by a factor of $10^5 - 10^6$. With $5 \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$, we should obtain

$$-\frac{1}{n^+(\text{O})} \frac{dn^+(\text{O})}{dt} \geq 5 \times 10^{-11} n(\text{O})$$

and we should find that, for $n(\text{O}) \geq 10^7 \text{ cm}^{-3}$, a time shorter than 10^4 sec would reduce the sunset concentration to less than 1 percent.

For example, no nighttime ionosphere at 300 km - 350 km, where $n(\text{O}) = 5 \times 10^8 \text{ cm}^{-3}$, can exist after a few hundred seconds. In order

to maintain an observed ion concentration, a nighttime production of more 1000 ions $\text{cm}^{-3} \text{ sec}^{-1}$ should be required at all latitudes.

The same remark must be applied to the dissociation of nitrogen.

Danilov introduces also the reaction



with a rate coefficient of the order of $3 \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$ which is


too great by a factor of several hundred thousands.

In fact, the problem must be considered as follows. The extent to which molecular oxygen is dissociated depends on the importance of the loss processes (2), in addition to the normal photodissociation $J(O_2)$ compared with the vertical transport, w , by diffusion. A continuity equation such as

$$\frac{dn(O_2)}{dt} = \frac{d[n(O_2)w]}{dt} - n(O_2) [J(O_2) + \gamma_2 n^+(O)] ,$$

since $J(O_2) > 10^{-6} \text{ sec}^{-1}$ with the use of $\gamma_2 \leq 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$, shows that the diffusion distribution of molecular oxygen represents, in a first approximation, the vertical distribution of that molecule. The disappearance of molecular oxygen in the F_2 region as claimed by Danilov should lead to such a strong departure from a diffusive equilibrium distribution, that an upward vertical flow of O_2 molecules is immediately supported by diffusion. It can easily reach more than $10^{11} \text{ molecules cm}^{-2} \text{ sec}^{-1}$ at 150 km and remains greater than $10^{10} \text{ molecules cm}^{-2} \text{ sec}^{-1}$ at 250 km. Thus, there is no difficulty in relating the dissociation of nitrogen to the vertical distribution of molecular oxygen.

In fact, atomic nitrogen reacts with O_2 according to the process




for which the rate coefficient b_1 can be written

$$b_1 = 1.5 \times 10^{-13} T^{1/2} e^{-3000/T}$$

showing the strong effect of the temperature. An important loss

process for atomic nitrogen occurs in the F_1 region where the temperature is high and where there is still a sufficient number of O_2 molecules. Furthermore, nitric oxide reacts easily with N according to



for which a rate coefficient b_2 may be written

$$b_2 = 1.5 \times 10^{-12} T^{1/2},$$

with practically no activation energy. In chemical equilibrium, (37) represents, therefore, the exact conditions. Thus, the mechanism of atomic nitrogen recombination and that of the formation of O_2^+ ions proposed by Danilov cannot be accepted since they are based on the assumption that radiative associations have rate coefficients of the order of $10^{-11} \text{ cm}^3 \text{ sec}^{-1}$.

It follows from the preceding remarks that an adequate approximation for (32) to (36) is introduced by neglecting reactions (3), (4), (11) and (13) in that $\gamma n(NO) \ll \gamma n(O_2)$ or $\gamma n(N_2)$.

In all the ionization equations (32) to (36) there is competition between reactions depending on the neutral constituent concentrations. A simple way to investigate the various effects is to introduce numerical factors having all possible values between 0 and 1. They are:

(a) for loss of atomic oxygen ions

$$A = \frac{\gamma_2 n(O_2)}{\gamma_1 n(N_2) + \gamma_2 n(O_2)} ; \quad 1 - A = \frac{\gamma_1 n(N_2)}{\gamma_1 n(N_2) + \gamma_2 n(O_2)} \quad (38)$$

(b) for production of atomic oxygen ions,

$$B = \frac{\gamma_8 n(O)}{\gamma_8 n(O) + \gamma_9 n(O_2)} ; \quad 1 - B = \frac{\gamma_9 n(O_2)}{\gamma_8 n(O) + \gamma_9 n(O_2)} \quad (39)$$

(c) for loss of molecular nitrogen ions,

$$C = \frac{\alpha(N_2) n_e}{\alpha(N_2) n_e + \gamma_8 n(O) + \gamma_9 n(O_2)} ; \quad 1 - C = \frac{\gamma_8 n(O_2) + \gamma_9 n(O_2)}{\alpha(N_2) n_e + \gamma_8 n(O_2) + \gamma_9 n(O_2)} \quad (40)$$

(d) for loss of molecular oxygen ions,


$$D = \frac{\alpha(O_2) n_e}{\alpha(O_2) n_e + \gamma_6 n(N) + \gamma_7 n(N_2)} ; \quad 1 - D = \frac{\gamma_6 n(N) + \gamma_7 n(N_2)}{\alpha(O_2) n_e + \gamma_6 n(N) + \gamma_7 n(N_2)} \quad (41)$$

3. - Molecular nitrogen ion


The ion N_2^+ is a minor constituent in the ionosphere [44] , [46] , [47] , even if its production by ultraviolet radiation $n(N_2)I(N_2) \equiv q(N_2)$ is important. Equation (32) becomes, after making the approximations justified before,

$$n^+(N_2) = \frac{q(N_2)}{\alpha(N_2) n_e + \gamma_8 n(O) + \gamma_9 n(O_2)} \quad (42)$$

showing that various loss processes dominate depending on the altitude involved. It must be pointed out here that Danilov^[48a] has introduced a concept which cannot be accepted. Instead of using the charge transfer or ion-atom interchange (10)



he considered the opposite endothermic reaction



with a rate coefficient of the order of $10^{-11} \text{ cm}^3 \text{ sec}^{-1}$.

It is not permissible to ignore an exothermic reaction when the opposite endothermic reaction is used, even if the ratio of the N_2^+ and N^+ concentrations that Danilov [48a] obtained agrees with the observed values. The reasoning is incorrect and the numerical values used are wrong. The relative concentration of N_2^+ in the E layer is very small, but its bands are observed in the twilight airglow, even though the nighttime intensity is very low ($\leq 5 \times 10^6 \text{ photons cm}^{-2} \text{ sec}^{-1}$) during quiet magnetic conditions. The twilight observations must be explained by a fluorescence effect of solar radiation on N_2^+ ions which are present in the F_2 region. The production of N_2^+ is due to solar radiation of $\lambda < 796 \text{ \AA}$ which are absorbed in the F_1 region; X rays of $\lambda < 10 \text{ \AA}$ are absorbed in the E layer; X rays shorter than 10 \AA are absorbed in the D region, and cosmic rays ionize N_2 in the lower part of the D region (Nicolet and Aikin [23]). There is, therefore, a production of N_2^+ ions in the whole ionosphere, i. e. where the electron concentration varies from 10^2 cm^{-3} to 10^6 cm^{-3} . In other words, the first term in the denominator of (42) corresponding to the dissociative recombination is not always the most important term; N_2^+ can be transformed, before dissociative recombination, by charge transfer process (8) and (9) involving O^+ and O_2^+ ; respectively. Thus the ionization equation (42) becomes, by using (40),

$$n^+(N_2) = \frac{C q(N_2)}{\alpha(N_2) n_e} \quad (43a)$$

or

$$n^+(N_2) = \frac{[1 - C] q(N_2)}{\gamma_8 n(O) + \gamma_9 n(O_2)} \quad (43b)$$

in which $0 \leq C \leq 1$ increases with height. With electron concentrations of the order of 10^6 cm^{-3} , i. e. in the F_2 layer, the lifetime of N_2^+ is very short since it depends on an exponential term approximately as follows

$$n^+(N_2) = n_o^+(N_2) e^{-\alpha(N_2)n_e t} + \frac{q(N_2)}{\alpha(N_2)n_e} [1 - e^{-\alpha(N_2)n_e t}] \quad (44)$$

leading to

$$\tau(N_2^+) = 1/\alpha(N_2) n_e \quad (45)$$


namely a few hundred seconds as a maximum lifetime.

Thus, photoequilibrium represents the ionization conditions for N_2^+ . Nevertheless, an exact analysis of aeronomic data is difficult since a determination of the factor C in equation (43a) is needed. An aeronomic determination cannot lead to a precise knowledge of rate coefficients for which experimental data are still lacking. Since the theoretical analysis shows that there must be a transformation of N_2^+ into O_2^+ in the D region, and into O^+ up to the F region, it is shown under which conditions charge transfer processes (8) and (9) should dominate.

Let us take $5 \times 10^{-7} \text{ cm}^3 \text{ sec}^{-1}$ as a working (maximum) value for $\alpha(N_2)$. In the D region (60-80 km), $\gamma_9 n(O_2) \geq \alpha_{N_2} n_e$ when

$$\begin{array}{lll}
 n(\text{O}_2) & = 10^{15} & = 10^{14} \text{ cm}^{-3} \\
 \alpha(\text{N}_2)n_e & = 5 \times 10^{-5} (*) & = 5 \times 10^{-4} \text{ sec}^{-1} \\
 \gamma_9 & \geq 10^{-19} & \geq 10^{-17} \text{ cm}^3 \text{ sec}^{-1}
 \end{array}$$

Such values of γ_9 , corresponding to charge transfer cross-sections less than 10^{-23} cm^2 show that there is no problem in considering the transformation of N_2^+ into O_2^+ in the D region. In addition, the following charge transfer process can be adopted for argon



and thus the ionospheric conditions resulting from the ionization of the principal atmospheric constituents by cosmic rays and by X rays lead to molecular oxygen ions.

As far as the E region (100 - 120 km) is concerned, the same analysis for O_2 and O gives

$$\begin{array}{lll}
 n(\text{O}_2), n(\text{O}) & = 10^{12} & = 10^{11} \text{ cm}^{-3} \\
 \alpha(\text{N}_2) n_e & = 5 \times 10^{-2} \text{ sec}^{-1} & \\
 \gamma_9, \gamma_8 & \geq 10^{-13} & \geq 10^{-12} \text{ cm}^3 \text{ sec}^{-1}
 \end{array}$$

The small amount of N_2^+ in the E region cannot be justified if only dissociative recombination is involved. It can be justified if the rate coefficients γ_8 or γ_9 reach values of the order of $10^{-12} \text{ cm}^3 \text{ sec}^{-1}$, i.e. a charge transfer cross-section of the order of $5 \times 10^{-18} \text{ cm}^2$

(*) Ionic recombination is neglected here; its effect is small at 80 km.

corresponding to 1/1000 times a resonance charge transfer. Without increasing the preceding value of γ_8 in the F region, a decrease with temperature of the dissociative recombination coefficient must be introduced. With, for example, $\alpha(N_2) = 5 \times 10^{-8} \text{ cm}^3 \text{ sec}^{-1}$, the following conditions are obtained for the F region:

Altitude	150	300	500 km
$n(O)$	$= 10^{10}$	$= 10^9$	$= 10^8 \text{ cm}^{-3}$
$\alpha(N_2)n_e$	$= 10^{-2}$	$= 5 \times 10^{-2}$	$= 10^{-2} \text{ sec}^{-1}$
γ_8	$\geq 10^{-12}$	$\geq 10^{-11}$	$\geq 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$

Thus, if the charge transfer process (8) has a rate coefficient of the order of $10^{-12} \text{ cm}^3 \text{ sec}^{-1}$ an additional production of atomic oxygen ions results from the general photo-ionization of molecular nitrogen. Bearing in mind that the production rate of N_2^+ is important in the F_1 layer, it is clear that the production of O^+ should be increased as compared with its direct production by photoionization of atomic oxygen. The transformation of N_2^+ into O^+ is a process different from an ion-atom interchange reaction leading to NO^+ indicating that the role played by N_2^+ in the E and F_1 layers does not correspond to that which Norton, Van Zandt and Denison [3] assumed.

From the present analysis of reactions involving N_2^+ ions, it appears that N_2^+ concentrations are not directly related to the nature of other ions but to the presence of electrons and the concentrations of the neutral constituents. In addition, N_2^+ may always be considered as essentially being in photo-ionization equilibrium in all ionospheric layers.


4 - Atomic nitrogen ion

Atomic nitrogen is a minor constituent^[43] and its ion, N^+ , exists only above the F_1 layer, being a certain fraction of O^+ (< 10%) in the F_2 layer^[44]. The N^+ concentration is given essentially by the following approximation of (33)

$$n^+(N) = \frac{n(N_2) I_2(N) + n(N) [I(N) + \gamma_{10} n^+(N_2)]}{\gamma_{12} n(O_2)} \quad (46)$$

The principal loss process of N^+ occurs by ion-atom interchange reactions with O_2 which apparently is very rapid. According to laboratory measurements made by Fite et al.^[17] a tentative value for the rate coefficient γ_{12} should be of the order of $5 \times 10^{-10} \text{ cm}^3 \text{ sec}^{-1}$. Even if such an experimental value is not precise, equation (46) shows that the absence of N^+ ions in the low ionosphere can be easily justified.

The first process of N^+ production is due to the dissociative photo-ionization of molecular nitrogen and requires the effect of solar radiation of $\lambda < 510 \text{ A}$. According to Weissler et al.^[45], the ionization cross-section of the process



is of the order of 10^{-19} cm^2 , i. e. about 100 times less than the ionization cross-section leading to N_2^+ . Since the ratio of molecular and atomic nitrogen concentrations $n(N_2)/n(N)$ is much greater than 100 at normal ionospheric levels, the ionization rate for atomic nitrogen at $\lambda < 510 \text{ A}$ is due to the dissociative photo-ionization of N_2 . In fact, since the optical depth for $\lambda < 800 \text{ A}$ depends on molecular nitrogen absorption, there is only the narrow spectral range

between 850 and 800 Å, which leads to a direct photo-ionization of N completely independent of N_2 . Thus, in using (43b), (46) is written as follows

$$n^+(N) = \frac{q(N) + \gamma_{10} n(N) [1-C] q(N_2) / [\gamma_8 n(O) + \gamma_9 n(O_2)]}{\gamma_{12} n(O_2)} \quad (48)$$

or, by the generally adequate approximation,

$$n^+(N) = \frac{n(N_2) I_2(N)}{\gamma_{12} n(O_2)} \quad (49)$$

When the optical depth becomes small enough (above the F_1 peak) the N^+ concentration, after a rapid increase, should be

$$n^+(N) \propto I_2(N) / \gamma_{12} \quad (50)$$

i.e. almost proportional to the ratio of rate coefficients since the variation of $n(N_2)/n(O_2)$ increases slowly with height.

From considerations of observational data obtained by Istomin^[44], it appears that $n^+(N)$ is not less than 10^4 cm^{-3} above the F_2 peak and can be obtained from an equation such as (49). At sufficiently high altitudes, the lifetime of N^+ ions is, however, very long, since

$$\tau(N^+) = 1 / \gamma_{12} n(O_2) \quad (51)$$

and (49) should only be used to determine the ion concentration for boundary conditions of the diffusive distribution above the F_2 peak. Also note that reaction (12) does not play a major role in creating


NO^+ , since reaction (1) involving O^+ and N_2 is, in fact, the essential process. From the present analysis, it appears that the N^+ concentration is not directly related to the presence of other ions but essentially to molecular oxygen. Thus, it can be neglected when the ionization conditions for all other ions are studied. Furthermore, the endothermic reaction $\text{N}^+ + \text{N}_2 \rightarrow \text{N}_2^+ + \text{N}$ used by Danilov^[48a] must be rejected since the interchange process transforming the molecular ion N_2^+ into the atomic ion N^+ is an exothermic reaction.

5 - Atomic oxygen ions


The principal reactions in which O^+ is involved lead to the following equation for photo-ionization equilibrium, (34)

$$n^+(\text{O})_{\text{eq}} = \frac{q(\text{O}) + n^+(\text{N}_2) \gamma_8 n(\text{O})}{\gamma_1 n(\text{N}_2) + \gamma_2 n(\text{O}_2)} \quad (52a)$$

The loss of O^+ ions by the associative process



with a rate coefficient certainly less than $10^{-16} \text{ cm}^3 \text{ sec}^{-1}$ is, of course, ignored in the presence of ion atom interchange reactions (1) and (2). The production of O^+ ions by the endothermic reaction



is not considered, since it cannot play an important role compared with other production sources.

With (39) and (43b), the steady state value (52a) becomes

$$n^+(O)_{eq} = \frac{q(O) + B(1 - C) q(N_2)}{\gamma_1 n(N_2) + \gamma_2 n(O_2)} \quad (52b)$$

in which $0 \leq B \leq 1$.

In the lower ionosphere B and C decrease to very small values while they increase with height in the F region. The maximum effect for an additional production of O^+ ions should occur for $B = 1$ and $C = 0$. Such conditions are never reached but are best approached in the F_1 layer. Thus, there is a contribution to the direct photo-ionization of atomic oxygen by the indirect photo-ionization of molecular nitrogen leading to an important production of the O^+ ion. At altitudes corresponding to the E layer, ion-atom interchange reactions (1) and (2) are important enough to transform O^+ ions into NO^+ and O_2^+ . If approximate ionization equilibrium can be retained for day-time conditions in the F_1 layer according to ionospheric observations, they must be rejected for the F_2 layer. Daytime and nighttime conditions for F_1 and F_2 layers clearly show that the loss coefficient

$$\gamma_1 n(N_2) + \gamma_2 n(O_2) = \beta(z) \quad (53)$$

becomes small so that

$$n^+(O) = n^+(O)_{t=0} e^{-\beta(z)t} + n^+(O)_{eq} [1 - e^{-\beta(z)t}] \quad (54)$$

In other words, the low electronic concentration in the F_1 layer for nighttime conditions mean that the principal loss processes (1) and (2) for atomic oxygen ions are large, and that steady state conditions can be used for day-time conditions. In the F_2 layer, the nighttime

concentration of O^+ ions is such that the steady state is far from being an adequate approximation for real conditions, which depend on diffusion processes. In fact, it is well known that diffusion leads to the F_2 peak and is active above it, so that photoequilibrium cannot be applied in that region.

From the analysis of ionospheric conditions by Danilov [48, 48a] and Ivanov-Kholodny [2], high values of γ of not less than $10^{-11} \text{ cm}^3 \text{ sec}^{-1}$ are deduced since $\beta(z)$ is at least 10^{-4} sec^{-1} and 10^{-3} sec^{-1} at 500 km and 300 km; respectively. All other results which have been obtained, however, lead to values less than 10^{-3} sec^{-1} at 300 km. For example, $\beta(300 \text{ km}) = 6.8 \times 10^{-4} \text{ sec}^{-1}$ given by Van Zandt et al. [49] is the largest value which is obtained for day-time conditions near sunspot maximum. A low value $\beta(300 \text{ km}) = 10^{-4} \text{ sec}^{-1}$ is deduced by Ratcliffe et al. [50] for nighttime data. A recent analysis made by Nisbet and Quinn [51] leads to even smaller values for nighttime data which must correspond to low temperatures of the thermopause.

It is clear, therefore, that equilibrium conditions cannot represent real conditions if $\beta(z) = \gamma n \leq 10^{-4} \text{ sec}^{-1}$. Equations (53) and (54) show that from such observational results γ cannot reach $10^{-11} \text{ cm}^3 \text{ sec}^{-1}$. A maximum average value should be $\gamma \leq 10^{-12} \text{ cm}^3 \text{ sec}^{-1}$. Furthermore, it is necessary to know the exact ratio γ_1/γ_2 in order to distinguish between the ion-atom interchange reactions of O^+ with N_2 and O_2 . It is not possible to obtain exact aeronomic conditions without knowing the ratio $\gamma_1 n(N_2)/\gamma_2 n(O_2)$. Again an apparent interpretation of ionospheric

behavior may be given with an arbitrary hypothesis since very many parameters are involved for which accurate values are as yet unknown.

6 - Molecular ions O_2^+ and NO^+


Mass-spectrometric measurements, [46] , [47] , [52] show that O_2^+ and NO^+ are the principal ions in the E layer and that O^+ becomes important only in the F_1 layer. Equilibrium conditions may be considered as a very good approximation to study the general behavior of molecular ions in a sunlit atmosphere, since they disappear via dissociative recombination.

Again neglecting reactions with NO, (35) and (36) are written as follows


$$n^+(NO) \left\{ \alpha(NO) n_e \right\} = q(NO) + [1-A] [\gamma_1 n(N_2) + \gamma_2 n(O_2)] n^+(O) + [\gamma_6 n(N) + \gamma_7 n(N_2)] n^+(O_2) \quad (55)$$

$$n^+(O_2) \left\{ \alpha(O_2) n_e + \gamma_6 n(N) + \gamma_7 n(N_2) \right\} = q(O_2) + A [\gamma_1 n(N_2) + \gamma_2 n(O_2)] n^+(O) + [1-B] [\gamma_8 n(O) + \gamma_9 n(O_2)] n^+(N_2) \quad (56)$$

It should again be made clear that we do not introduce, in the presence of the ion-atom interchange (2),



the production of molecular oxygen by the radiative association



introduced by Danilov^[19b] since the rate coefficient he has used, i. e. $5 \times 10^{-11} \text{ cm}^3 \text{ sec}^{-1}$, is too great by a factor of about 10^6 . Equations (55) and (56) show how $n^+(\text{NO})$ and $n^+(\text{O}_2)$ are related to the same loss reactions of atomic oxygen ions. An exact knowledge of the factor A defined by (38) is first required. But the ratio $n^+(\text{NO})/n^+(\text{O}_2)$ is subject to the effect of reactions (6) and (7) leading to a transformation of O_2^+ into NO^+ . Equation (55) leads to

$$\frac{n^+(\text{NO})}{n^+(\text{O}_2)} \geq \frac{\gamma_6 n(\text{N}) + \gamma_7 n(\text{N}_2)}{\alpha(\text{NO}) n_e} \quad (57)$$

In the E region, where $n_e = 10^5 \text{ cm}^{-3}$ and $n(\text{N}_2) = 10^{13}$ to 10^{12} cm^{-3} , a value $\gamma_7 \geq 10^{-15} \text{ cm}^3 \text{ sec}^{-1}$ leads to $n^+(\text{NO}) > n^+(\text{O}_2)$ if $\alpha(\text{NO}) = 10^{-7}$ to $10^{-8} \text{ cm}^3 \text{ sec}^{-1}$. Thus, if one assumes that the cross-section $Q(\text{O}_2^+ - \text{N}_2)$ can reach 10^{-20} cm^2 , there is a transformation of ions O_2^+ into NO^+ at the bottom of the E layer. Such a value of γ_7 would lead to a complete transformation of all ions into NO^+ in the D region, since N_2^+ can also be transformed into O_2^+ according to charge transfer process (9).

There is, therefore, an important problem to be resolved before determining the exact behavior of the D region and lower E layer. All mass-spectrometric observations, except perhaps during disturbed conditions, seem to show an increase of NO^+ compared with O_2^+ toward low altitudes. In addition, reaction (6) involving nitrogen

atoms should be considered as a process transforming O_2^+ into NO^+ since atomic nitrogen reaches its maximum concentration in the E layer^[43]. A sufficiently high value of γ_6 is required but is not impossible. As far as the reaction $O_2^+ + NO \rightarrow NO^+ + O_2$ is concerned, it should not be compared with (6) since $n(NO)$ is a very small fraction of $n(N)$ in the E layer according to equation (37). In any case, the present analysis shows that the behavior of the lower ionosphere cannot be studied by ignoring the possible effect of reactions (6) and (7).

Above a certain altitude, it is clear that O^+ is more important than O_2^+ in the production of the NO^+ ion. Equation (55) leads to

$$\frac{n^+(NO)}{n^+(O)} \geq \frac{\gamma_1 n(N_2)}{\alpha(NO) n_e} \quad (58)$$

which becomes, where $n^+(O) \approx n_e$,

$$n^+(NO) \geq \gamma_1 n(N_2) / \alpha(NO) \quad (59)$$

Equations (58) and (59) should be applied to the F_1 and F_2 layers, respectively; the NO^+ production essentially depends on the ion-atom interchange reaction (1).

Analysis of (56) shows that the direct production of O_2^+ ions by photoionization is an important process since all solar radiation of $\lambda < 1026 \text{ \AA}$ is involved. But the final concentration $n^+(O_2)$ is related to the effect of its transformation into NO^+ by reactions (6) and (7). Above a certain altitude, (56) leads to

$$\frac{n^+(O_2)}{n^+(O)} \geq \frac{\gamma_2 n(N_2) + q(O_2)/n^+(O)}{\alpha(O_2) n_e} \quad (60)$$

which becomes, where $n^+(O) \approx n_e$,

$$n^+(O_2) \geq \frac{q(O_2)}{\alpha(O_2)n_e} + \frac{\gamma_2 n(O_2)}{\alpha(O_2)} \quad (61)$$

Equations (60) and (61) should be applied to the F_1 and F_2 layers, respectively; O_2^+ production depends on the ion-atom interchange reaction (2) together with direct photo-ionization.

The general ionization equations (55) and (56) can be written, using equilibrium conditions (43b) and (52b),

$$n^+(NO) \left\{ \alpha(NO)n_e \right\} = q(NO) + [1-A] q(O) + [1-C] [B-AB] q(N_2) \\ + n^+(O_2) [\gamma_6 n(N) + \gamma_7 n(N_2)] \quad (62)$$

and

$$n^+(O_2) \left\{ \alpha(O_2)n_e \right\} = q(O_2) + A q(O) + [1-C] [1-(B-AB)] q(N_2) \\ - n^+(O_2) [\gamma_6 n(N) + \gamma_7 n(N_2)] \quad (63)$$

This analysis shows that, in the D region, the last term of the second member of equation (62) and (63) cannot be neglected when reactions (6) and (7) are involved. If the cross-section for reaction (7) reaches 10^{-20} cm^2 , it may even play a role in the lower E layer. When the transformation of O_2^+ into NO^+ can be ignored, it is clear from (62) and (63) that atmospheric conditions are simplified, since then one can write

$$n^+(\text{NO}) \left\{ \alpha(\text{NO}) n_e \right\} = [1-A] q(\text{O}) + [1-C][B-AB] q(\text{N}_2) + q(\text{NO}) \quad (64)$$

and

$$n^+(\text{O}_2) \left\{ \alpha(\text{O}_2) n_e \right\} = A q(\text{O}) + [1-C][1 - (B-AB)] q(\text{N}_2) + q(\text{O}_2) \quad (65a)$$

Application of (64) and (65a) in the F_1 layer requires the knowledge of a large number of parameters. Numerous experimental and aeronomic data are needed before the whole ionosphere will be susceptible to explanation even under photo-equilibrium conditions.

Note again that the conditions in the lower E layer or upper D region should not be studied by using an equation such as (65a) but, with (63) and (41), by

$$n^+(\text{O}_2) \left\{ \alpha(\text{O}_2) n_e \right\} = D \left\{ A q(\text{O}) + [1-C][1 - (B-AB)] q(\text{N}_2) + q(\text{O}_2) \right\} \quad (65b)$$

with A, B, C and D between 0 and 1. Factor D may be small in the lower ionosphere and certainly reach unity in the F_1 layer, C is small in the E layer but does not reach unity in the F_1 layer while B has its smallest value in the lower part of the ionosphere but increases with height; particularly in the diffusion region. A should decrease with height since $n(\text{O}_2)/n(\text{N}_2)$ decreases in the diffusion region.

In conclusion, the five equations (43), (48), (52), (62) and (65) show how each ion is related to the various productions $q(\text{N}_2)$, $q(\text{O})$, $q(\text{O}_2)$, $q(\text{NO})$ and $q(\text{N})$. Any aeronomic analysis must deal with the vertical distribution of the various factors A, B, C and D. An exact knowledge of these factors requires a precise determination of the rate coefficients of reactions (1) and (2), (8) and (9), (6) and (7)

and (12) with that of the recombination coefficients $\alpha(N_2)$, $\alpha(O_2)$ and $\alpha(NO)$ i. e. the knowledge of at least 10 parameters when the vertical distribution of neutral constituents (N_2 , O_2 , O and NO) with their ionization and absorption rate coefficient is perfectly known along with the solar flux. Since temperature effects are also important, it is clear that only approximate solutions can be obtained. Some simplification may be introduced when a specific ionospheric region is considered as shown in the analysis of the theoretical behavior of each ion. Nevertheless, the general complexity of equations (43) to (65) demonstrates that contradictory conclusions can be reached when a parameter is not well chosen. More in situ observations, particularly of ion densities, are needed before sufficient aeronomic data will be available to determine rate coefficients of reactions yet unknown from laboratory measurements.

7 - Hydrogen and helium ions

It is not the purpose of this work to discuss the behavior of H^+ and He^+ ions which play an important role above the F_2 peak where diffusion controls the electron and ion densities. However, it may be pointed out that charge transfer processes give the basic conditions (see for example, Bates and Patterson^[53]). At sufficiently low altitudes, the photo-ionization of H is less important than the charge transfer $H + O^+ \rightarrow H^+ + O$. For atomic hydrogen ions, the steady state is given by

$$\frac{n^+(H)}{n^+(O)} = \frac{9}{8} \frac{n(H)}{n(O)} \quad (66)$$

where the ratio 9/8 is the ratio of products of statistical weights considering



as the essential process. In the lower ionosphere, $n(\text{H})/n(\text{O})$ is small, less than 10^{-5} , and since $n^+(\text{O}) < n_e$ the concentration of atomic hydrogen ions is very small in the thermosphere. Above the F_2 peak, the ratio $n(\text{H})/n(\text{O})$ increases and $n^+(\text{O}) \approx n_e$. At high temperatures of the order of 2000°K $n(\text{H})/n(\text{O})$ is still of the order of 10^{-5} , and $n^+(\text{H})$ is less than 10 cm^{-3} . However, the ratio $n(\text{H})/n(\text{O})$ is very sensitive to the temperature; it reaches at 500 km about 10^{-3} for $T = 1000^\circ \text{K}$ and increases up to more than 10^{-1} at 700°K . It is, therefore, clear that H^+ becomes an important ion near 1000 km when the temperature decreases to about 1000°K .

For helium ions, the boundary conditions for the diffusive region are subject to the effect of its ionization by ultraviolet radiation. It will play an important role when the temperature is sufficiently high i. e. $T > 1000^\circ \text{K}$ only in the diffusive region and will remain a minor ion in the normal ionospheric layers.

REFERENCES

- [1] K. Watanabe and H. E. Hinteregger, Photoionization rates in the E and F regions, J. Geophys. Res., 67, 999, (1962).
- [2] G. S. Ivanov-Kholodny, Intensity of sun's short wave radiation and rate of ionization and recombination processes in the ionosphere, Geomagnetizm i Aeronomiya, 2, 377, (1962).
- [3] R. B. Norton, T. E. Van Zandt and J. S. Denison, A model of the atmosphere and Ionosphere in the E and F₁ regions, Proceedings International Conference on Ionosphere, Inst. Phys. and Phys. Soc. London, (1963).
- [4] L. A. Hall, K. R. Damon and H. E. Hinteregger, Solar extreme ultraviolet photon flux measurements in the upper atmosphere of August 1961, Proc. Third Int. Space Science Symp., North-Holland Publ. Cy., Amsterdam p. 745, (1963).
- [5] C. R. Detweiler, D. L. Garrett, J. P. Purcell and R. Tousey, The intensity distribution in the ultraviolet solar spectrum, Annales Geophys., 17, 263, (1961).
- [6] H. Zirin, L. A. Hall and H. E. Hinteregger, Analysis of the solar emission spectrum from 1300 to 250 Å as observed in August 1961, Proc. Third Int. Space Science Symp., North-Holland Publ. Cy., Amsterdam, p. 760, (1963).
- [7] H. E. Hinteregger, Absorption spectrometric analysis of the upper atmosphere in the EUV region, J. Atmos. Sciences, 19, 351, (1962).
- [8] M. Nicolet, Density of the heterosphere related to temperature, Smithsonian Institution, Res. Space Science, Special Report No. 75, (1961).
- [9] J. W. Townsend, Jr., Composition of the upper atmosphere, Chapter 7 in Physics and Medicine of the atmosphere and Space, John Wiley and Sons, New York, 1960.
- [9a] E. J. Schaefer, The dissociation of oxygen measured by a rocket-borne mass spectrometer, J. Geophys. Res., 68, 1175 (1963).
- [10] A. A. Pokhunkov, Variation in the night atmosphere of the mean molecular weight of the air in the 100 to 210 km altitude range according to mass spectrometric measurements, Iskusstvennye Sputniki Kemli, 12, 133, (1962).
- [11] L. A. Antonova and G. S. Ivanov-Kholodny, Corpuscular hypothesis for the ionization of the night-time ionosphere, Geomagnetizm i Aeronomiya, 1, 164, (1961).

- [12] Yu. I. Galperin, On the question of energy sources in the upper atmosphere, *Izv. Akad. Nauk SSSR, Ser. Geofiz.*, 2, 252, (1962).
- [13] B. J. O'Brien, Lifetime of outer zone electrons and their participation into the atmosphere, *J. Geophys. Res.* 67, 3687, (1962).
- [14] J. A. Gledhill and H. O. Van Rooyen, Cape Town anomaly and auroral emission, *Nature*, 196, 973, (1962).
- [15] P. H. G. Dickinson and J. Sayers, Ion charge exchange reactions in oxygen afterglows, *Proc. Phys. Soc.*, 76, 137, (1960).
- [16] G. F. O. Langstroth and J. B. Hasted, Contribution to the general discussion in Inelastic collisions of atoms and simple molecules, A general discussion of the Faraday Society, n° 33, 298 (1962).
- [16a] R. L. Potter, Ion-atom interchange in air, *J. Chem. Phys.*, 23, 2462, (1955).
- [16b] V. I. Krassovsky, Effective coefficient of recombination in the ionosphere, *Izv. Akad. Nauk, SSSR, Ser. Geofiz.*, n° 4, 504, (1957).
- [16c] V. L. Talrose, M. I. Maikin and I. K. Larin, The reaction of $O^+ + N_2 \rightarrow NO^+ + N$, in Inelastic Collisions of atoms and simple molecules, A general discussion of the Faraday Society, n° 33, 257, (1962).
- [17] W. L. Fite, J. A. Rutherford, W. R. Snow, and V. A. J. van Lint, Ion-neutral collisions in afterglows, in Inelastic collisions of atoms and simple molecules, A general discussion of the Faraday Society, n° 33, 264, (1962).
- [18] M. Hertzberg, Ion-neutral reactions, *J. Atm. Terr. Phys.*, 20, 177, (1961).
- [19] A. D. Danilov, Molecular ions in the upper atmosphere, *Dokl. Akad. Nauk SSSR*, 137, 1098, (1961).
- [19a] A. D. Danilov, On the problem of O_2^+ formation in the high atmosphere *Iskusstvennye Sputniki Zemli*, 7, 56, (1961).
- [19b] A. D. Danilov, On the question of NO^+ formation in the high atmosphere *Iskusstvennye Sputniki Zemli*, 5, 60, (1960).
- [19c] A. D. Danilov, Nitrogen dissociation in the upper atmosphere, *Geomagnetism i Aeronomiya*, 1, 174, (1961).
- [20] D. R. Bates and M. Nicolet, Ion-atom interchange, *J. Atmos. Terr. Phys.*, 18, 65, (1960); 21, 286, (1961).

- [21] E. V. Stupochenko, I. P. Stakhanov, E. V. Samuilov, A. S. Pleshanov, I. B. Rozhdestvenskii, Thermodynamic properties of the air in the temperature interval from 1000 to 12000°K and the pressure intervals from 0.001 to 1000 atmosphere, ARS J., 30, 98, (1960).
- [22] H. Friedman, Solar radiation, Astronautics, 7 (8), 14, (1962).
- [23] M. Nicolet and A. C. Aikin, The formation of the D region of the Ionosphere, J. Geophys. Res., 65, 1469, (1960).
- [23a] M. Nicolet, Contribution a l etude de l ionosphere, Institut Roy. Meteor., Bruxelles, Memoires, 19, (1945).
- [23b] M. Nicolet, Aeronomic conditions in the mesosphere and lower thermosphere, Penn State Univ., Scient. Rep., n° 102, (1958).
- [24] M. Nicolet, Meteor ionization and the night-time E layer, p. 99 in Meteors: Special Supplement (vol. 2) to J. Atm. Phys., Pergamon Press, London, (1955).
- [25] V. G. Istomin, Extraterrestrial ions in the terrestrial ionosphere, Iskustvennye Sputniki Zemli, 11, 98, (1961).
- [26] D. R. Bates and A. Dalgarno, Electronic recombination, Chapter 7 in Atomic and Molecular Processes, ed. D. R. Bates, Academic Press, New York, (1962).
- [27] A. C. Faire and K. S. W. Champion, Measurements of dissociative recombination and diffusion in nitrogen at low pressures, Phys. Rev. 113, 1, (1959).
- [28] W. H. Kasner, W. A. Rogers and M. A. Biondi, Electron-ion recombination coefficients in nitrogen and in oxygen, Phys. Rev. Letters, 7, 321, (1961).
- [29] J. Sayers, Recent Laboratory studies of recombination cross-sections, p. 212 in Solar Eclipses and the Ionosphere, Ed. Beynon and Brown, Pergamon Press, London, (1956).
- [30] M. Mentzoni, C. Montgomery and R. Row, Electron recombination and relaxation in heated nitrogen and dry air, Bull. Amer. Phys. Soc., I, 636, (1962).
- [31] R. C. Gunton and E. C. Y. Inn, Rates of electron removal by recombination attachment and ambipolar diffusion in nitric oxide plasmas, J. Chem. Phys., 35 1896, (1961).
- [32] J. P. Doering and B. H. Mahan, Photoionization of nitric oxide, J. Chem. Phys., 36, 669, (1962).

- [32] J. P. Doering and B. H. Mahan, Photoionization of nitric oxide, J. Chem. Phys., 36, 669, (1962).
- [33] S. C. Lin, cited by A. Dalgarno, [39], Charged particles in the upper atmosphere, Annales Géophys., 17, 16, (1961).
- [34] M. W. Syverson, R. P. Stein, T. M. Shaw, M. Scheibe and R. C. Gunton, Electron-NO⁺ recombination in shock-heated air, Bull. Amer. Phys. Soc., 7, 378, (1962).
- [35] R. C. Whitten and I. G. Popoff, A Model of solar-flare ionization in the D-region, J. Geophys. Res., 66, 2779, (1961) and corrected ibidem 67, 3000, (1962).
- [36] I. G. Popoff and R. C. Whitten, D-region ionization by solar x rays, J. Geophys. Res., 67, 2986, (1962).
- [37] J. A. Ratcliffe, A survey of solar eclipses and the ionosphere, in: Solar Eclipses and the ionosphere, Ed. Beynon and Brown, Pergamon Press, (1956).
- [38] J. E. Titheridge, Ionization below the night-time F-layer, J. Atm. Terr. Phys., 17, 126, (1959).
- [39] A. Dalgarno, Charged particles in the upper atmosphere, Annales Géophys., 17, 16, (1961).
- [40] L. M. Chanin, A. V. Phelps and M. A. Biondi, Measurement of the attachment of slow electrons in oxygen, Phys. Rev. Letters, 2, 344, (1959) and Phys. Rev., 128, 219, (1962).
- [41] L. M. Branscomb, D. S. Burch, S. J. Smith and S. Geltman, Photodetachment cross-section and electron affinity of atomic oxygen, Phys. Rev., 111, 504, (1958).
- [42] R. K. Curran, Formation of NO₂⁻ by charge transfer at very low energies, Phys. Rev., 125, 910, (1962).
- [43] M. Nicolet, Aeronomic chemical reactions, Chapter 2 in Physics and Medicine of the Atmosphere and Space, Ed. Benson and Strughold, John Wiley and Sons, New York, (1960).
- [44] V. G. Istomin, Nitrogen ions in the upper atmosphere of the earth and night-time ionization in the E-region, Dokl. Akad. Nauk, SSSR, 137, 1102, (1961).
- [45] G. L. Weissler, J. A. R. Samson, M. Ogawa and G. R. Cook, Photoionization analysis by mass spectrometry, J. Opt. Soc. Amer., 49, 338, (1959).

- [46] C. Y. Johnson, Aeronomic parameters from mass spectrometry, Annales Geophys., 17, 100, (1961).
- [47] H. A. Taylor and H. C. Brinton, Atmospheric ion composition measured above Wallops Island, Virginia, J. Geophys. Res., 66, 2587. (1961).
- [48] A. D. Danilov, Molecular ions in the upper atmosphere, Dokl. Akad. Nauk. SSSR, 137, 1098, (1961).
- [48a] A. D. Danilov, The formation of molecular ion in the high atmosphere, Iskusstvennye Sputniki Zemli, 8, 72, (1962).
- [49] T. E. Van Zandt, R. B. Norton, and G. H. Stonehocker, Photochemical rates in the equatorial F_2 region from the 1958 eclipse, J. Geophys. Res., 65, 2003, (1960).
- [50] J. A. Ratcliffe, E. R. Schmerling, C. S. G. K. Setty and J. O. Thomas, The rates of production and loss of electrons in the F region of the ionosphere, Phil. Trans. R. Soc., A 248, 621, (1956).
- [51] J. S. Nisbet and T. P. Quinn, The recombination coefficient of the night-time F layer, J. Geophys. Res., 68, No. 4, (1963).
- [52] V. G. Istomin, Absolute concentrations of ions between 100 and 200 km in the terrestrial atmosphere, Iskisstvennye Sputniki Zemli, 11, 94, (1961).
- [53] D. R. Bates and T. N. L. Patterson, Helium ions in the upper atmosphere, Planet. Space Science, 9, 599, (1962).