

INSTITUT D'AERONOMIE SPATIALE DE BELGIQUE

3 - Avenue Circulaire

B - 1180 BRUXELLES

AERONOMICA ACTA

A - N° 77 - 1970

Ultraviolet solar radiation related to mesospheric processes

by M. ACKERMAN

BELGISCH INSTITUUT VOOR RUIMTE-AERONOMIE

3 - Ringlaan

B - 1180 BRUSSEL

FOREWORD

This paper has been presented at the 4th ESRIN-ESLAB symposium held in Frascati from July 6 to 10, 1970 and will be published in the proceedings of the symposium.

AVANT-PROPOS

Cet article a été présenté au 4^e symposium ESRIN-ESLAB tenu à Frascati du 6 au 10 juillet 1970 et sera publié dans les proceedings de ce symposium.

VOORWOORD

Dit artikel werd voorgedragen op het 4^e ESRIN-ESLAB symposium gehouden te Frascati van 6 tot 10 juli 1970 en zal gepubliceerd worden in de proceedings van dit symposium.

VORWORT

Diese Arbeit würde zum 4. Symposium ESRIN-ESLAB in Frascati (6. zum 10. Juli 1970) vorgestellt und wird in den "Proceedings" dieses Symposium herausgegeben werden.

ULTRAVIOLET SOLAR RADIATION RELATED TO MESOSPHERIC PROCESSES

by

M. ACKERMAN

Abstract

The solar ultraviolet radiation involved in the mesospheric processes is subject to atmospheric attenuation at practically all wavelengths. For this reason, solar fluxes are discussed as well as absorption cross sections of molecular oxygen and of ozone.

Résumé

Le rayonnement ultraviolet solaire lié aux phénomènes mesosphériques est atténué par l'atmosphère elle-même à presque toutes les longueurs d'onde. Pour tenir compte de ce phénomène, des valeurs du flux solaire sont discutées de même que celles des sections efficaces d'absorption de l'oxygène moléculaire et de l'ozone.

Samenvatting

De ultraviolette zonnestraling, verbonden aan de mesosferische verschijnselen, wordt in de atmosfeer door praktisch alle golflengten geabsorbeerd. Om rekening te houden met deze atmosferische verschijnselen wordt de zonneflux alsmede de werkzame absorptiedoorsneden van moleculaire zuurstof en ozon voorgesteld.

Zusammenfassung

Die ultraviolette Sonnenstrahlung, die in den mesosphärischen Prozessen einkommt, wird beinahe an allen Wellenlängen durch die Atmosphäre absorbiert. Der Sonnenfluss sowie die Durchquerschnitte von molekularen Sauerstoff und Ozon werden vorgestellt.

INTRODUCTION

The atmospheric layer of strong negative temperature gradient called the mesosphere and located at altitudes between roughly 50 and 85 km is still part of the homosphere since the mean molecular mass is there maintained constant at all altitudes by the mixing of the major constituents. However the chemical reactions are very efficient in this region and strongly affect the concentration of minor constituents. These processes are of course not strictly limited in altitude and continue to play a role in the neighbouring atmospheric regions, namely the stratosphere below and the thermosphere above, by transport processes.

These interferences lead to some difficulties in defining the solar ultraviolet radiation of interest to the mesosphere where the solar photons of sufficient energy to initiate chemical chain reactions, penetrate with different degrees of attenuation. The two main absorbing species involved in this attenuation are molecular oxygen and ozone of which the photodissociation produces atomic oxygen, providing the main oxidizing agent in the chemosphere. Typical total numbers of molecules above 50 and 85 km are respectively $4 \times 10^{21} \text{ cm}^{-2}$ and $2 \times 10^{19} \text{ cm}^{-2}$ for O_2 and 10^{17} cm^{-2} and 10^{13} cm^{-2} for O_3 . Since maximum absorption cross sections are of the order of 10^{-17} cm^2 oxygen and to a lesser degree ozone will control the penetration of solar ultraviolet radiation in the mesosphere.

The data actually available on the solar ultraviolet flux from 1100 Å to 3500 Å will be discussed here. The short wavelength limit will allow inclusion of the Lyman α radiation which plays a role in the photodissociation of O_2 , H_2O , CO_2 , etc. and which constitutes the bridge between the chemosphere and the ionosphere owing to its important role in the formation of the D region by ionizing nitric oxide. The long wavelength limit has been fixed at the beginning of the ultraviolet absorption of ozone in the Huggins bands.

On the other hand, to allow the evaluation of the number of photons available at different altitudes, absorption cross section data on O_2 and O_3 will be discussed.

Finally, adopted values for the three parameters will be listed for fixed wavelength intervals suitable for photochemical computations.

ABSORPTION CROSS SECTION OF OZONE

Between 3500 Å and 2000 Å, ozone absorbs in a wide band discovered by Hartley and peaking near 2500 Å. At this wavelength, unit optical depth for a typical ozone distribution takes place at the lower boundary of the mesosphere for an overhead sun and at 70 km for a solar zenith distance of 90°. This indicates the important mesospheric role of the Hartley band of ozone which has a bell shape presenting a diffuse structure corresponding to a complicated set of diffuse bands. On the long wavelength side, between 3100 Å and 3500 Å, the structure appears more clearly and was discovered by Huggins in the spectrum of Sirius near the horizon.

The photodissociation of ozone may lead to molecular and atomic oxygen in various excited states, as was already pointed out more than 30 years ago by Nicolet (1939). The absorption cross section of ozone has been well known since the work of Vigroux (1953), Watanabe et al (1953) and Inn and Tanaka (1953). A slight disagreement between the values given by Vigroux (1953) and Inn and Tanaka (1953) at wavelengths smaller than 2700 Å prompted the measurements of Hearn (1961). These have been confirmed by Griggs (1968). New measurements of Vigroux (1969) have definitely shown the correctness of the data of Inn and Tanaka (1953). Various sets of data are represented in Fig. 1. Below 2000 Å, only one set of data exists to our knowledge, those of Watanabe et al (1953). Below 1800 Å ozone has so far not been shown to play a role neither in the photochemistry nor in the transparency of the earth atmosphere, since oxygen is there the main absorbant and since the solar ultraviolet intensity is becoming so much smaller compared to that in the 2000 to 3500 Å region.




Fig. 1.- Absorption cross section of ozone versus wavelength in the Hartley and Huggins bands.

For the sake of completeness, absorption cross section values in the Chappius bands are represented in Fig. 2.

It appears that absorption cross sections of ozone are now very well known. However the question about the quantum yield of the various photodissociation processes is not yet fully answered.

ABSORPTION CROSS SECTION OF MOLECULAR OXYGEN

Many data exist on the absorption cross section of molecular oxygen. A critical analysis is necessary before adopting useful aeronomic values. This was done several times in the past, by Nicolet and Mange (1954) for instance. Many new measurements have been made in the last few years which justify a new analysis. This will be made for successive wavelength intervals.

1160 Å - 1370 Å

The absorption by molecular oxygen changes rapidly with wavelength in this region, presenting optical windows that allow the penetration of the solar radiation in the mesosphere. Absorption cross sections have for instance been obtained by Watanabe et al (1953, 1958).

One optical window which is particularly well known for its fundamental importance in mesospheric problems allows the deep penetration of the solar Lyman α radiation. Values of absorption cross section of O_2 at 1215.67 Å have been obtained by various authors (Preston, 1940 ; Watanabe et al, 1953 ; Ditchburn et al, 1954 ; Lee, 1955 ; Watanabe et al, 1958 ; Metzger and Cook, 1964 ; Ogawa, 1968 and Gailly, 1969) and range from 8.5×10^{-21} to $1.04 \times 10^{-20} \text{ cm}^2$. This leads to unit optical atmospheric depth at about 75 km altitude for an overhead sun. The Lyman α line has been shown to be slightly on the short wavelength wing of the optical window and pressure effects on the absorption have not yet been satisfactorily interpreted.




Fig. 2.- Absorption cross section of ozone versus wavelength in the Chappuis bands.

1370 Å - 1750 Å

This region is characterized by the Schumann-Runge continuum which presents high values of the absorption cross section measured by various authors (Landenburg and Van Voorhis, 1933 ; Schneider, 1940 ; Ditchburn and Heddle, 1953 ; Watanabe and Marmo, 1956 ; Huffman et al, 1964 ; Metzger and Cook, 1964 and Hudson et al, 1966) as shown in Fig. 3, leading to a complete extinction of the solar radiation in the thermosphere.

1750 Å - 2431 Å

Many absorption cross section values discussed recently by Ackerman et al (1970) have been obtained since the beginning of this century for the Schumann-Runge bands of O₂ which extend from 1750 Å to 2000 Å. While they indicate the important role that the bands can play in the mesospheric photochemistry they are however not very valuable for precise aeronautical evaluations.

This situation has recently been considerably improved. New laboratory measurements concerning the absorption properties have been obtained by Ackerman et al (1969) and concerning the structure of the bands by Ackerman and Biaumé (1970) and have led to a detailed evaluation of the absorption cross section taking temperature effects into account (Ackerman et al, 1970). First applications of these new data will be presented by Kockarts (1970) in this symposium.

Above 2000 Å the Herzberg continuum extends up to 2439 Å. Its absorption cross section ranges from about 10^{-23} to 10^{-24} cm². Such low values imply that in this wavelength region, molecular oxygen influences the atmospheric transparency only in the stratosphere at heights around 25 km. In fact the strong absorption of ozone limits the role of oxygen to the narrow optical window at 2000 Å which is essential in the process of stratospheric ozone formation.




Fig. 3.- Absorption cross section of molecular oxygen versus wavelength and versus wavenumber in the Schumann continuum.

SOLAR FLUX

When the first spectroscopic measurements of the solar ultraviolet radiation were undertaken, the conclusion was drawn that the considerable intensity decay taking place at 3000 Å towards shorter wavelengths was due to solar absorption itself. This phenomenon was attributed to the absorption by ozone after the presence of this atmospheric constituent was firmly established by Fowler and Lord Rayleigh in (1917), and after ground-based study of its absorption of the solar radiation at various zenith distances was performed by Fabry and Buisson in (1921).

At shorter wavelengths, rocket borne experiments have brought information on the solar ultraviolet radiation since 1946. These data have already been reviewed several times (Hinteregger, 1965). In the wavelength range here considered the available values of solar radiation intensities were until recently those published in 1961 by Detwiler et al (1961). In the range from 2000 Å to 1400 Å, these authors believed that the accuracy of their data was better than + 20% and were fairly sure that there were no errors greater than a factor of 1.5, whereas below 1300 Å errors as great as a factor of two or more were possible.

Using also photographic detection, Bonnet (1968) has obtained values of the solar flux from about 2000 Å to 3000 Å by integrating the flux measured at different places of the solar disk using a rocket-borne instrument. His data indicated a much more important discontinuity centered at 2085 Å, than had been previously measured. However his values were practically in agreement with the previous ones outside of the discontinuity.

In 1968 also, Ackerman et al (1968) reported observations made during the first flight of a balloon borne photoelectric monochromator viewing the whole solar disk which indicated an even larger amplitude for the discontinuity. Their results showed also that the solar ultraviolet intensity below 2000 Å had been previously overestimated. This has since been confirmed by subsequent flights which will be described elsewhere, one of which being discussed in detail by Frimout (1969).

Measurements in the solar spectrum between 1400 and 1875 Å with a photoelectric rocket-borne spectrometer have been published in 1969 by Parkinson and Reeves (1969) and correspond to a solar black body temperature of the order of 4500°K or less, namely much lower than the earlier values given by Detwiler et al (1961).

This year finally, Widing et al (1970) analysing rocket spectra photographed in 1966, have deduced new flux values which are situated as shown in Fig. 4, between the two extremes. They are in agreement with the data of Ackerman et al (1968) at 1950 Å.

DISCUSSION AND CONCLUSION

The solar ultraviolet of interest to the mesosphere may now be defined more easily. It appears to cover the whole wavelength interval that we have examined after excluding the range covered by the Schumann continuum. Below this feature, considering the absolute intensity, only the Lyman α line plays an important role. Since the sun is radiating more and more strongly towards longer wavelengths above 1700 Å the solar photons have an increasingly important effect in the mesosphere. Below 2000 Å, the radiation has very much the character of a continuum, while above, it presents a very complicated structure. In some applications intensity data at high resolution would be very useful.

Until two or three years ago the situation was rather comfortable from the aeronomical point of view since only one set of data existed. All authors were referring to the work of Detwiler et al (1961). The situation now appears different since new data have been obtained showing that these authors were a little too optimistic about the accuracy of their measurements.

The new values necessarily imply making a choice. This choice has to be precisely specified to allow the comparison of the results obtained in photochemical evaluations, for instance. One set of adopted values




Fig. 4.- Flux of solar photons at one A.U. versus wavelength from 1000 Å to 3000 Å.

is presented in Table I with values of absorption cross sections of O_2 and O_3 . The adopted solar ultraviolet flux is based on the data given by Detwiler et al (1961) from 1163 to 1428 Å, by Widing et al (1970) from 1428 Å to 1942 Å, by Ackerman et al (1968) and by Frimout (1969) from 1942 Å to 2299 Å, by Detwiler et al (1961) and by Tousey (1963) from 2299 Å to 3625 Å and by Johnson (1954) at longer wavelength. This is of course not a final choice and new data are now required.

The question of variability with time arises more strongly than ever. The dispersion of the experimental results shows at least that experimental techniques have to be considerably improved before an answer can be given. For Lyman α , a rather important variability may be taken as certain.

Finally, if ozone and oxygen have up to now been considered as the absorbers of the ultraviolet radiation related to the mesosphere, one has still to be cautious since other absorbers might be involved. Indications of such effects have been recently given in the 1300 Å region by Reid and Withbroe (1970) and in the 2000 Å region by Ackerman et al (1968).

REFERENCES

- M. ACKERMAN and F. BIAUME, "Structure of the Schumann-Runge bands from the O-O to the 13-0 band", J. Mol. Spectry, 35 (1970) 73-00.
- M. ACKERMAN, F. BIAUME and G. KOCKARTS, "Absorption cross sections of the Schumann-Runge bands of molecular oxygen", Planet. Space Science, 00(1970) 000.
- M. ACKERMAN, F. BIAUME and M. NICOLET, "Absorption in the spectral range of the Schumann-Runge bands", Can. J. Chem., 47 (1969) 1834-1840.
- M. ACKERMAN, D. FRIMOUT et R. PASTIELS, "Mesure du rayonnement ultraviolet solaire par ballon stratosphérique", Ciel et Terre, 84 (1968) 408-419.
- R.M. BONNET, "Stigmatic spectra of the sun between 1800 Å and 2800 Å", Space Research VII (1968) 458-472.
- C.R. DETWILER, D.L. GARRETT, J.D. PURCELL and R. TOUSEY, "The intensity distribution in the ultraviolet solar spectrum, Ann. Geophys., 17 (1961) 9-18.
- R.W. DITCHBURN, J.E.S. BRADLEY, C.G. CANNON and G. MUNDAY, "Absorption cross sections for Lyman α and neighbouring lines" in Rocket Exploration of the upper atmosphere (ed. by R.L. Boyd and M.J. Seaton), Pergamon Press, London, 1954, p 327-334.
- R.W. DITCHBURN and D.W.O. HEDDLE, "Absorption cross section in the vacuum ultraviolet. I. continuous absorption of oxygen (1800 to 1300 Å)", Proc. Roy. Soc., 220 A (1953) 61-70.
- C. FABRY and H. BUISSON, "Etude de l'extrémité ultraviolette du spectre solaire", J. Phys. et Radium, II S VI (1921) 197-226.
- A. FOWLER and R.J. STRUTT, "Absorption bands of atmospheric ozone in the spectra of sun and stars", Proc. Roy. Soc., A 93 (1917) 577-586.
- D. FRIMOUT, "De meting van de zonneintensiteit in het optisch venster van 2000 to 2200 Å", Thesis Rijksuniversiteit, Gent, Academiejaar 1969-1970.

- T.D. GAILLY, "Optical absorption coefficient of molecular oxygen near 1215 Angstroms", J. Opt. Soc. Amer., 59 (1969) 536-538.
- M. GRIGGS, "Absorption coefficients of ozone in the ultraviolet and visible region", J. Chem. Phys., 49 (1968) 857-859.
- A.G. HEARN, "The absorption of ozone in the ultraviolet and visible regions of the spectrum", Proc. Phys. Soc., 78 (1961) 932-940.
- H.E. HINTEREGGER, "Absolute intensity measurements in the extreme ultraviolet spectrum of solar radiation", Space Sci. Rev., 4 (1965) 461-497.
- R.D. HUDSON, V.L. CARTER and J.A. STEIN, "An investigation of the effect of temperature on the Schumann-Runge absorption continuum of oxygen, 1580-1950 Å", J. Geophys. Res., 71 (1966) 2295-2298.
- R.E. HUFFMAN, Y. TANAKA and J.C. LARABEE, "Nitrogen and oxygen absorption cross sections in the vacuum ultraviolet", Disc. Farad. Soc., 37 (1964) 159-165.
- E.C.Y. INN and Y. TANAKA, "Absorption coefficient of ozone in the ultraviolet and visible regions", J. Opt. Soc. Amer., 43 (1953) 870-873.
- F.S. JOHNSON, "The solar constant", J. Meteorol., 11 (1954) 431-439.
- G. KOCKARTS, this symposium.
- R. LANDENBURG and C.C. VAN VOORHIS, "The continuous absorption of oxygen between 1750 and 1300 Å and its bearing upon the dispersion", Phys. Rev., 43 (1933) 315-321.
- P. LEE, "Photodissociation and photoionization of Oxygen (O₂) as inferred from measured absorption coefficients", J. Opt. Soc. Amer., 45 (1955) 703-709.
- P.H. METZGER and R.G. COOK, "A reinvestigation of the absorption cross section of molecular oxygen in the 1050-1800 Å region", J. Quant. Spectry. Radiative Transfer, 4 (1964) 107-116.
- M. NICOLET, "Le problème atomique dans l'atmosphère supérieure", Mém. Inst. Météor. Belg. XI (1939).
- M. NICOLET and P. MANGE, "The dissociation of oxygen in the high atmosphere", J. Geophys. Res., 59 (1954) 15-45.
- M. OGAWA, "Absorption coefficients of O₂ at the Lyman alpha line and its vicinity", J. Geophys. Res., 73 (1968) 6759-6763.

- W.H. PARKINSON and E.M. REEVES, "Measurements in the solar spectrum between 1400 and 1875 Å with a rocket borne spectrometer", Solar Phys., 10 (1969) 342-347.
- W.H. PRESTON, "The origin of radio fade outs and the absorption coefficient of gases for light wavelength 1215.7 Å", Phys. Rev., 57 (1940) 885-887.
- R.H.G. REID and G.L. WITHBROE, "The density and vibrational distribution of molecular oxygen in the lower thermosphere", Harvard College Observatory TR-16, 1970.
- E.G. SCHNEIDER, "An estimate of the absorption of air in the extreme ultraviolet", J. Opt. Soc. Amer., 30 (1940) 128-132.
- R. TOUSEY, "The extreme ultraviolet spectrum of the sun", Space Sci. Rev., 2 (1963) 3-69.
- E. VIGROUX, Contribution à l'étude expérimentale de l'ozone, Masson et Cie, 1953.
- E. VIGROUX, "Coefficients d'absorption de l'ozone dans la bande de Hartley", Ann. Géophys., 25 (1969) 169-172.
- K. WATANABE, E.C.Y. INN and M. ZELIKOFF, "Absorption coefficients of oxygen in the vacuum ultraviolet", J. Chem. Phys., 21 (1953) 1026-1030.
- K. WATANABE and F.F. MARMO, "Photoionization and total absorption cross section of gases. II. O₂ and N₂ in the region 850-1500 Å", J. Chem. Phys. 25 (1956) 967-971.
- K. WATANABE, M. SAKAI, J. MOTTI and T. NAKAYAMA, Absorption cross section of O₂, NO and NO₂ with an improved photoelectric method, Contribution nr 11, Hawaii Institute of geophysics, 1958.
- K. WATANABE, M. ZELIKOFF and E.C.Y. INN, "Absorption coefficients of several atmospheric gases" Geophys. Res. Pap. nr 21, A.F.C.R.L. (1953).
- K.G. WIDING, J.D. PURCELL and G.D. SANDLIN, "The UV continuum 1450-2100 Å and the problem of the solar temperature minimum", Solar Phys., 12 (1970) 52-62.

TABLE I.1.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 Å

N°	$\Delta\lambda$ (Å)	$\Delta\nu$ (cm^{-1})	q ($\text{cm}^{-2}\text{sec}^{-1}$)	$\sigma(O_2)$ (cm^2)	$\sigma(O_3)$ (cm^2)
1	Ly α 1.215,67	82.259	3.00×10^{11}	1.00×10^{-20}	2.32×10^{-17}
2	1.170-1.163	85.500-86.000	1.03×10^8	2.00×10^{-20}	7.80×10^{-18}
3	1.176-1.170	85.000-85.500	2.66	1.25×10^{-18}	7.97
4	1.183-1.176	84.500-85.000	1.12	2.55×10^{-19}	8.66
5	1.190-1.183	84.000-84.500	1.24	3.00×10^{-20}	9.51
6	1.198-1.190	83.500-84.000	1.82	3.75×10^{-19}	1.25×10^{-17}
7	1.205-1.198	83.000-83.500	1.90	4.45×10^{-18}	1.84
8	1.212-1.205	82.500-83.000	7.40	8.35	2.19
9	1.220-1.212	82.000-82.500	2.28×10^9	6.00×10^{-19}	2.30
10	1.227-1.220	81.500-82.000	3.67	2.35	2.26
11	1.235-1.227	81.000-81.500	1.36	4.50	2.06
12	1.242-1.235	80.500-81.000	1.61	3.35	1.30
13	1.250-1.242	80.000-80.500	1.32	1.75×10^{-17}	8.91×10^{-18}
14	1.258-1.250	79.500-80.000	1.41	8.95×10^{-19}	7.24
15	1.266-1.258	79.000-79.500	3.11	4.30	6.09
16	1.274-1.266	78.500-79.000	1.06	1.10	5.66
17	1.282-1.274	78.000-78.500	1.37	2.05	5.87
18	1.290-1.282	77.500-78.000	1.02	4.43	6.47
19	1.299-1.290	77.000-77.500	1.14	5.55	8.14
20	1.307-1.299	76.500-77.000	7.29	4.20	1.24×10^{-17}
21	1.316-1.307	76.000-76.500	2.20	6.85	1.52
22	1.324-1.316	75.500-76.000	1.59	1.45×10^{-18}	1.47
23	1.333-1.324	75.000-75.500	2.21	2.25	1.51

TABLE I.2.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 Å

N°	$\Delta\lambda$ (Å)	$\Delta\nu$ (cm^{-1})	q ($\text{cm}^{-2}\text{sec}^{-1}$)	$\sigma(O_2)$ (cm^2)	$\sigma(O_3)$ (cm^2)
24	1.342-1.333	74.500-75.000	1.24×10^{10}	2.30×10^{-18}	1.51×10^{-17}
25	1.351-1.342	74.000-74.500	1.99×10^9	4.55	1.65
26	1.360-1.351	73.500-74.000	3.09	7.23	1.54
27	1.370-1.360	73.000-73.500	2.57	9.50	1.35
28	1.379-1.370	72.500-73.000	2.74	1.23×10^{-17}	1.05
29	1.389-1.379	72.000-72.500	3.10	1.32	7.97×10^{-18}
30	1.408-1.389	71.000-72.000	7.60	1.36	7.17
31	1.428-1.408	70.000-71.000	1.01×10^{10}	1.40	6.28
32	1.449-1.428	69.000-70.000	1.30	1.48	5.66
33	1.470-1.449	68.000-69.000	1.82	1.41	5.23
34	1.492-1.470	67.000-68.000	2.33	1.29	4.47
35	1.515-1.492	66.000-67.000	2.66	1.15	3.69
36	1.538-1.515	65.000-66.000	2.90	9.91×10^{-18}	2.93
37	1.562-1.538	64.000-65.000	3.60	8.24	2.19
38	1.587-1.562	63.000-64.000	4.75	6.58	1.63
39	1.613-1.587	62.000-63.000	6.40	4.97	1.20
40	1.639-1.613	61.000-62.000	5.49	3.45	9.77×10^{-19}
41	1.667-1.639	60.000-61.000	1.19×10^{11}	2.08	8.66
42	1.695-1.667	59.000-60.000	1.76	1.23	8.14
43	1.724-1.695	58.000-59.000	2.32	7.22×10^{-19}	8.17
44	1.739-1.724	57.500-58.000	1.44	4.58	8.57
45	1.754-1.739	57.000-57.500	1.83	2.74	8.40
46	1.770-1.754	56.500-57.000	2.34	*	8.11
47	1.786-1.770	56.000-56.500	2.62	*	7.99

* See Ackerman et al (1970).

TABLE I.3.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 Å

N°	$\Delta\lambda$ (Å)	$\Delta\nu$ (cm ⁻¹)	q (cm ⁻² sec ⁻¹)	$\sigma(O_2)$ (cm ²)	$\sigma(O_3)$ (cm ²)
48	1.802-1.786	55.500-56.000	2.88×10^{11}	*	7.86×10^{-19}
49	1.818-1.802	55.000-55.500	3.14	*	7.63
50	1.835-1.818	54.500-55.000	3.81	*	7.29
51	1.852-1.835	54.000-54.500	4.43	*	6.88
52	1.869-1.852	53.500-54.000	4.95	*	6.40
53	1.887-1.869	53.000-53.500	5.94	*	5.88
54	1.905-1.887	52.500-53.000	6.59	*	5.31
55	1.923-1.905	52.000-52.500	7.26	*	4.80
56	1.942-1.923	51.500-52.000	9.85	*	4.38
57	1.961-1.942	51.000-51.500	1.27×10^{12}	*	4.11
58	1.980-1.961	50.500-51.000	1.39	*	3.69
59	2.000-1.980	50.000-50.500	1.53	*	3.30
60	2.020-2.000	49.500-50.000	1.60	*	3.26
61	2.041-2.020	49.000-49.500	1.74	1.14×10^{-23}	3.26
62	2.062-2.041	48.500-49.000	2.31	1.05	3.51
63	2.083-2.062	48.000-48.500	4.20	1.00	4.11
64	2.105-2.083	47.500-48.000	7.30	9.55×10^{-24}	4.84
65	2.128-2.105	47.000-47.500	9.42	8.93	6.26
66	2.150-2.128	46.500-47.000	1.06×10^{13}	8.28	8.57
67	2.174-2.150	46.000-46.500	1.34	7.60	1.17×10^{-18}
68	2.198-2.174	45.500-46.000	1.32	6.92	1.52
69	2.222-2.198	45.000-45.500	1.73	6.28	1.97
70	2.247-2.222	44.500-45.000	1.80	5.65	2.55
71	2.273-2.247	44.000-44.500	1.82	5.03	3.24

See Ackerman et al (1970)

TABLE I.4.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 Å

N°	$\Delta\lambda$ (Å)	$\Delta\nu$ (cm ⁻¹)	q (cm ⁻² sec ⁻¹)	$\sigma(O_2)$ (cm ²)	$\sigma(O_3)$ (cm ²)
72	2.299-2.273	43.500-44.000	2.26×10^{13}	4.40×10^{-24}	4.00×10^{-18}
73	2.326-2.299	43.000-43.500	2.40	3.76	4.83
74	2.353-2.326	42,500-43.000	2.25	3.09	5.79
75	2.381-2.353	42.000-42.500	2.21	2.44	6.86
76	2.410-2.381	41.500-42.000	2.32	1.75	7.97
77	2.439-2.410	41.000-41.500	2.50	6.74×10^{-25}	9.00
78	2.469-2.439	40.500-41.000	2.73		1.00×10^{-17}
79	2.500-2.469	40.000-40.500	2.88		1.07
80	2.532-2.500	39.500-40.000	3.02		1.11
81	2.564-2.532	39.000-39.500	3.97		1.12
82	2.597-2.564	38.500-39.000	7.13		1.11
83	2.632-2.597	38.000-38.500	4.37		1.03
84	2.667-2.632	37.500-38.000	1.12×10^{14}		9.43×10^{-18}
85	2.703-2.667	37.000-37.500	1.25		8.23
86	2.740-2.703	36.500-37.000	1.16		6.81
87	2.778-2.740	36.000-36.500	1.19		5.31
88	2.817-2.778	35.500-36.000	1.38		3.99
89	2.857-2.817	35.000-35.500	1.70		2.84
90	2.899-2.857	34.500-35.000	2.46		1.92
91	2.941-2.899	34.000-34.500	3.90		1.14
92	2.985-2.941	33.500-34.000	3.99		6.60×10^{-19}
93	3.030-2.985	33.000-33.500	3.86		3.69
94	3.077-3.030	32.500-33.000	5.08		1.97

TABLE I.5.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 \AA

N°	$\Delta\lambda(\text{\AA})$	$\Delta\nu(\text{cm}^{-1})$	$q(\text{cm}^{-2}\text{sec}^{-1})$	$\sigma(O_3)(\text{cm}^2)$
95	3.100 (\pm 25)	32.520-32.000	5.92×10^{14}	1.05×10^{-19}
96	3.150	32.000-31.496	6.05	5.23×10^{-20}
97	3.200	31.496-31.008	6.94	2.91
98	3.250	31.008-30.534	8.12	1.50
99	3.300	30.534-30.075	9.71	7.78×10^{-21}
100	3.350	30.075-29.630	8.97	3.72
101	3.400	29.630-29.197	9.44	1.71
102	3.450	29.197-28.777	1.01×10^{15}	7.46×10^{-22}
103	3.500	28.777-28.369	1.03	2.66
104	3.550	28.369-27.972	1.03	1.09
105	3.600	27.972-27.586	1.04	5.49×10^{-23}
106	3.650	27.586-27.211	1.18	-
107	3.700	27.211-26.846	1.23	-
108	3.750	26.846-26.490	1.24	-
109	3.800	26.490-26.144	1.17	-
110	3.850	26.144-25.806	1.11	-
111	3.900	25.806-25.478	1.09	-
112	3.950	25.478-25.157	1.19	-
113	4.000	25.157-24.845	1.54	-
114	4.050	24.845-24.540	1.90	-
115	4.100	24.540-24.242	1.99	2.91
116	4.150	24.242-23.952	1.99	3.14
117	4.200	23.952-23.669	2.02	3.99
118	4.250	23.669-23.392	2.01	6.54
119	4.300	23.392-23.121	1.94	6.83
120	4.350	23.121-22.851	1.98	8.66

TABLE I.6.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 Å

N°	$\Delta\lambda(\text{Å})$	$\Delta\nu(\text{cm}^{-1})$	$q(\text{cm}^{-2}\text{sec}^{-1})$	$\sigma(O_3)(\text{cm}^2)$
121	4.400 (+ 25)	22.851-22.599	2.25×10^{15}	1.25×10^{-22}
122	4.450	22.599-22.346	2.39	1.49
123	4.500	22.346-22.099	2.48	1.71
124	4.550	22.099-21.858	2.49	2.12
125	4.600	21.858-21.622	2.48	3.57
126	4.650	21.622-21.390	2.50	3.68
127	4.700	21.390-21.164	2.55	4.06
128	4.750	21.164-20.942	2.61	4.89
129	4.800	20.942-20.725	2.59	7.11
130	4.850	20.725-20.513	2.46	8.43
131	4.900	20.513-20.504	2.44	8.28
132	4.950	20.504-20.100	2.53	9.09
133	5.000	20.100-19.900	2.48	1.22×10^{-21}
134	5.050	19.900-19.704	2.49	1.62
135	5.100	19.704-19.512	2.50	1.58
136	5.150	19.512-19.324	2.43	1.60
137	5.200	19.324-19.139	2.43	1.78
138	5.250	19.139-18.957	2.52	2.07
139	5.300	18.957-18.779	2.58	2.55
140	5.350	18.779-18.605	2.64	2.74
141	5.400	18.605-18.433	2.67	2.88
142	5.450	18.433-18.265	2.70	3.07
143	5.500	18.265-18.100	2.68	3.17
144	5.550	18.100-17.937	2.66	3.36
145	5.600	17.937-17.778	2.66	3.88

TABLE I.7.- Flux of solar photons, q , at one A.U., absorption cross section of O_2 and of O_3 , $\sigma(O_2)$ and $\sigma(O_3)$, for wavelength intervals $\Delta\lambda$ and wavenumber intervals $\Delta\nu$ from Ly α to 7300 Å

N°	$\Delta\lambda(\text{Å})$	$\Delta\nu(\text{cm}^{-1})$	$q(\text{cm}^{-2}\text{sec}^{-1})$	$\sigma(O_3)(\text{cm}^2)$
146	5.650 (\pm 25)	17.778-17.621	2.67×10^{15}	4.31×10^{-21}
147	5.700	17.621-17.467	2.67	4.67
148	5.750	17.467-17.316	2.69	4.75
149	5.800	17.316-17.667	2.71	4.55
150	5.850	17.667-17.021	2.71	4.35
151	5.900	17.021-16.878	2.71	4.42
152	5.950	16.878-16.736	2.72	4.61
153	6.000	16.736-16.598	2.72	4.89
154	6.050	16.598-16.461	2.71	4.84
155	6.100	16.461-16.326	2.70	4.54
156	6.150	16.326-16.194	2.70	4.24
157	6.200	16.194-16.064	2.70	3.90
158	6.250	16.064-15.936	2.69	3.60
159	6.300	15.936-15.810	2.68	3.43
160	6.350	15.810-15.686	2.67	3.17
161	6.400	15.686-15.564	2.66	2.74
162	6.450	15.564-15.444	2.65	2.61
163	6.500 (\pm 50)	15.384-15.265	3.95	2.40
164	6.600	15.265-15.038	5.22	2.07
165	6.700	15.038-14.815	5.18	1.72
166	6.800	14.815-14.598	5.14	1.37
167	6.900	14.598-14.388	5.09	1.11
168	7.000	14.388-14.184	5.04	9.13×10^{-22}
169	7.100	14.184-13.986	4.99	7.93
170	7.200	13.986-13.793	4.94	6.40
171	7.300	13.793-13.605	4.90	5.14