

I N S T I T U T D ' A E R O N O M I E S P A T I A L E D E B E L G I O U E

3 - Avenue Circulaire
B - 1180 BRUXELLES

AERONOMICA ACTA

A - N° 165 - 1976

Magnetic holes in the solar wind

by

J.M. TURNER, L.F. BURLAGA, N.F. NESS

and J. LEMAIRE

B E L G I S C H I N S T I T U U T V O O R R U I M T E - A E R O N O M I E

3 - Ringlaan
B - 1180 BRUSSEL

FOREWORD

The Article "Magnetic holes in the solar wind" by T.M. Turner, L.F. Burlaga, N.F. Ness, J. Lemaire, will be published in Journal of Geophysical Research.

AVANT-PROPOS

L'article "Magnetic holes in the solar wind" par J.M. Turner, L.F. Burlaga, N.F. Ness, J. Lemaire, sera publié dans Journal of Geophysical Research.

VOORWOORD

De tekst "Magnetic holes in the solar wind" van J.M. Turner, L.F. Burlaga, N.F. Ness, J. Lemaire, zal in Journal of Geophysical Research gepubliceerd worden.

VORWORT

Das Artikel "Magnetic holes in the solar wind" von J.M. Turner, L.F. Burlaga, N.F. Ness, J. Lemaire, wird in Journal of Geophysical Research veröffentlicht.

MAGNETIC HOLES IN THE SOLAR WIND

by

J.M. TURNER*, L.F. BURLAGA**, N.F. NESS** and J. LEMAIRE***

* *Morehouse College, Atlanta, Georgia 30314, USA.*

** *Goddard Space Flight Center, Greenbelt, Maryland 20771, USA.*

*** *Institut d'Aéronomie Spatiale de Belgique, B-1180 Brussels, Belgium.*

Abstract

An analysis of high resolution magnetic field measurements from the GSFC magnetometer on Explorer 43 showed that low magnetic field intensities ($< 1 \gamma$) in the solar wind at 1 AU occur as distinct depressions or "holes", in otherwise nearly average conditions. These magnetic holes are new kinetic-scale phenomena, having a characteristic dimension on the order of 20,000 km. They occur at a rate of 1.5/day in the 18-day interval (March 18 to April 6, 1971) that was considered. Most magnetic holes are characterized by both a depression in $|\vec{B}|$ and a change in the magnetic field direction, and some of these are possibly the result of magnetic merging. However, in other cases the direction does not change; such holes are not due to merging, but might be a diamagnetic effect due to localized plasma inhomogeneities.

Résumé

L'analyse des magnétogrammes à haute résolution obtenus à l'aide de Explorer 43 ont montré l'existence de dépressions dans l'intensité du champ magnétique interplanétaire. L'intensité minimum du champ magnétique au sein de ces "magnetic holes" est parfois inférieure à 1γ . Ces dépressions isolées de $|\vec{B}|$ ont des dimensions de l'ordre de 20.000 km. Elles sont observées à raison d'environ 1.5 par jour au cours de la période de 18 jours qui a été examinée. La plupart des "magnetic holes" sont caractérisés par une variation de $|\vec{B}|$ et par un changement dans la direction du champ magnétique. Il est suggéré que ces "magnetic holes" sont dus à des inhomogénéités dans le plasma du vent solaire produisant un effet diamagnétique détecté par les magnétomètres.

Samenvatting

De analyse van hoge resolutie - magnetogrammen, bekomen met behulp van de GSFC magnetometer in Explorer 43 toonde het bestaan aan van lage waarden van de intensiteit van het interplanetaire magneetveld. De minimale intensiteit van het magneetveld binnen zulke "magnetische gaten" is soms kleiner dan 1γ . Deze gelokaliseerde depressies hebben afmetingen van ongeveer 20.000 km. Gedurende de 18 daagse periode van 18 maart tot 6 april 1971, werden zij ongeveer 1.5 per dag waargenomen. De meeste van deze "magnetische gaten" gingen gepaard met een daling van de magnetische inductie en een omkeer in de zin van het magneetveld. Sommige van deze "magnetische gaten" zijn te wijten aan diamagnetische effecten die het gevolg zijn van gelokaliseerde plasma inhomogeniteiten.

Zusammenfassung

Eine Analyse der magnetischen Daten des GSFC Magnetometer der mit Explorer 43 geflogen ist, hat merkwürdige Vertiefungen in der Intensität des Interplanetaren Magnetischen Feldes erwiesen. Diese Vertiefungen die "magnetic holes" genannt worden sind, haben eine Breite von 20.000 km und sind zwischen 18 März und 6 April 1971, mit eine Frequenz von 1.5 pro Tag beobachtet worden. Für die meisten der "magnetic holes" wechselt nicht nur die Intensität aber auch in Richtung. Einige dieser "magnetic holes" sind wahrscheinlich durch "magnetic merging" gebildet. In anderen Fälle when die Richtung nicht ändert, sind diese "magnetic holes" durch diamagnetische Effekte erklärbar.

INTRODUCTION

Regions of very low intensity magnetic fields can be seen in high resolution measurements of the interplanetary magnetic field near 1 AU. We define low intensity by $|\vec{B}| < 1 \gamma$ which is to be compared with the average intensity of 5γ and the most probable value of 6γ . Most low field intensities were found to occur in isolated regions in the form of discrete "holes" imbedded in a background of otherwise uniform fields of nearly average intensity. The existence and the characteristics of these magnetic holes are the subjects of this paper.

Our analysis is based on ≈ 18 days of interplanetary data from Explorer 43 (IMP-I) in the period March 18 to April 9, 1971, the interval during which the GSFC plasma analyzer was operating. Low field regions ($|\vec{B}| < 1 \gamma$) were initially identified in plots of 15 s magnetic field averages. The "holes" thus found have a very small radial extent, but the high sampling rate of the magnetometer (12.5/s) resolved the structure of every event. The plasma sampling rate was much lower (a spectrum was measured in approximately one minute and successive spectra were obtained at four-minute intervals) and the structure of holes could not be resolved, but the plasma instrument did provide measurements of the pre- and post-hole states. The magnetic field and plasma experiments are described in reports by Fairfield (1973) and Ogilvie and Burlaga (1974), respectively.

RESULTS

We identified 28 magnetic "holes" using the criterion $|\vec{B}| < 1 \gamma$ and the data set discussed above. Typical examples are shown in Figure 1, which contains plots of magnetic field intensity for 10-min intervals. Nearly all holes are essentially isolated depressions in magnetic field intensity which is otherwise nearly average. They are distinct entities, not just random fluctuations in low intensity, disturbed field regions. Thus, the lowest magnetic field intensities in the solar wind near 1 AU, like the highest field intensities, apparently are the result of special physical processes distinct from those which produce the most probable fields.




Fig. 1.- Representative examples of magnetic holes. In each panel, 15 s averages of the magnetic field intensity are plotted versus time. The events are labeled by decimal day and universal time. Despite the variety of shapes and widths, there is a common characteristic, viz., a distinct depression to $|\vec{B}| < 1 \gamma$ in an otherwise normal magnetic field profile.

Given 28 holes in 18 days of data, one obtains an occurrence rate of 1.5/day. This is intermediate between the rate for shocks ($\approx .05/\text{day}$), (Chao and Lepping, 1974) and that for directional discontinuities ($\approx 25/\text{day}$) (Burlaga, 1972). The period used in this study is a representative solar wind state, in the sense that there were several well-defined streams (Burlaga and Ogilvie, 1973), 2 shocks (Ogilvie and Burlaga, 1974), and the types of "Alfven waves" that are often observed (Belcher and Davis, 1971, Burlaga and Turner, 1976). Thus, the rate of 1.5 holes per day, or ≈ 40 per solar rotation, is probably typical. Figure 2 shows the relations between the holes and the features just mentioned, and one can see that they are distributed fairly uniformly with respect to the streams, with perhaps some preference for the regions of decreasing speed. This gives us further reason to expect that the occurrence rate of $\approx 1.5/\text{day}$ is representative and not very strongly biased by conditions in our limited sample of data.

The "widths" of the holes ranged from ≈ 2 s to ≈ 130 s, with a median of 50 s. Since they are convected radially past the spacecraft at a speed on the order of 400 km/s, their thickness along the radial direction is on the order of 2×10^4 km, and since the proton Larmor radius near (but not in) the holes, R_L , is typically ≈ 100 km, the radial thickness of the holes is on the order of $200 R_L$. If the holes are field-aligned, the actual thickness is somewhat smaller, $\approx 150 R_L$. Because of their small size, magnetic holes are kinetic scale phenomena in the classification scheme of Burlaga (1969).

Turning now to the change in direction of the magnetic field across the holes, we find that it may change abruptly by a large amount, it may vary irregularly, or it may not change at all. Of the 28 events, 8 had little or no directional change, 9 were similar to D-sheets and 11 fell into neither of those categories. In the following, we shall discuss several examples of such changes. The plots to be presented are based on the high resolution data obtained at 12.5 samples/sec and are displayed in a coordinate system in which \hat{y} is the average field direction for 2 seconds before the event, \hat{z} is the direction of minimum variance for points in the interval during which the transition takes place, and \hat{x} is orthogonal to \hat{y} and \hat{z} and forms a right-handed coordinate system. It should be stressed that the coordinate system varies from event to event. In any case, however, tangential "discontinuities" in this system are indicated by $B_z = 0$.




Fig. 2.- Relation between magnetic holes and mesoscale interplanetary conditions.

An example in which the magnetic field direction changes abruptly across the holes is shown in Figure 3. The change in direction is centered about the time of minimum intensity, and B_z is essentially zero in the transition layer, indicating a tangential "discontinuity". The width of the March 27 event is 8 s, which is typical for directional discontinuities in the solar wind. The magnetic field direction changes by 180° in the March 27 event (Table I), and the magnetic field intensity drops to nearly zero, 0.12γ . In this respect, the structure resembles a D-sheet. Observations of D-sheets have been discussed by Burlaga (1968) and Burlaga and Scudder (1974) presented evidence that some D-sheets are the result of Sweet's mechanism, by which magnetic field is annihilated. The magnetic field intensity depressions in the D-sheets discussed heretofore are much broader than that in Figure 3 and occur much more infrequently than holes. For the March 27 event, the annihilation hypothesis predicts that the minimum intensity in the hole is $B_{\min} = 0.15\gamma$; this is in very good agreement with the observed value, 0.12γ . Unfortunately, the orientation is such that we cannot test for the subalfvenic streaming toward the current sheet which is predicted by Sweet's mechanism (see Burlaga and Scudder (1974) and references therein).

Another magnetic hole that resembles a D-sheet is the March 28, 1637 UT event, described in Table 1. In this case one can determine that the thickness is $20 R_L$. The observations suggest a subalfvenic streaming toward the current sheet ($V_o/V_A = 0.04$), where V_o is the flow speed normal to the current sheet and V_A is the Alfvén speed outside and adjacent to the current sheet. The value of B_{\min} predicted by the annihilation hypothesis is very close to the measured value (Table I).

Figure 4 shows an event that resembles a thin D-sheet, but which is not entirely consistent with the annihilation hypothesis. There is essentially no B_z component, indicating that the directional "discontinuity" is tangential. The "width" is only 8 sec, and the thickness along the normal (\hat{z}) direction is only $4 R_L$. The velocity measurements were not sufficiently accurate to determine whether or not there was a subalfvenic flow toward the current sheet. The important feature is that the observed minimum field is significantly smaller than that predicted by the merging model using the measured angular separation, ω , between the fields preceding and following the hole, \vec{B}_1 and \vec{B}_2 , respectively (see Table I).




Fig. 3.- A magnetic hole which might be the site of magnetic merging. The magnetic field direction rotates through 180° in a plane, and its intensity drops to nearly zero.




Fig. 4.- A magnetic hole which resembles a magnetic merging region but in which the minimum field intensity is lower than expected from the merging hypothesis.

TABLE I : Magnetic Holes that Resemble D-Sheets

| Event Time | ω | $\frac{B_{\min} \text{ (predicted)}}{B_{\min} \text{ (observed)}}$ | Thickness (R_L) |
|----------------|----------|--|------------------------|
| March 27, 0440 | 180° | 1.2 | — |
| March 28, 1637 | 129° | 1.2 | 20 |
| April 6, 1638 | 167° | 2.7 | 14 |
| April 1, 1025 | 131° | 7.1 | 4 |

Thus, either merging can operate in a way that is not understood or there is an entirely different process involved instead of or in addition to merging. There were other events which has minimum fields significantly smaller than predicted by the merging model (e.g., April 6, 1638 UT; in Table I). The events in this category had normals which were nearly radial. It should be noted that all of the holes discussed in this paper differ from the D-sheets discussed by Burlaga (1968) in that here the depression is confined to a region the size of that in which the direction changes, whereas it is much broader in D-sheets.

A distinctly different type of magnetic hole (which we call a linear hole) is shown in Figure 5; here is a smooth, symmetrical depression in magnetic field intensity, but no change in direction. The change is seen only in the B_y component, which is the average field direction, and in the intensity. Four such linear holes were found among the 28 events (see Table II). Their width along the radial direction is similar to that of other magnetic holes. Table II indicates that there was possibly a change in one or more of the plasma parameters across the linear holes, but more examples are needed before one can draw a general conclusion. Four other linear holes were identified with the same basic characteristics, but differed in that the field intensity did not vary smoothly in the hole. In these cases, the field intensity varied irregularly outside of the holes as well, and it is likely that the nonuniformity in the holes is due to external conditions. The occurrence of linear holes relative to streams is indicated by the 'L's in Figure 2.

Linear magnetic holes are certainly not produced by a merging process, since a change in the direction of \vec{B} is a necessary signature for merging. A possible explanation is that they are diamagnetic responses to localized plasma inhomogeneities. Indeed, high values of $\beta \equiv nkT/(B^2/8\pi)$ (where n , T are the density and temperature, respectively, of the protons, and B is the magnetic field intensity) were observed adjacent to the holes (see Table II). One can model linear holes using the theory for diamagnetic boundary layers developed by Sestero (1964) and Lemaire and Burlaga (1976). In particular, one can regard a hole as two adjacent boundary layers across which $|\vec{B}|$ changes. In one layer, from $z \rightarrow -\infty$ to z_0 where $|\vec{B}|$ is a minimum, the magnetic field intensity decreases; in the other layer, from z_0 to $z \rightarrow +\infty$, the magnetic field intensity increases. The model implies a localized plasma inhomogeneity




Fig. 5.- A linear magnetic hole. In this case, the hole is certainly not the result of merging. This, and perhaps all magnetic holes might be a diamagnetic effect due to a localized plasma inhomogeneity.

TABLE II : Linear Magnetic Holes
 (1 = pre event, 2 = post event)

| EVENT | (n_1/n_2) | (T_1/T_2) | (V_1/V_2) | (β_1) | (β_2) | ω |
|-------------------|-------------|-------------|-------------|-------------|-------------|----------|
| March 23, 552 | .92 | .86 | 1.01 | 3.80 | 2.02 | 6° |
| March 24, 1607 | 1.15 | .90 | 1.03 | 1.38 | 1.18 | 17° |
| March 24, 1633 | 1.03 | 1.02 | 1.02 | 1.11 | 1.02 | 5° |
| March 27, 1823 | 1.11 | 1.09 | .97 | .40 | .24 | 13° |

(dense and/or hot plasma) which "excludes" the magnetic field. An electric field is set up along the "normal" to the current sheet and particles drift in this field and in the gradient of $|\vec{B}|$, thereby providing the current which maintains the structure in a steady state. This model allows magnetic field enhancements as well as holes, if the plasma inhomogeneity is due to a decrease in density and/or temperature. An observation of such an event is shown in Figure 6. We did not attempt to study the statistics of such events. Of course, the application of this model to magnetic holes is only speculative, and it doesn't explain the origin of the plasma inhomogeneities. High resolution plasma measurements are needed to understand the true nature and origin of magnetic holes. Multispacecraft measurements are needed to determine their spatial structure and to follow their evolution.

SUMMARY AND DISCUSSION

In the high time resolution data from Explorer 43 (IMP-I), we found that the lowest magnetic field intensities ($< 1\gamma$) in the solar wind at 1 AU nearly always occur as distinct depressions or "holes" in otherwise nearly average interplanetary magnetic fields. These magnetic holes are new kinetic scale phenomena, convecting past a fixed spacecraft in some tens of seconds and having dimensions on the order of tens of proton Larmor radii. They occur at a rate of 1.5/day during the 18-day period which was considered, a rate intermediate between that of shocks and that of directional discontinuities.

The direction of \vec{B} changes across most magnetic holes, much as it does in the current sheets associated with directional discontinuities with no change in $|\vec{B}|$, i.e., it rotates in a plane and has a thickness of several proton gyroradii. However, there are some magnetic holes at which there is virtually no change in the direction of \vec{B} . Some of the directional holes resemble D-sheets, although there is an important difference in that the depression in $|\vec{B}|$ has the same dimension as the change in direction at holes whereas it is much broader than the change in direction at D-sheets. In particular, some holes are possibly the result of magnetic merging. However, the linear holes are certainly not the result of merging, which requires a change in the direction of \vec{B} . These linear holes (and perhaps all holes) are possibly diamagnetic effects due to the presence of localized plasma inhomogeneities, but

IMPI
23 MARCH 71




Fig. 6.- The antithesis of a magnetic hole. Here the magnetic field intensity increases in a 7 s. interval while the direction remains constant. Presumably the plasma pressure was low during the magnetic field enhancement.

we can neither observe such small inhomogeneities because of the low plasma data sampling rates nor offer an unambiguous explanation for their origin.

ACKNOWLEDGMENTS

Dr. Fairfield, a Co-Investigator on the Explorer 43 magnetic field experiment, was probably the first person to notice linear magnetic holes. He provided some plots of the magnetic field data and critically examined a draft of this paper. The plasma data were kindly provided by Dr. K.W. Ogilvie. One of us (J.L.) was supported by a National Academy of Sciences/National Research Council Resident Research Associateship. He thanks N. Ness and K. Ogilvie for their hospitality and support at the Laboratory for Extraterrestrial Physics.

REFERENCES

- BELCHER, J.W. and L. DAVIS, Jr., Large amplitude Alfvén waves in the interplanetary medium, 2, *J. Geophys. Res.*, **76**, 3534, 1971.
- BURLAGA, L.F., Microscale structures in the interplanetary medium, *Solar Physics*, **4**, 67, 1968.
- BURLAGA, L.F., Directional discontinuities in the interplanetary magnetic field, *Solar Physics*, **7**, 54, 1969.
- BURLAGA, L.F., Microstructure of the interplanetary medium in *The Solar Wind*, ed. C.P. Sonett, P.J. Coleman, Jr., and J.M. Wilcox, NASA SP-308, 1972.
- BURLAGA, L.F. and K.W. OGILVIE, Solar wind temperature and speed, *J. Geophys. Res.*, **78**, 2028, 1973.
- BURLAGA, L.F. and J.D. SCUDDER, Sweet's mechanism in the solar wind, *Astrophys. J. (Letters)*, **191**, 1149, 1974.
- BURLAGA, L.F. and J.M. TURNER, "Alfvén" waves in the solar wind, *J. Geophys. Res.*, **81**, 73, 1976.
- CHAO, J.K. and R.P. LEPPING, A correlative study of SSC's interplanetary shocks, and solar activity, *J. Geophys. Res.*, **79**, 1799, 1974.
- FAIRFIELD, D.H., Whistler waves observed upstream from collisionless shocks, *J. Geophys. Res.*, **79**, 1368, 1973.
- LEMAIRE, J. and L.F. BURLAGA, Diamagnetic boundary layers in the solar wind : kinetic theory, submitted to *Astrophys. and Space Sci.*, 1976.
- OGILVIE, K.W. and L.F. BURLAGA, A discussion of interplanetary shock waves, with two examples, *J. Geophys. Res.*, **79**, 2324, 1974.
- SESTERO, A., Structure of plasma sheaths, *Physics of Fluids*, **1**, 44, 1964.

- 100 - BIAUME, F., Détermination de la valeur absolue de l'absorption dans les bandes du système de Schumann-Runge de l'oxygène moléculaire, 1972.
- 101 - NICOLET, M. and W. PEETERMANS, The production of nitric oxide in the stratosphere by oxidations of nitrous oxide, 1972.
- 102 - VAN HEMELRIJCK, E. et H. DEBEHOGNE, Observations au Portugal de phénomènes lumineux se rapportant à une expérience de lâcher de barium dans la magnétosphère, 1972.
- 103 - NICOLET, M. et W. PEETERMANS, On the vertical distribution of carbon monoxide and methane in the stratosphere, 1972.
- 104 - KOCKARTS, G., Heat balance and thermal conduction, 1972.
- 105 - ACKERMAN, M. and C. MULLER, Stratospheric methane from infrared spectra, 1972.
- 106 - ACKERMAN, M. and C. MULLER, Stratospheric nitrogen dioxide from infrared absorption spectra, 1972.
- 107 - KOCKARTS, G., Absorption par l'oxygène moléculaire dans les bandes de Schumann-Runge, 1972.
- 108 - LEMAIRE, J. et M. SCHERER, Comportements asymptotiques d'un modèle cinétique du vent solaire, 1972.
- 109 - LEMAIRE, J. and M. SCHERER, Plasma sheet particle precipitation : A kinetic model, 1972.
- 110 - BRASSEUR, G. and S. CIESLIK, On the behavior of nitrogen oxides in the stratosphere, 1972.
- 111 - ACKERMAN, M. and P. SIMON, Rocket measurement of solar fluxes at 1216 Å, 1450 Å and 1710 Å, 1972.
- 112 - CIESLIK, S. and M. NICOLET, The aeronomic dissociation of nitric oxide, 1973.
- 113 - BRASSEUR, G. and M. NICOLET, Chemospheric processes of nitric oxide in the mesosphere and stratosphere, 1973.
- 114 - CIESLIK, S. et C. MULLER, Absorption raie par raie dans la bande fondamentale infrarouge du monoxyde d'azote, 1973.
- 115 - LEMAIRE, J. and M. SCHERER, Kinetic models of the solar and polar winds, 1973.
- 116 - NICOLET, M., La biosphère au service de l'atmosphère, 1973.
- 117 - BIAUME, F., Nitric acid vapor absorption cross section spectrum and its photodissociation in the stratosphere, 1973.
- 118 - BRASSEUR, G., Chemical kinetic in the stratosphere, 1973.
- 119 - KOCKARTS, G., Helium in the terrestrial atmosphere, 1973.
- 120 - ACKERMAN, M., J.C. FONTANELLA, D. FRIMOUT, A. GIRARD, L. GRAMONT, N. LOUISNARD, C. MULLER and D. NEVEJANS, Recent stratospheric spectra of NO and NO₂, 1973.
- 121 - NICOLET, M., An overview of aeronomic processes in the stratosphere and mesosphere, 1973.
- 122 - LEMAIRE, J., The "Roche-Limit" of ionospheric plasma and the formation of the plasmopause, 1973.
- 123 - SIMON, P., Balloon measurements of solar fluxes between 1960 Å and 2300 Å, 1974.
- 124 - ARJIS, E., Effusion of ions through small holes, 1974.
- 125 - NICOLET, M., Aéronomie, 1974.
- 126 - SIMON, P., Observation de l'absorption du rayonnement ultraviolet solaire par ballons stratosphériques, 1974.
- 127 - VERCHEVAL, J., Contribution à l'étude de l'atmosphère terrestre supérieure à partir de l'analyse orbitale des satellites, 1973.
- 128 - LEMAIRE, J. and M. SCHERER, Exospheric models of the topside ionosphere, 1974.
- 129 - ACKERMAN, M., Stratospheric water vapor from high resolution infrared spectra, 1974.
- 130 - ROTH, M., Generalized invariant for a charged particle interacting with a linearly polarized hydromagnetic plane wave, 1974.
- 131 - BOLIN, R.C., D. FRIMOUT and C.F. LILLIE, Absolute flux measurements in the rocket ultraviolet, 1974.
- 132 - MAIGNAN, M. et C. MULLER, Méthodes de calcul de spectres stratosphériques d'absorption infrarouge, 1974.
- 133 - ACKERMAN, M., J.C. FONTANELLA, D. FRIMOUT, A. GIRARD, N. LOUISNARD and C. MULLER, Simultaneous measurements of NO and NO₂ in the stratosphere, 1974.

- 134 - NICOLET, M., On the production of nitric oxide by cosmic rays in the mesosphere and stratosphere, 1974.
- 135 - LEMAIRE, J. and M. SCHERER, Ionosphere-plasmasheet field aligned currents and parallel electric fields, 1974.
- 136 - ACKERMAN, M., P. SIMON, U. von ZAHN and U. LAUX, Simultaneous upper air composition measurements by means of UV monochromator and mass spectrometer, 1974.
- 137 - KOCKARTS, G., Neutral atmosphere modeling, 1974.
- 138 - BARLIER, F., P. BAUER, C. JAECK, G. THULLIER and G. KOCKARTS, North-South asymmetries in the thermosphere during the last maximum of the solar cycle, 1974.
- 139 - ROTH, M., The effects of field aligned ionization models on the electron densities and total flux tubes contents deduced by the method of whistler analysis, 1974.
- 140 - DA MATA, L., La transition de l'homosphère à l'hétérosphère de l'atmosphère terrestre, 1974.
- 141 - LEMAIRE, J. and R.J. HOCH, Stable auroral red arcs and their importance for the physics of the plasmopause region, 1975.
- 142 - ACKERMAN, M., NO, NO₂ and HNO₃ below 35 km in the atmosphere, 1975.
- 143 - LEMAIRE, J., The mechanisms of formation of the plasmopause, 1975.
- 144 - SCIALOM, G., C. TAIEB and G. KOCKARTS, Daytime valley in the F1 region observed by incoherent scatter, 1975.
- 145 - SIMON, P., Nouvelles mesures de l'ultraviolet solaire dans la stratosphère, 1975.
- 146 - BRASSEUR, G. et M. BERTIN, Un modèle bi-dimensionnel de la stratosphère, 1975.
- 147 - LEMAIRE, J. et M. SCHERER, Contribution à l'étude des ions dans l'ionosphère polaire, 1975.
- 148 - DEBEHOGNE, H. et E. VAN HEMELRIJCK, Etude par étoiles-tests de la réduction des clichés pris au moyen de la caméra de triangulation IAS, 1975.
- 149 - DEBEHOGNE, H. et E. VAN HEMELRIJCK, Méthode des moindres carrés appliquée à la réduction des clichés astrométriques, 1975.
- 150 - DEBEHOGNE, H. et E. VAN HEMELRIJCK, Contribution au problème de l'aberration différentielle, 1975.
- 151 - MULLER, C. and A.J. SAUVAL, The CO fundamental bands in the solar spectrum, 1975.
- 152 - VERCHEVAL, J., Un effet géomagnétique dans la thermosphère moyenne, 1975.
- 153 - AMAYENC, P., D. ALCAYDE and G. KOCKARTS, Solar extreme ultraviolet heating and dynamical processes in the mid-latitude thermosphere, 1975.
- 154 - ARIJS, E. and D. NEVEJANS, A programmable control unit for a balloon borne quadrupole mass spectrometer, 1975.
- 155 - VERCHEVAL, J., Variations of exospheric temperature and atmospheric composition between 150 and 1100 km in relation to the semi-annual effect, 1975.
- 156 - NICOLET, M., Stratospheric Ozone : An introduction to its study, 1975.
- 157 - WEILL, G., J. CHRISTOPHE, C. LIPPENS, M. ACKERMAN and Y. SAHAI, Stratospheric balloon observations of the southern intertropical arc of airglow in the southern american area, 1976.
- 158 - ACKERMAN, M., D. FRIMOUT, M. GOTTIGNIES, C. MULLER, Stratospheric HCl from infrared spectra, 1976.
- 159 - NICOLET, M., Conscience scientifique face à l'environnement atmosphérique, 1976.
- 160 - KOCKARTS, G., Absorption and photodissociation in the Schumann-Runge bands of molecular oxygen in the terrestrial atmosphere, 1976.
- 161 - LEMAIRE, J., Steady state plasmopause positions deduced from McIlwain's electric field models, 1976.
- 162 - ROTH, M., The plasmopause as a plasma sheath : A minimum thickness, 1976.
- 163 - FRIMOUT, D., C. LIPPENS, P.C. SIMON, E. VAN HEMELRIJCK, E. VAN RANSBEECK et A. REHRI, Lâchers de monoxyde d'azote entre 80 et 105 km d'altitude. Description des charges utiles et des moyens d'observation, 1976.
- 164 - LEMAIRE, J. and L.F. BURLAGA, Diamagnetic boundary layers : a kinetic theory, 1976.