

I N S T I T U T D ' A E R O N O M I E S P A T I A L E D E B E L G I Q U E

3 - Avenue Circulaire  
B - 1180 BRUXELLES

# AERONOMICA ACTA

A - N° 279 - 1984

Photodissociation effects of solar UV radiation

by

P.C. SIMON and G. BRASSEUR

B E L G I S C H I N S T I T U U T V O O R R U I M T E - A E R O N O M I E

3 - Ringlaan  
B - 1180 BRUSSEL

## FOREWORD

This paper has been presented at the European Geophysical Society (EGS), Meeting Structure of the Upper and Middle Atmosphere in Leeds, 23-27 August, 1982, held in honour of Prof. Marcel Nicolet.

## AVANT-PROPOS

Ce travail a été présenté à la réunion - Structure of the Upper and Middle Atmosphere - du "European Geophysical Society" (EGS), organisée à Leeds, 23-27 août 1982, en l'honneur du Prof. Marcel Nicolet. Il sera publié dans "Planetary and Space Science".

## VOORWOORD

Dit werk werd voorgesteld tijdens de vergadering - Structure of the Upper and Middle Atmosphere - van de "European Geophysical Society" (EGS), gehouden te Leeds, 23-27 augustus 1982, ter ere van Prof. Marcel Nicolet. Het zal verschijnen in "Planetary and Space Science".

## VORWORT

Diese Arbeit wurde präsentiert während der "European Geophysical Society" (EGS) Tagung - Structure of the Upper and Middle Atmosphere - organisiert in Leeds, 23-27 August 1982, zu Ehren des Professors Marcel Nicolet. Diese Arbeit wird in "Planetary and Space Science" herausgegeben werden.

# PHOTODISSOCIATION EFFECTS OF SOLAR UV RADIATION

by P.C. SIMON and G. BRASSEUR\*

Institut d'Aéronomie Spatiale de Belgique  
3, Avenue Circulaire, B-1180 Bruxelles - Belgium

## Abstract

This paper reviews the current knowledge of ultraviolet solar irradiances beyond 120 nm measured during solar cycle 21. Parameters required to calculate the atmospheric attenuation of the UV radiation and the photodissociation rate of neutral constituents are discussed. The effect of scattered sunlight is also considered.

## Résumé

Ce travail fait le point sur la connaissance actuelle de l'éclairement solaire dans l'ultraviolet au-delà de 120 nm, mesuré durant le cycle solaire 21. Les paramètres, nécessaires pour calculer l'atténuation atmosphérique de la radiation UV et le taux de photodissociation des éléments composants neutres, sont discutés. L'effet de la diffusion du rayonnement solaire est également pris en considération.

---

\* Aspirant au Fonds National de la Recherche Scientifique

## Samenvatting

Dit werk bekijkt de huidige kennis van de zonnestraling in het ultraviolet boven 120 nm, gemeten tijdens zonnecyclus 21. De parameters, nodig om de atmosferische verzwakking van de UV straling en de fotodissociatiegraad van de neutrale componenten te berekenen, worden besproken. Het verstrooiingseffekt van de zonnestraling wordt eveneens onderzocht.

## Zusammenfassung

Diese Arbeit erwägt die heutige Kenntnis über die Sonnenausstrahlung im Ultraviolett über 120 nm, gemessen während des Sonnenzykluses 21. Die Parameter, nötig um die atmosphärische Schwächung der UV Sonnenstrahlen und der Photodissoziationsgrad von den neutralen Komponenten zu berechnen, werden besprochen. Der Diffusionseffekt der Sonnenausstrahlung wird auch untersucht.

## 1. INTRODUCTION

The chemical and the thermal structure of planetary atmospheres is driven by the radiation emitted by the Sun and its absorption by optically active atmospheric constituents. An accurate knowledge of the solar irradiance and its variation is therefore fundamental in aeronomy and in climatology. Most of the solar energy is confined in the wavelength range below 4000 nm while the radiation emitted by the Earth and the surrounding atmosphere has a wavelength greater than 4000 nm.

Photochemical processes of the neutral constituents in the middle atmosphere and in the lower thermosphere are initiated by ultraviolet radiation in the wavelength interval ranging from 120 nm to the visible part of the spectrum. Furthermore the H $\gamma$  Lyman  $\alpha$  line is responsible for the ionization of nitric oxide (Nicolet, 1945) leading to the formation of the D region.

When penetrating in the atmospheric layers, the solar irradiance is progressively attenuated by absorption and scattering. The main absorbing gases are molecular oxygen and ozone. An accurate analysis of the radiative transfer in the ultraviolet requires therefore a perfect knowledge of (1) the solar irradiance at the top of the atmosphere, (2) the absorption cross sections of O<sub>2</sub> and O<sub>3</sub>, (3) the integrated amount of these absorbing species (4) the scattering cross section and (5) the albedo.

The distribution of the trace species concentration depends directly on the photodissociation rates in the atmosphere. The photodissociation frequency of a molecule X is given by

$$J(X; z, \chi) = \int_{\lambda} \varepsilon(X; \lambda) \sigma(X; \lambda) q(\lambda, z, \chi) d\lambda \quad (1)$$

where  $q(\lambda, z, \chi)$  is the total photon flux into a volume element at an altitude  $z$ , for a solar zenith angle  $\chi$  and for wavelength  $\lambda$ .  $\sigma(X, \lambda)$  is the absorption cross section and  $\varepsilon(X, \lambda)$  the photodissociation efficiency for the given molecule. The absorption and scattering processes will therefore influence the atmospheric composition.

The purpose of this paper is to review the key parameters which are used as input data in photochemical models of the atmosphere. In particular, the solar irradiance, the absorption cross section of molecular oxygen and ozone and the Rayleigh scattering will be discussed. In order to clarify the analysis, the solar spectrum will be divided according to the various spectral regions of molecular oxygen corresponding to the penetration of solar radiation in different atmospheric layers.

## 2. SOLAR IRRADIANCE

Solar ultraviolet irradiance measurements from 135 to 400 nm will be reviewed. The solar H $\alpha$  Lyman  $\alpha$  line will not be discussed here since its absolute value and its variations are analyzed by Bossy (1983, this volume). On the other hand, measurements obtained since the first observation reported by Friedman et al. (1951) have been discussed in several papers (e.g. Vidal-Madjar, 1977; Simon 1978; Simon 1981).

### 2.1. The wavelength interval 135-175 nm

During solar cycle 20, several rocket measurements have been made in the wavelength interval corresponding to the Schumann-Runge continuum of molecular oxygen. Heroux and Swirbalus (1976) and Samain and Simon (1976) have published the lowest irradiance values. The latter values refer to a quiet solar full disk. Rottman (1981) has proposed revised values for his observations performed in 1972 and 1973. With three other flights made in 1975, 1976 and 1977, he has

defined a so-called reference spectrum corresponding to minimum condition between solar cycle 20 and 21, by taking an average value provided by the five flights.

The irradiance values obtained during cycle 21 for a high solar activity level have been published by Mount et al. (1980) and Mount and Rottman (1981). They are systematically higher than the reference spectrum of Rottman (1981). The differences represent long-term variability which is confirmed by the AE-E satellite observations reported by Hinteregger (1981) even if the latter correspond to a limited field of view on the solar disk of 6 arc min x 6 arc min situated near the disk center. New data obtained in 1982 have been very recently published by Mount and Rottman (1983). These observations, corresponding to an intermediate level of solar activity are comparable to the minimum values proposed by Rottman (1981). They are even lower than the values obtained in 1976 and 1977 with a similar instrument. Values obtained by Heath (1980) during the rising phase of solar cycle 21 in November 1978 are also in better agreement with the minimum values proposed by Rottman (1981) rather than with those obtained during the maximum of solar cycle 21.

On the basis of the former observations performed since 1978, the enhancement of solar irradiance with the 11-year solar activity seems correlated only with the highest activity levels around the cycle maximum in 1979 and 1980. This is in contradiction with the two-component plage model of Cook et al. (1980) which proposes a linear relationship between the solar irradiance in the 117.5-210 nm wavelength interval and the Zurich sunspot number. A three-component model recently published by Lean et al. (1982) and based on spatially resolved observations of the Ca II K chromosphere is in better agreement with the 11-year cycle variation deduced from the LASP data (Rottman, 1981; Mount et al., 1980; Mount and Rottman, 1981). According to this latter model, the irradiance increase in the 155-160 nm wavelength interval

reaches levels higher than 25 percent after the beginning of 1978 or before mid 1971 corresponding to smooth sunspot numbers larger than 80.

The ratios of irradiance values integrated over 5 nm from 135 to 175 nm taking as a reference the irradiance values proposed by Brasseur and Simon (1981) are given in figure 1. This graph clearly shows that the previous values of Heroux and Swirbalus (1976) are 30 percent lower than the reference spectrum. On the other hand, the divergences between the five observations of Rottman (1981) give a good idea of the reproductibility reached in this wavelength range by means of snapshot measurements which include the variability related to the 27-day solar rotational period.

Observations made by the Solar Mesosphere Explorer (SME) indicate a 27 day variability of the order of 12 percent in the integrated 130-175 nm wavelength range (Rottman et al., 1982). A detailed discussion of this question is given by Rottman (1983, this volume).

Accuracy and precision of the irradiance observations need to be improved in order to quantitatively measure the solar variability during the 11-year cycle. The accuracy of  $\pm 8$  percent at the two  $\sigma$  level quoted by Mount and Rottman (1983) in the wavelength interval of 115-186.5 nm seems very promising in regard to the previous observations usually made with an accuracy of  $\pm 15$  percent at the one  $\sigma$  level. This improvement has been reached by using the Synchrotron Users Radiation Facility (SURF) at the National Bureau of Standards as a calibration standard.

## 2.2. The wavelength interval 175-210 nm

Solar irradiance and its variability are very poorly known in the wavelength range corresponding to the Schumann-Runge bands of mole-






Figure 1 Observed solar irradiance in the spectral range 135-210 nm expressed as ratios relative to the solar reference spectrum of Brasseur and Simon (1981). Values are integrated over 5 nm intervals.

cular oxygen. Figure 1 gives the ratios of several observed irradiance values integrated over 5 nm taking as a reference the values obtained by Samain and Simon (1976) corresponding to a quiet sun and quoted by Brasseur and Simon (1981). Differences at wavelengths below 185 could probably be due to long-term solar variability. Divergences between individual observations are not uniform with wavelength. It should be mentioned that, according to the two-component model of Cook et al. (1980) and the three-component model of Lean et al. (1982), the variability during the rising phase of solar cycle 21 beyond 182 nm does not exceed 30 percent. Ratios from 180 to 190 nm obtained on the basis of Rottman's measurement should be excluded because of larger errors. In addition, the data of Mount et al. (1980) are not reliable beyond 190 nm because of experimental problems (Mount and Rottman, 1981). Consequently, the quantitative variation over the 11-year cycle is still masked by the uncertainties of all measurements in this wavelength range. Additional observations with accuracy and precision higher than or at least comparable to those obtained by Mount and Rottman (1983) are required.

The variability of the solar irradiance with the 27 day rotation period has been observed by Nimbus 7 (Heath, 1980) and by SME (Rottman et al., 1982). Heath (1980) observes a maximum variability of 5 percent around 180 nm while Rottman et al. report a variability of 10 percent for the whole spectral interval of 175 to 190 nm.

### 2.3. The spectral region beyond 210 nm

Figure 2a presents the ratio of irradiance values integrated over 5 nm from 205 to 270 nm taking as a reference the values proposed by Brasseur and Simon (1981). Systematic divergences clearly appear between most measurements in the 205-240 wavelength interval. They are probably due to experimental errors. Differences between all the flights do not exceed 20 percent. They are within 12 percent when

considering only the balloon observations reported by Simon et al. (1982a,b) and within 10 percent when keeping only the 1980 and 1982 rocket flight results published by Mount and Rottman (1981, 1983). All these measurements confirm that the values of Broadfoot (1972) have to be lowered by at least 25 percent below 260 nm. The lowest values of Mount et al. (1980) are not reliable in this wavelegnth range because of experimental problems encountered during this flight (Mount and Rottman, 1981). Above 240 nm, the divergences are within 15 percent. Data reported by Mount and Rottman (1981) have been revised on the basis of improved calibration methods obtained with the Synchrotron Users Radiation Facility (SURF) at the National Bureau of Standard (see Mount and Rottman, 1983).

Figure 2b represents the same comparison for integrated irradiance values over 5 nm between 270 and 350 nm. The agreement between all the observations is good ( $\pm 10$  percent) up to 295 nm. Beyond this wavelength, the data obtained in 1980 and revised by Mount and Rottman (1983), as well as those of Broadfoot (1972), become significantly lower than the others. Balloon observations also exhibit larger discrepancies up to 15 percent, at larger wavelengths. It should be noted that the quantum yield of the solar blind detectors used for rocket and balloon observations decreases rapidly beyond 250 nm and becomes very small for wavelengths greater than 300 nm. Nevertheless, the agreement with the values obtained by Heath (1980) is within 5 percent for the 1979 balloon observation (Simon et al., 1982b).

Beyond 330 nm, new irradiance values have been published by Neckel and Labs (1981). These latter values which supersede those previously published by Labs and Neckel (1970) are probably the most reliable in the near ultraviolet and in the visible. According to Table 3 in Simon (1981), they are in good agreement with the observation reported by Heath (1980). The measurements of Arvesen et al. (1969) have to be carefully adjusted because of errors in their wavelength




Figure 2 Observed solar irradiance in the spectral ranges 205-275 nm (bottom) and 270-340 nm (top) expressed as ratios relative to the solar reference spectrum of Brasseur and Simon (1981). Values are integrated over 5 nm intervals. The labeling is as follows : B 70 : Broadfoot (1972); S 76 : Simon et al. (1982a) flight dated July 1, 1976.

scale below 400 nm and of uncertainties due to changes in the spectral irradiance scale of the National Bureau of Standards in 1973.

Long-term variability in this spectral range is generally considered as being negligible from the aeronomic point of view and is also masked by the uncertainties in available measurements. Variations less than 2 percent related with the 27-day rotational period of the sun have been reported by Heath (1980) from Nimbus 7 observations. They are confirmed by the recent SME observations (Rottman, 1983).

#### 2.4. Summary

In conclusion, solar irradiance values are probably known with an uncertainty of  $\pm 20$  percent from 135 to 175 nm for solar minimum condition. The long term variations are certain, reaching a factor of two around 150 nm between the maximum and the minimum of activity during the rising phase of solar cycle 21. In the wavelength interval of 175-200 nm, the uncertainty is not smaller ( $\pm 20$  percent) and the long-term variations are still masked by the accuracy and the precision of available observations. Solar irradiances can be estimated within  $\pm 15$  percent of accuracy around 210 nm and with  $\pm 5$  percent at 300 nm. Further observations with an accuracy of  $\pm 5$  percent and a precision better than  $\pm 2$  percent are still needed in order to provide useful irradiance values and to determine the possible long-term variability beyond 175 nm. Such goals are only achievable by referring the calibration to a common stable source as the synchrotron to insure the continuity of future observations. It is also required to compare the different instrument performances when observing the sun from the same platform. Intercomparison of solar spectrometers should make future measurements on board balloon, rocket and space shuttle more comparable and should help to the validation of satellite observations, particularly for the instrument drift in orbit.

### 3. THE ABSORPTION CROSS SECTION OF MOLECULAR OXYGEN

#### 3.1. The Lyman $\alpha$ window region

Below 130 nm, the absorption cross section of molecular oxygen varies rapidly with wavelength. Near the H $\gamma$  Lyman  $\alpha$  solar line, the minimum values of the oxygen cross-section allows the solar radiation to reach the mesosphere, down to the altitude of 70 km. Since this solar line is responsible for the dissociation of water vapor, methane, carbon dioxide and the ionization of nitric oxide, an accurate knowledge of the optical depth in this wavelength region is required.

Many laboratory measurements have been made (e.g. Watanabe, 1958; Ogawa, 1968; Dose et al., 1975) leading to the working value of  $1 \times 10^{-20}$  cm<sup>2</sup> at 121.6 nm. Hall (1972) has pointed out that the variation in the molecular oxygen cross section across the solar Lyman  $\alpha$  line which has the appearance of a doublet (Purcell and Tousey, 1960) requires the calculation of an effective cross section dependent of the oxygen content above a given altitude. On the other hand, Smith and Miller (1974) and Weeks (1975) have pointed out the temperature dependence of the molecular oxygen absorption cross section near Lyman  $\alpha$ . Carver et al. (1977) have measured the oxygen absorption cross sections at 0.02 nm intervals between 121.4 and 121.86 nm at three temperatures of 294, 195 and 82 K. The effective cross sections have been calculated as a function of the integrated Lyman  $\alpha$  extinction for the three temperatures. Results are significantly lower than the effective cross sections published by Hall (1972) and based on the Ogawa's measurements.

Nicolet and Peetermans (1980) using the data of Carver et al. (1977), have deduced simple expressions, adopting an average temperature of 190 K, for the photodissociation rate calculations.

The optical depth is given by the expression :

$$\tau_{\text{Ly } \alpha}(\text{O}_2) = 4.17 \times 10^{-19} N(\text{O}_2)^{0.917} \quad (2a)$$

for  $N(\text{O}_2) > 1 \times 10^{19}$  molecules/cm<sup>2</sup> corresponding to an effective cross section given by :

$$\sigma_{\text{Ly } \alpha}(\text{O}_2) = 4.17 \times 10^{-19} N(\text{O}_2)^{-0.083} \quad (2b)$$

For  $N(\text{O}_2) \leq 1 \times 10^{19}$  molecules/cm<sup>2</sup>, a cross section value of  $1 \times 10^{-20}$  cm<sup>2</sup> can be adopted.

### 3.2. The Schumann-Runge continuum

The heating in the lower thermosphere as well as the formation of the atomic oxygen due to the photodissociation of molecular oxygen in this atmospheric layer has to be attributed to the absorption of solar radiation in the 135-175 nm wavelength interval.

The absorption cross section of molecular oxygen in this range has been reviewed several times. The first analysis was made by Nicolet and Mange (1954) and followed by those of Watanabe (1958), Ackerman (1971) and Hudson (1971). The latter author has also extensively discussed the sources of errors in the ultraviolet absorption cross section measurements. He claimed that the total uncertainty with the laboratory techniques available at that time, is larger than 5 percent. Furthermore, errors on the cross section values are larger when photographic emulsions are used instead of photoelectric techniques detection. Actually, the first investigations of molecular oxygen absorption cross section have been conducted using photographic techniques and yielded values 20-25 percent higher than those obtained by photoelectric measurements. Later on, the data obtained by Goldstein and Mastrup (1966) did not confirm the higher cross section values obtained by

photographic emulsions. Figure 3 compares several photoelectric measurements between 135 and 160 nm obtained since 1953 at room temperature with those of Goldstein and Mastrup (1966). The divergences between the photoelectric observations do not exceed 10 percent and are therefore within the  $\pm 10$  percent of accuracy quoted by most investigators. Specific measurements have been performed only around the peak of the absorption spectrum in order to obtain reliable cross section values of the order of  $1.5 \times 10^{-17} \text{ cm}^2$  in this region (e.g., Watanabe and Marno, 1956; Huffman et al., 1964). Measurements performed between 160 and 180 nm are displayed in figure 4. The temperature dependence of the absorption cross section is illustrated by the data obtained around 300 and 600 K by Hudson et al. (1966) and by Lean and Blake (1981). The positive temperature effect becomes significant for wavelengths greater than 160 nm and should be taken into account in photodissociation calculations in the thermosphere. According to Blake et al. (1980), the temperature coefficient becomes negative at shorter wavelengths but the change is limited to less than - 4 percent at 149 nm instead the - 10 percent at 160 nm as measured by Hudson et al. (1966). A semi-empirical two terms model of the temperature dependence of the molecular oxygen absorption cross section is proposed by Lean and Blake (1981).

The Schumann-Runge continuum adjoins the Schumann-Runge band system at 175 nm and underlies the bands below about 183 nm. Therefore, this continuum contributes to the absorption cross section minima in the band system and should be taken into account in cross section calculations discussed in the following section.

### 3.3. The Schumann-Runge band system

The attenuation of the solar radiation in the wavelength range of 175-200 nm is driven by the absorption of the molecular oxygen in the Schumann-Runge band system. The absorption takes place mainly in the






Figure 3 Absorption cross section of molecular oxygen between 135 and 160 nm.




Figure 4 Absorption cross section of molecular oxygen between 160 and 185 nm.

mesosphere, between the altitudes of 60 and 90 km. However the penetration is large enough so that the radiation in this spectral range contributes significantly to the photodissociation of several constituents in the stratosphere.

Several studies of the photodissociation of molecular oxygen in the middle atmosphere have been performed by many authors such as Kockarts (1971), Hudson and Mahle (1972), Fang et al. (1974), Park (1974), Kockarts (1976), Blake (1979), Nicolet and Peetermans (1980) and Frederick and Hudson (1979).

In situ observations at 40 km altitude of solar irradiance between 184 and 202 nm have been reported by Frederick et al. (1981) and have provided a check on the absorption cross sections of molecular oxygen. The differences with detailed attenuation calculations are of the order of 10 percent above 194 nm, 20 percent between 189 and 194 nm and as large as 40 percent below 189 nm. According to Frederick et al. (1981), such disagreements could be attributed to uncertainties in spectroscopic data but errors associated with the observations should not be neglected.

The cross section values in the Schumann-Runge band system which vary by about 5 orders of magnitude between 175 and 205 nm are calculated on the basis of experimental spectroscopic parameters for the Schumann-Runge rotational lines, adding the contribution of underlying continua which become important near both ends of the band system. Nicolet and Peetermans (1980) emphasize the difficulties of obtaining accurate experimental values of the oscillator strength and of the predissociation linewidth. Figure 5 illustrates the linewidth measured since the work of Ackerman and Biaumé (1970), including the new values recently obtained by Gies et al. (1981). Theoretical values deduced from models are not reported in this figure. The large differences between the available data should play an important role on the photo-




Figure 5 Experimental values of the linewidth ( $\text{cm}^{-1}$ ) in the Schumann-Runge bands of molecular oxygen.

dissociation rates of several trace species. Indeed, if smaller linewidths are used, the absorption by molecular oxygen is more concentrated in the line centers. Consequently the solar radiation in the wavelength interval of 175-200 nm penetrates more deeply in the stratosphere and enhances the photodissociation rates at lower altitudes.

Recent studies on the photodissociation of the molecular oxygen in the Schumann-Runge bands (Kockarts, 1976; Blake, 1979; Nicolet and Peetermans, 1980; Frederick and Hudson, 1979) are based on detailed computation of the rotational structure of the band system. They require lengthy computation and, in order to reduce the computer time and cost, these authors have also developed parameterization methods which can be applied either to calculate the photodissociation rate of molecular oxygen itself or the penetration of the solar radiation in the wavelength range of 175-200 nm. These approximations are only valid when the absorption cross section of the trace species can be averaged over each given wavelength intervals (1 nm,  $500 \text{ cm}^{-1}$ , intervals between successive band heads). On the contrary, when constituents such as nitric oxide are characterized by rotational structures in their cross sections, special treatments are required (Nicolet, 1979; Allen and Frederick, 1982).

Kockarts (1971) has introduced reduction factors defined by the expressions :

$$R_b(M) = \frac{1}{n} \sum_{i=1}^{n_b} \exp(-\tau_{ib}) \quad (3)$$

and

$$R_b(O_2) = \frac{1}{n} \sum_{i=1}^{n_b} \epsilon_b \sigma_{ib} \exp(-\tau_{ib}) \quad (4)$$

where  $b$  denotes each band in the Schumann-Runge band system  
 $\sigma_{ib}$  the  $O_2$  cross section calculated every  $0.5 \text{ cm}^{-1}$   
 $\epsilon_b$  the predissociation probability corresponding to band  $b$   
 $\tau_{ib}$  the optical depth every  $0.5 \text{ cm}^{-1}$   
 $n_b$  the number of intervals in band  $b$ .

The reduction factors are calculated using the molecular parameters taken from Ackerman et al. (1970). Approximations of these reduction factors taking into account the atmospheric temperature profile have been given by Kockarts (1976) :

$$\ln R_b(x) = -a \exp \left[ \sum_{i=1}^6 c_i (x-x_0)^i \right] \quad (5)$$

where  $x$  is the natural logarithm of the total content of molecular oxygen in  $\text{cm}^{-2}$ . The same expressions can also be applied for intervals of  $500 \text{ cm}^{-1}$  and  $1 \text{ nm}$ . The coefficients  $x_0$ ,  $a$  and  $c_1$  through  $c_6$  are given in table 2, 3, 5, 6, 8 and 9 of Kockarts (1976) for each type of wavelength intervals. The calculations are only valid for altitudes above 30 km. The errors, using these approximations, are generally less than 5 percent for most of the intervals. The same set of coefficients can be used for different solar zenith angles and the choice of the wavelength interval is guided by the aeronomic problem to be studied.

In his model, Blake (1979) has adjusted the linewidth values, used as a parameter, in order to improve the agreement between the calculated and the measured cross section value. The best fit is obtained with a linewidth in good agreement with those observed by Lewis et al. (1978) and the cross section model is in good agreement with the measurements of Ackerman et al. (1970) except at a few wavelengths. Blake (1979) proposed an expression giving the mean transmission  $T$  in the middle atmosphere for two wavelength intervals, namely 1.0 and 0.2 nm :

$$T = \exp - \left[ \frac{X}{(1+\beta x N(O_2))^{1/2}} + Y \right] N(O_2) \quad (6)$$

where  $N$  designates the oxygen column density.  $X$ ,  $Y$  and  $\beta$  are coefficients given by Blake (1979) in his tables 3 and 4.

The effective local cross section  $\sigma_e$  is obtained with a polynomial function :

$$\ln(\sigma_e) = \exp \left[ \sum_{n=0}^7 a_n X^n \right] \quad (7)$$

where

$$X = \{ \log [N(O_2)] - 20 \} / 3.5 \quad (8)$$

The  $a_n$  coefficients are given by Blake (1979) in his tables 5 and 6. The standard deviation over all points is 6 percent for the 1.0 nm model and 7 percent for the 0.2 nm model.

Nicolet and Peetermans (1980) have calculated an effective cross section following the expression :

$$\sigma_e(O_2) = \sigma_m \left[ \frac{\sigma_M}{\sigma_m} \right]^{1/(1+e^p)} \quad (9)$$

where  $\sigma_M$  and  $\sigma_m$  are the maximum and the minimum photodissociation cross section defined by :

$$\overline{\sigma}_M(T) = \frac{1}{n} \sum_{i=1}^n \sigma_i(T) \quad (10)$$

when  $N(O_2) \rightarrow 0$ , and by :

$$\sigma_m(T) = [\sigma_i(T)]_m \quad (11)$$

which corresponds to the smallest values of  $\sigma$  in the adopted range when  $N(O_2) \rightarrow \infty$ . In these expressions,  $\sigma_i(T)$  is the absorption cross section corresponding to a narrow interval labelled  $i$  and  $n$  is the number of intervals. Values of  $\sigma_M$  and  $\sigma_m$  are given for three temperatures, 190, 230 and 270 K in table 7 in Nicolet and Peetermans (1980). Parameter  $p$  is a function given by the polynomial expression :

$$p = \sum_{n=0}^n p_i [\ln N(O_2)]^i \quad (12)$$

Values using only two terms ( $n = 1$ ) in expression (12) are given by Nicolet and Peetermans (1980) in their table 8 for the aforementioned temperatures. The transmission related to the mean optical depth  $\tau(O_2)$  is defined by the expression :

$$\exp [-\tau(O_2)] = \exp (-e^d) \quad (13)$$

where  $d$  is given by a second polynomial function :

$$d = \sum_{i=0} d_i [\ln N(O_2)]^i \quad (14)$$

for which coefficient values using only two terms ( $n = 1$ ) are given in table 9 of Nicolet and Peetermans (1980).

The temperature effect cannot be neglected when accurate values are required. Moreover precise calculations should take into account the specific effect of the solar zenith angle since, due to the temperature dependence of the cross section values, optical paths with the same  $O_2$  column content but different zenith angles can lead to significantly different solar attenuation and dissociation rates. When the calculation with a 10 percent precision is sufficient, the expressions with two terms can be used with an average temperature of 230 K and for a maximum total content of molecular oxygen less than  $5 \times 10^{23}$  molecules/cm<sup>2</sup>, corresponding to stratospheric levels above 30 km.

Allen and Frederick (1982) have also recently published a simple polynomial representation of the effective photodissociation cross section, taking into account the effect of the altitude  $z$  and of the zenith angle  $\chi$ :

$$\log \sigma_i^e (z, \chi = 0^\circ) = \sum_j a_{ij} X_i(z)^{j-1} \quad (15)$$



where  $X_i(z)$  is either the temperature  $T(z)$  or the logarithm of the pressure  $\log P(z)$ . The coefficients  $a_{ij}$  are presented by Allen and Frederick (1982) in table 2, where indice  $i$  refers to the  $500 \text{ cm}^{-1}$  wavelength intervals in the spectral region of the Schumann-Runge bands. The error averaged over all altitudes is 9 percent with a maximum value of 22 percent at 50 km. Allen and Frederick (1982) stated that such an error is within the uncertainties of the detailed calculations made by Frederick and Hudson (1979), namely  $\pm 40$  percent.

The effective cross section variation with the zenith angle  $\chi$  is given by :

$$\sigma_i^e(z, \chi) / \sigma_i^e(z, 0) = (\sec \chi)^{-C_i(z)} \quad (16)$$

where  $C_i(z)$  is given by :

$$\log C_i(z) = \sum_j b_{ij} [\log N(z)]^{j-1} \quad (17)$$

where  $N(z)$  is the zenith oxygen column density above the altitude  $z$ . Coefficients  $b_{ij}$  are presented by Allen and Frederick (1982) in table 3.

### 3.4 The Herzberg continuum

The penetration of the solar radiation in the wavelength range of 200-240 nm is driven by the absorption of the molecular oxygen in the Herzberg continuum but also by the absorption of ozone which becomes predominant beyond 220 nm. In addition, extinction by Rayleigh scattering contributes for at least 10 percent of the total attenuation. It contributes also to the minimum values in the Schumann-Runge band system as the underlying Herzberg continuum. The absorption cross sections of molecular oxygen are very weak beyond 200 nm (lower than  $10^{-23} \text{ cm}^2$ ) and accurate values are difficult to obtain. Laboratory measurements require long path lengths or high pressure in the ab-

sorption cell. In this latter case, the absorption is contaminated by the presence of  $O_4$  which is formed by association of two oxygen molecules. Several determinations of the absorption cross section of molecular oxygen in the Herzberg continuum have been performed in the last 20 years. The values measured by Hasson and Nicholls (1971) are somewhat higher than the data obtained by Ditchburn and Young (1962) on which the tables published by Ackerman (1971) are based. The absorption spectrum reported by Shardanand and Prasad Rao (1977) is characterized by the smallest cross section at wavelengths shorter than 230 nm, i.e. in the region which contributes the most to the photodissociation frequency in the middle and lower stratosphere. Other data such as the values by Ogawa (1971) or the tabulation by Hudson and Reed (1979) or by Nicolet (1978) are intermediate.

Recently, Frederick and Mentall (1982) as well as Herman and Mentall (1982a) have derived the molecular oxygen absorption cross section spectrum which is required to explain the observed attenuation of the solar irradiance between two different heights located in the range of 30 to 40 km. The observations have been performed by means of the same balloon borne double monochromator.

Frederick and Mentall (1982), have deduced from their observations on September 15, 1980, that, in the 200-210 nm range, the Hasson and Nicholls (1971) cross sections should be multiplied by factors in the range of 0.52 to 0.68 and the Shardanand and Prasad Rao (1977) values by factors in the range of 0.85 to 0.93. It should be noted however that, when applied to the two sets of laboratory data, these factors do not lead to the same value of the corrected cross sections. In principle, they are calculated assuming that the cross section of ozone is perfectly known and that the ozone column between the two levels of observation is measured without error.

Herman and Mentall (1982a) have derived with the same instrument on April 15, 1980 absorption cross sections which are about 30 percent

smaller than the laboratory results of Sharda and Prasad Rao (1977) from 200 to 210 nm and about 50 percent smaller than those of Hasson and Nicholls (1971). These values deduced from in situ measurements are thus somewhat smaller than those suggested by Frederick and Mentall (1982).

Figure 6 shows that the dispersion in the available data is quite important and leads to significant uncertainties in the photodissociation rate of several trace species. One dimensional models have been used very recently by Froidevaux and Yung (1982) and by Brasseur et al. (1983) in order to estimate the importance of the stratospheric composition changes due to absorption cross section values 30 to 50 percent lower than those currently used in photodissociation calculation.

#### 4. THE ABSORPTION CROSS SECTION OF OZONE

The ozone absorption cross section has been measured in the ultraviolet up to 350 nm and in the visible range. Only one measurement with an estimated uncertainty of  $\pm 10$  percent is available below 200 nm (Tanaka et al., 1953). Numerical values have been deduced from this work for modelling purposes (Ackerman, 1971).

Beyond 200 nm, the Hartley bands peak around 250 nm with a value of  $1.146 \times 10^{-17} \text{ cm}^2$  for the absorption cross section (Hearn, 1961) which decreases below to  $10^{-22} \text{ cm}^2$  through to the Huggins bands beyond 300 nm. The values measured by Inn and Tanaka (1953) are widely used in the photodissociation calculation from 200 to 300 nm. Beyond this wavelength, the temperature dependence of the absorption cross section has to be taken into account. The work of Vigroux (1953) has long been the only available reference.

Since the review published by Ackerman (1971), new measurements are now in progress at the National Bureau of Standards (USA) and at




Figure 6 Absorption cross section of molecular oxygen between 200 and 250 nm.

the University of Reims (France). Preliminary results for the wavelength range of 245 to 345 nm at 295 K and 243 K have been reported by Bass and Paur (1981). These measurements are relative to the peak absorption value measured by Hearn (1961) at 254 nm. Absolute measurements performed by Daumont et al. (1983) at 293 and 223 K between 300 and 350 nm are systematically 12 percent lower than those reported by Bass and Paur (1981). The uncertainty in the new values of the cross section is estimated to be 2 percent or less. The latter measurements resolve with a better precision the features of the absorption spectrum (the instrumental width is about 0.02 nm). Consequently, these data with resolved structure, especially beyond 300 nm, can be used to analyze the solar backscattered radiation measured by satellites (Mc Peters and Bass, 1982).

The absorption of solar radiation by ozone takes place in the mesosphere and in the stratosphere for wavelengths greater than 180 nm. At higher altitude, the ozone density is too small to play any role in the penetration of the ultraviolet solar radiation. The photodissociation of ozone leading to the formation of the excited state of atomic oxygen has been investigated for a long time, particularly by Nicolet (1939).

## 5. SCATTERING

When penetrating in the atmosphere, the solar radiation is not only absorbed by optically active constituents but also scattered by the gas molecules and the aerosols. The irradiance  $q$  at a given altitude  $z$ , wavelength  $\lambda$  and solar zenith angle  $\chi$  is given by :

$$q(\lambda, z, \chi) = q_{\infty} e^{-(\tau_a + \tau_s)} + \int_{\Omega} I(\lambda, z, \Omega) d\Omega \quad (18)$$

where  $I(\lambda, z, \Omega)$  is the intensity of the light which is scattered in the solid angle  $\Omega$ .

The optical depth corresponding to the absorption ( $\tau_a$ ) can be expressed as :

$$\tau_a(\lambda, z, \chi) = \sec \chi [\sigma_a(O_2; \lambda) \int_z^\infty n(O_2) dz' + \sigma_a(O_3; \lambda) \int_z^\infty n(O_3) dz'] \quad (19)$$

since the main absorbing gases are molecular oxygen and ozone. The attenuation due to scattering processes appears through a second contribution of the optical depth ( $\tau_s$ ) which can be written as :

$$\tau_s(\lambda, z, \chi) = \sec \chi [\sigma_R \int_z^\infty n_R dz' + \sigma_M \int_z^\infty n_M dz'] \quad (20)$$

where  $\sigma$  and  $n$  are respectively the scattering cross section and the concentration of the Rayleigh (R) and Mie (M) scatters.

The first effect is governed by air molecules while the second contribution is due to aerosol particles. As indicated by Meier et al. (1982), test calculations have shown that, for standard atmospheric load, Mie scattering has a small effect on the value of the solar irradiance. The cross section related to Rayleigh scattering is given by (Fröhlich and Shaw, 1980) :

$$\sigma_R = \frac{32 \pi (\mu - 1)}{3 n_R^2 \lambda^4} f(\delta) \quad (21)$$

where  $\mu$  is the refractive index of the air which depends on the pressure, the temperature, the relative humidity and varies with wavelength. The function :

$$f(\delta) = (6 + 3 \delta)/(6 - 7 \delta) \quad (22)$$

where  $\delta$  is the depolarization factor, accounts for the anisotropy of the molecules. For a standard mixture of air it has a value close to 1.016 (Fröhlich and Shaw, 1980). When calculating explicitly the Rayleigh scattering cross section can be expressed as :

$$\sigma_R = \frac{3.93 \times 10^{-16}}{[\lambda(\text{nm})]^4 + x} \quad (23)$$

where  $x$  is given by (Fröhlich and Shaw, 1980) :

$$x = 74 \lambda + 5 \times 10^{-5} / \lambda - 0.084 \quad (24)$$

$\lambda$  being expressed in nm.

The effect of scattering on the ultraviolet and visible radiation field in the troposphere and stratosphere has been recently analyzed by Meier et al. (1982) and Nicolet et al. (1982). Their results show that multiple scattering and ground albedo can lead to large amplifications of the solar irradiance in the photochemically active spectral region. These authors have calculated the corresponding enhancement factors as a function of height, solar zenith angle and wavelength. Numerical tables of these factors are given in convenient format for computational use. From their solar ultraviolet observations performed at the altitude of 40 km, Herman and Mentall (1982b) have determined the relative importance of scattering versus absorption. Their analysis shows that the scattered flux in the upper stratosphere is of the order of 1 percent of the direct flux between 220 and 300 nm and becomes larger than 10 percent beyond 310 nm.

## 5. CONCLUSIONS

Significant improvements are required in solar irradiance measurements. Accuracy of the order of 5 percent is needed for the calculation of photodissociation frequencies. Such an objective is achievable by appropriate calibration and measurement strategies. In particular, the magnitude of the solar variability related to the rotation period of the Sun and the 11-year cycle should be further investigated.

New in situ measurements of the atmospheric attenuation of the solar irradiance have questioned the accepted values of the absorption cross section of molecular oxygen in the Herzberg continuum. The low absorption recently observed in this wavelength range needs to be confirmed.

The penetration of the solar radiation in the Schumann-Runge bands is strongly dependent on spectrometric parameters which are not sufficiently well known for an accurate determination of the photodissociation frequencies in this part of the spectrum.

The optical depth around 210 nm is also driven by the ozone absorption. New values of the corresponding cross section in the vicinity of this wavelength are required in order to confirm the presently available data. Determination of the solar penetration should also consider the effect of multiple scattering which plays an important role above 300 nm and cannot be neglected in the calculation of the attenuation around 200 nm.



## REFERENCES

- ACKERMAN, M. (1971). Ultraviolet solar radiation related to mesospheric processes, in Mesospheric Models and Related Experiments (Ed. G. Fiocco), p. 149, Reidel, Dordrecht, Holland.
- ACKERMAN, M. and BIAUME, F. (1970). Structure of the Schumann-Runge bands from the (0-0) to the (13-0) band. J. Molec. Spectrosc. 35, 73.
- ACKERMAN, M., BIAUME, F. and KOCKARTS, G. (1970). Absorption cross sections of the Schumann-Runge bands of molecular oxygen. Planet. Space Sci. 18, 1639.
- ALLEN, M. and FREDERICK, J.E. (1982). Effective photodissociation cross sections for molecular oxygen and nitric oxide in the Schumann-Runge bands. J. Atmos. Sci., 39, 2066.
- ARVESEN, J.C., GRIFFIN, R.N., Jr. and PEARSON, B.D., Jr. (1969). Determination of extraterrestrial solar spectral irradiance from a research aircraft. Appl. Opt. 8, 2215.
- BASS, A.M. and PAUR, R.J. (1981). UV absorption cross-sections for ozone : the temperature dependence. J. Photochem. 17, 141.
- BLAKE, A.J. (1979). An Atmospheric absorption model for the Schumann-Runge bands of oxygen. J. Geophys. Res., 84, 3272.
- BLAKE, A.J., McCOY, D.G., GIES, H.P.F. and GIBSON, S.T. (1980). Photoabsorption in molecular oxygen. Interim Sci. Rep. Department of Physics, Univ. of Adelaide (Australia).
- BOSSY, L. (1983) same issue.
- BRASSEUR, G. and SIMON, P.C. (1981). Stratospheric chemical and thermal response to long-term variability in solar UV irradiance, J. Geophys. Res. 86, 7343.
- BRASSEUR, G., DE RUDDER, A. and SIMON, P.C. (1983). Implication for stratospheric composition of a reduced absorption cross section in the Herzberg continuum of molecular oxygen, Geophys. Res. Lett., 10, 20.
- BROADFOOT, A.L. (1972). The solar spectrum 2100-3200 Å, Astrophys. J. 173, 681.

- BRUECKNER, G.E., BARTOE, J.D.F., KJELDSETH MOE, O. and VAN HOOSIER, M.E. (1976). Absolute solar ultraviolet intensities and their variation with solar activity. Part I : the wavelength region 1750-2100 Å. *Astrophys. J.* 209, 935.
- CARVER, J.H., GIES, H.P., HOBBS, T.I., LEWIS, B.R. and McCOY, D.G. (1977). Temperature dependence of the molecular oxygen photoabsorption cross section near the H Lyman- $\alpha$  line, *J. Geophys. Res.* 82, 1955.
- COOK, J., BRUECKNER, G. and VAN HOOSIER, M. (1980). Variability of the solar flux in the far ultraviolet 1175-2100 Å. *J. Geophys. Res.* 85, 2257.
- DAUMONT, D., MALICET, J. and BRION, J. (1983). Private communication.
- DITCHBURN, R.W. and YOUNG, P.A. (1962). The absorption of molecular oxygen between 1850 and 2500 Å, *J. Atm. Terr. Phys.*, 24, 127.
- DOSE, V., SCHMOCKER, U. and SELE, G. (1975). Photoabsorption coefficient in molecular oxygen in the vicinity of the hydrogen Lyman  $\alpha$  line. *Z. Phys. A.* 274, 1.
- FANG, T.M., WOFYSY, S.C. and DALGARNO, A. (1974). Opacity distribution functions and absorption in the Schumann-Runge bands of molecular oxygen. *Planet. Space Sci.* 22, 413.
- FREDERICK, J.E. and HUDSON, R.D. (1979). Predissociation linewidths and oscillator strengths for the (2-0) to (13-0) Schumann-Runge bands of O<sub>2</sub>. *J. Molec. Spectrosc.* 74, 247.
- FREDERICK, J.E. and MENTALL, J.E. (1982). Solar irradiance in the stratosphere : implication for the Herzberg continuum absorption of O<sub>2</sub>. *Geophys. Res. Lett.* 9, 461.
- FREDERICK, J.E., HUDSON, R.D. and MENTALL, J.E. (1981). Stratospheric observations of the attenuated solar irradiance in the Schumann-Runge band absorption region of molecular oxygen. *J. Geophys. Res.* 86, 9885.

- FRIEDMAN, J., LICHTMAN, S.W. and BYRAN, E.T. (1951). Photon counter measurements of solar X-rays and extreme U.V. light. *Phys. Rev.* 83, 1025.
- FROHLICH, C. and SHAW, G.E. (1980). New determination of Rayleigh scattering in the terrestrial atmosphere. *Appl. Opt.* 19, 1773.
- FROIDEVAUX, L. and YUNG, Y.L. (1982). Radiation and Chemistry in the stratosphere : sensitivity to O<sub>2</sub> absorption cross sections in the Herzberg continuum, *Geophys. Res. Lett.* 9, 854.
- GIES, H.P.F., GIBSON, S.T., McCOY, D.G. and BLAKE, A.J. and LEWIS, B.R. (1981). Experimentally determined oscillator strengths and linewidths for the Schumann-Runge band system of molecular oxygen-III. The (7-0) to (19-0) bands, *J. Quant. Spectrosc. Radiat. Transfer.* 26, 469.
- GOLDSTEIN, R. and MASTRUP, F.N. (1966). Absorption coefficients of the O<sub>2</sub> Schumann-Runge continuum from 1270 Å to 1745 Å using a new continuum source. *J. Opt. Soc. Amer.* 56, 765.
- HALL, J.E. (1972). Atmospheric pressure, density and scale height calculated from H Lyman- $\alpha$  absorption allowing for the variation in cross-section with wavelength, *J. Atm. Terr. Phys.* 34, 1337.
- HASSON, V. and NICHOLLS, R.W. (1971). Absolute spectral absorption measurements on molecular oxygen from 2460-1920 Å : II continuum measurements 2430-1920 Å. *J. Phys. B., Atm. Molec. Phys.* 4, 1789.
- HEARN, A.G. (1961). The absorption of ozone in the ultraviolet and visible regions of the spectrum. *Proc. Phys. Soc. (London)*. 78, 932.
- HEATH, D.F. (1980). A review of observational evidence for short and long term ultraviolet flux variability of the sun, in *Proceedings of International Conference on Sun and Climate, Centre National d'Etudes Spatiales*, p. 163, France.
- HEROUX, L. and SWIRBALUS, R.A. (1976). Full-disk solar fluxes between 1230 and 1940 Å. *J. Geophys. Res.* 81, 436.

- HERMAN, J.R. and MENTALL, J.E. (1982a). O<sub>2</sub> absorption cross sections (187-225 nm) from stratospheric solar flux measurements. *J. Geophys. Res.*, 87, 8967.
- HERMAN, J.R. and MENTALL, J.E. (1982b). The direct and scattered solar flux within the stratosphere. *J. Geophys. Res.* 87, 1319.
- HINTEREGGER, H.E. (1981). Representations of solar EUV fluxes for aeronomical applications, in *The Mesosphere and Thermosphere* (Eds. G. Schmidtke and K.S.W. Champion), p. 39, Pergamon Press, Oxford, U.K.
- HUDSON, R.D. (1971). Critical review of ultraviolet photoabsorption cross sections for molecules of astrophysical and aeronomic interest. *Rev. Geophys. and Space Phys.* 9, 305.
- HUDSON, R.D. and MAHLE, S.H. (1972). Photodissociation rates of molecular oxygen in the mesosphere and lower thermosphere. *J. Geophys. Res.* 77, 2902.
- HUDSON, R.P. and REED, E.I. (1979). The stratosphere : present and future, NASA reference publication 1049, 432 pp.
- HUDSON, R.D., CARTER, V.L. and STEIN, J.A. (1966). An investigation of the effect of temperature on the Schumann-Runge absorption continuum of oxygen, 1580-1950 Å. *J. Geophys. Res.* 71, 2295.
- HUFFMAN, R.E., TANAKA, Y. and LARRABEE, J.C. (1964). Nitrogen and oxygen absorption cross-sections in the vacuum ultraviolet. *Disc. Faraday Soc.* 37, 159.
- INN, E.C.Y. and TANAKA, Y. (1953). Absorption coefficient of ozone in the ultraviolet and visible regions. *J. Opt. Soc. Amer.* 43, 870.
- KOCKARTS, G. (1971). Penetration of solar radiation in the Schumann-Runge bands of molecular oxygen, in *Mesospheric Models and Related Experiments* (Ed. G. Fiocco), p. 160 Reidel, Dordrecht, Holland.
- KOCKARTS, G. (1976). Absorption and Photodissociation in the Schumann-Runge bands of molecular oxygen in the terrestrial atmosphere. *Planet. Space Sci.* 24, 589.

- LABS, D. and NECKEL, H. (1970). Transformation of the absolute solar radiation data into the "International Practical Temperature Scale of 1968". *Solar Phys.* 15, 79.
- LEAN, J.L. and BLAKE, A.J. (1981). The effect of temperature on thermospheric molecular oxygen absorption in the Schumann-Runge continuum. *J. Geophys. Res.* 86, 211.
- LEAN, J.L., WHITE, O.R., LIVINGSTON, W.C., HEATH, D.F., DONNELLY, R.F. and SKUMANICH, A. (1982). A three-component model of the variability of the solar ultraviolet flux : 145-200 nm., *J. Geophys. Res.* 87, 10, 307.
- LEWIS, B.R., CARVER, J.H., HOBBS, T.I., McCOY, D.G. and GIES, H.P.F. (1978). Experimentally determined oscillator strengths and linewidths for the Schumann-Runge band system of molecular oxygen -I. The (6-0) to (14-0) bands. *J. Quant. Spectrosc. Rad. Trans.* 20, 191.
- LEWIS, B.R., CARVER, J.H., HOBBS, T.I., Mc COY, D.G. and GIES, H.P.F. (1979). Experimentally determined oscillator strengths and linewidths for the Schumann-Runge band system of molecular oxygen - II. The (2-0) to (5-0) bands. *J. Quant. Spectrosc. Rad. Trans.* 21, 213.
- Mc PETERS, R.D. and BASS, A.M. (1982). Anomalous atmospheric spectral features between 300 and 310 nm interpreted in light of new ozone absorption coefficient measurements. *Geophys. Res. Lett.* 9, 227.
- MEIER, R.R., ANDERSON, D.E., Jr., and NICOLET, M. (1982). Radiation field in the troposphere and stratosphere from 240-1000 nm -I. General analysis. *Planet. Space Sci.* 30, 923.
- METZGER, P.H. and COOK, G.R. (1964). A reinvestigation of the absorption cross sections of molecular oxygen in the 1050-1800 Å region. *J. Quant. Spectrosc. Rad. Trans.* 4, 107.
- MOUNT, G.H. and ROTTMAN, G.J. (1981). The solar spectral irradiance 1200-3184 Å near solar maximum : July 15, 1980. *J. Geophys. Res.* 86, 9193.

- MOUNT, G.H. and ROTTMAN, G.J. (1983). The solar absolute spectral irradiance 1150-3173 Å : 17 May 1982. J. Geophys. Res. in press.
- MOUNT, G.H., ROTTMAN, G.J. and THIMOTHY, J.G. (1980). The solar spectral irradiance 1200-2550 Å at solar maximum. J. Geophys. Res. 85, 4271.
- NECKEL, H. and LABS, D. (1981). Improved data of solar spectral irradiance from 0.33 to 1.25 μ. Solar Phys. 74, 231.
- NICOLET, M. (1939). Le problème aéronomique dans l'atmosphère supérieure, Mém. Inst. r. météorol. Belg. 11, 1.
- NICOLET, M. (1945). Contribution to the study of ionospheric structure. Mém. Inst. r. météorol. Belg. 19, 1.
- NICOLET, M. (1978). Etude des réactions chimiques de l'ozone dans la stratosphère, Institut royal météorologique de Belgique, Bruxelles, Belgique, 536 pp.
- NICOLET, M. (1979). Photodissociation of nitric oxide in the mesosphere and stratosphere : simplified numerical relations for atmospheric model calculations. Geophys. Res. Lett. 6, 866.
- NICOLET, M. and MANGE, P. (1954). The dissociation of oxygen in the high altitude. J. Geophys. Res. 59, 15.
- NICOLET, M. and PEETERMANS, W. (1980). Atmospheric absorption in the O<sub>2</sub> Schumann-Runge band spectral range and photodissociation rates in the stratosphere and mesosphere. Planet. Space Sci. 28, 85.
- NICOLET, M., MEIER, R.R. and ANDERSON, D.E., Jr. (1982). Radiation field in the troposphere and stratosphere - II. Numerical analysis. Planet. Space Sci. 30, 935.
- OGAWA, M. (1968). Absorption coefficients of O<sub>2</sub> at the Lyman-alpha line and its vicinity. J. Geophys. Res. 73, 6759.
- OGAWA, M. (1971). Absorption cross-sections of O<sub>2</sub> and CO<sub>2</sub> continua in the Schumann and far U.V. regions. J. Chem. Phys. 54, 2550.
- OGAWA, S. and OGAWA, M. (1975). Absorption cross section of O<sub>2</sub> (a<sup>1</sup>Δg) and O<sub>2</sub> (X<sup>3</sup>Σg) in the region from 1087 to 1700 Å. Canad. J. Phys. 53, 1845.

- PARK, J.H. (1974). The equivalent mean absorption cross-sections for O<sub>2</sub> Schumann-Runge bands : application to the H<sub>2</sub>O and NO photo-dissociation rates. J. Atmos. Sci. 31, 1893.
- PURCELL, J.D. and TOUSEY, R. (1960). The profile of solar hydrogen Lyman  $\alpha$ . J. Geophys. Res. 65, 370.
- ROTTMAN, G.J. (1981). Rocket measurements of the solar spectral irradiance during solar minimum, 1972-1977. J. Geophys. Res. 86, 6697.
- ROTTMAN, G.J., BARTH, C.A., THOMAS, R.J., MOUNT, G.H., LAWRENCE, G.M., RUSCH, D.W., SANDERS, R.W., THOMAS, G.E. and LONDON, J. (1982). Solar spectral irradiance, 120 to 190 nm, October 13, 1981 -January 3, 1982. Geophys. Res. Lett. 9, 587.
- ROTTMAN, G.J. (1983), 27-day variations observed in solar ultraviolet (120-300 nm) irradiance. Planet. Space Sci. (same issue).
- SAMAIN, D. and SIMON, P.C. (1976). Solar flux determination in the spectral range 150-210 nm. Solar Phys. 49, 33.
- SHARDANAND and PRASAD RAO, A.D. (1977). Collision-induced absorption of O<sub>2</sub> in the Herzberg continuum. J. Quant. Spectrosc. Radiat. Transfer 17, 433.
- SIMON, P.C. (1975). Nouvelles mesures de l'ultraviolet solaire dans la stratosphere. Bull. Acad. r. Belg. Cl. Sci. 61, 399.
- SIMON, P.C. (1978). Irradiation solar flux measurements between 120 and 400 nm. Current position and future needs. Planet. Space Sci. 26, 355.
- SIMON, P.C. (1981). Solar irradiance between 120 and 400 nm and its variations. Solar Phys. 74, 273.
- SIMON, P.C., PASTIELS, R. and NEVEJANS, D. (1982a). Balloon observations of solar ultraviolet irradiance at solar minimum. Planet. Space Sci. 30, 67.

- SIMON, P.C., PASTIELS, R., NEVEJANS, D. and GILLOTAY, D. (1982b). Balloon observations of solar ultraviolet irradiance during solar cycle 21 in Proceedings of the Symposium on the Solar Constant and the Spectral Distribution of Solar Irradiance (Eds. J. London and C. Fröhlich), IAMAP Third Scientific Assembly, p. 95, Boulder, USA.
- SMITH, L.G. and MILLER, K.L. (1974). The measurement of O<sub>2</sub> number density by absorption of Lyman  $\alpha$ . J. Geophys. Res. 79, 1965.
- TANAKA, Y., INN, E.C.Y. and WATANABE, K. (1953). Absorption coefficients of gases in the vacuum ultraviolet, Part IV. Ozone J. Chem. Phys. 21, 1651.
- VIDAL-MADJAR, A. (1977). The solar spectrum at Lyman  $\alpha$  in The Solar output and its variation (Ed. O. White), p. 213, Colorado Associated University Press, Boulder, USA.
- VIGROUX, E. (1953), Contribution à l'étude expérimentale de l'absorption de l'ozone. Ann. Phys. 2, 709.
- WATANABE, K. (1958). Ultraviolet absorption processes in the upper atmosphere in Advances in Geophysics (Eds. H.E. Landsberg and J. Van Mieghem) 5, 153.
- WATANABE, K. and MARNO, F.F. (1956). Photoionization and total absorption cross section of gases - II O<sub>2</sub> and N<sub>2</sub> in the region 850-1500 Å. J. Chem. Phys. 25, 965.
- WATANABE, K., INN, E.C.Y. and ZELIKOFF, M. (1953). Absorption coefficients of oxygen in the vacuum ultraviolet. J. Chem. Phys. 21, 1026.
- WEEKS, L.H. (1975). Determination of O<sub>2</sub> density from Lyman  $\alpha$  ion chambers. J. Geophys. Res. 80, 3655.