

issn 0065-3713

INSTITUT D'AERONOMIE SPATIALE DE BELGIQUE

3 - Avenue Circulaire

B - 1180 BRUXELLES

AERONOMICA ACTA

A - N° 314 - 1986

Remote sensing ozone measurements from stratospheric
balloon during map/globus campaign 1983

by

P.C. SIMON, W. PEETERMANS, E. PLATEAU,

P. RIGAUD, J.-P. NAUDET, D. HUGUENIN,

D. OFFERMAN and H. RIPPEL

BELGISCH INSTITUUT VOOR RUIMTE-AERONOMIE

3 - Ringlaan

B - 1180 BRUSSEL

**REMOTE SENSING OZONE MEASUREMENTS FROM STRATOSPHERIC BALLOON
DURING MAP/GLOBUS CAMPAIGN 1983**

P.C. SIMON and W. PEETERMANS

Institut d'Aéronomie Spatiale de Belgique, Bruxelles, Belgium

E. PLATEAU

Faculté des Sciences, Université de l'Etat à Mons, Belgium.

P. RIGAUD and J.-P. NAUDET

Laboratoire de Physique et Chimie de l'Environnement

Orléans, France

D. HUGUENIN

Observatoire de Genève, Switzerland

D. OFFERMANN and H. RIPPEL.*

Physics Department, University of Wuppertal, FRG

Abstract

Remote sensing measurements of ozone in ultraviolet, visible and infrared domains performed in September 1983 by means of stratospheric balloons during the MAP/Globus campaign in France are presented and discussed. Ozone concentrations deduced from planet and solar occultation measurements in the Huggins and Chappuis absorption bands are given with an uncertainty of ± 10 percent. The results obtained from absorption measurements in the Chappuis bands give systematically higher concentration values than those deduced from in situ techniques. Discrepancies of the order of 20 percent are found out between 24 and 34 km of altitude. They are larger below the ozone maximum and above 35 km. The analysis of the data shows an aerosol contribution in the measured optical depth in the Chappuis bands, for altitude below 28 km giving overestimated inferred ozone concentrations. Discrepancies at higher altitudes with in situ soundings are difficult to explain.

* Now at Kayser-Threde GmbH, Munich, FRG.

Résumé

Des observations de l'ozone dans l'ultraviolet, le visible et l'infrarouge ont été réalisées en septembre 1983 à l'aide de ballons stratosphériques au cours de la campagne MAP/Globus. Les concentrations d'ozone déduites des mesures d'occultation dans les bandes de Huggins et de Chappuis, en utilisant le soleil ou la planète Venus comme source, sont présentées avec une incertitude de $\pm 10\%$. Les résultats obtenus par absorption dans les bandes de Chappuis sont systématiquement plus élevés que ceux déduits des méthodes d'observation in situ. Un désaccord de l'ordre de 20% est mis en évidence entre 24 et 34 km, pour augmenter en-dessous du maximum de l'ozone et au-dessus de 35 km d'altitude. Ces écarts importants peuvent être expliqués par une extinction supplémentaire due aux aérosols en-dessous de 28 km, mais sont difficiles à comprendre aux altitudes supérieures.

Samenvatting

Ozonmetingen in ultraviolette, zichtbare en infrarode spectrale gebieden, uitgevoerd met behulp van stratosferische ballons in september 1983 tijdens de MAP/Globus campagne in Frankrijk, worden voorgesteld en besproken. Ozonconcentraties, afgeleid uit occultatiemetingen in de absorptiebanden van Huggins en Chappuis met de zon of de planeet Venus als bron, worden met een onzekerheid van $\pm 10\%$ gegeven. De resultaten die bekomen werden uit absorptiemetingen in de Chappuisbanden geven systematisch hogere concentratiewaarden dan die afgeleid uit in situ technieken. Afwijkingen van 20% worden gevonden tussen 24 en 34 km en deze worden nog groter beneden het ozonmaximum en boven de 35 km. Deze belangrijke verschillen kunnen uitgelegd worden door een aanvullende 'uitdoving' te wijten aan de aërosols beneden de 28 km, maar zijn moeilijk te verklaren op grotere hoogten.

Zusammenfassung

Ozonmessungen im Ultraviolett, im Infrarot und in sichtbaren Gebieten, getan mit der Hilfe von stratosphärischen Ballons im September 1983 während der MAP/Globus Kampagne in Frankreich, werden präsentiert und besprochen. Ozonkonzentrationen, abgeleitet aus Bedeckungsmessungen in den Absorptionsbanden von Huggins und Chappuis mit der Sonne oder dem Planet Venus als Quelle, werden mit einer Ungewissheit von $\pm 10\%$ gegeben. Die Resultaten bekommen aus Absorptionsmessungen in den Chappuisbanden geben systematisch höhere Konzentrationswerte als die abgeleitet aus in situ Techniken. Abweichungen von 20% werden gefunden zwischen 24 und 34 Km und diese steigen noch unter dem Ozonmaximum und über 35 Km. Diese wichtige Unterschiede können erklärt werden durch eine ergänzende Extinktion, den Aërosols zuzuschreiben, unter 28 Km, aber diese sind schwer erklärlich auf grösseren Höhen.

INTRODUCTION

Quantitative measurement of ozone concentration in the stratosphere is of fundamental importance in order to understand the aeronomic processes controlling the ozone budget and to distinguish between possible changes due to natural variations (e.g. solar variability response, Brasseur and Simon, 1981; Garcia et al. 1984) and those due to anthropogenic influences.

Several methods are presently used to derive ozone vertical profile from balloon observations. Among those, remote sensing methods based on solar occultation in the ultraviolet and in the visible ranges similar to satellite observations, on infrared emission and on planet occultation in the visible range have been used during the MAP/Globus campaign in 1983 in order to compare results obtained from different techniques. Previous campaigns (Aimedieu et al., 1983; Hilsenrath et al., 1985) have already pointed out systematic differences among various methods of observations.

The purpose of this paper is to present the intercomparison of remote sensing measurements of ozone performed in September 1983 by means of balloon observations in the frame of the MAP/Globus project (Offermann, 1983). Ground-based remote measurements made during the same epoch are reported by de La Noë et al. (1987).

PRINCIPLE OF MEASUREMENTS AND INSTRUMENTATION

The data reported and analyzed in this work were obtained from three different instrumentations separately integrated in three gondolas carried by stratospheric balloons. Table 1 summarized the information concerning these three flights.

The first flight carried a stabilized platform of the Geneva Observatory (Huguenin and Magnan, 1978) in which a double grating

TABLE 1.

Flight nr	Gondola	Date	Time UT	Observation Method	Institution
29	G5	13-14 Sept.	night	Planet occultation (Chappuis bands)	LPCE, Orléans
34	G1	23 Sept.	4h30	IR emission (9.6 μ m)	Univ. of Wuppertal
37	G2b	28 Sept.	Sunset	Solar occultation (Huggins and Chappuis bands)	IASB, Bruxelles

spectrometer JY DH 10 was integrated. This instrumentation has been extensively described by Rigaud et al (1983). Ozone is measured by its absorption in the Chappuis bands, from 647 to 672 nm, during the night, by observing Venus occultation by the Earth.

The second flight measured the thermal infrared emission of atmospheric trace gases by means of an Ebert-Fastie grating spectrometer working in the wavelength interval 5-19 μm . Ozone emission at 9.6 μm were measured by vertical limb scanning performed by a mirror, covering $\pm 2.4^\circ$ with respect to the balloon horizon. The whole instrument is cooled by liquid helium. An extensive description of instrumentation and calibration methods is given by Rippel et al. (1986).

The third flight measured the ozone content by absorption in the ultraviolet and visible domains, using the Sun as a source. The instrument is a double grating spectrometer with holographic gratings of 10 cm focal length made by Jobin-Yvon. A silicon photodiode EG & G type UV444-B is used as detector. The wavelength range extends from 250 to 700 nm and the band-pass of the spectrometer (FWHM) is 1 nm, with a triangular shape. A high-pass filter with a cutt-off at 300 nm has been set in front of the detector. Wavelength scans below this limit provide accurate measurement of the instrument "optical zero" including the electronic and telemetry offset and possible residual stray-light. Two wavelength intervals, namely 320-340 nm and 540-660 nm, corresponding to the Huggins and Chappuis absorption bands of ozone are used to measure the solar absorption during sunset. In addition, differential absorption method as developed for nitric dioxide measurements between 430 and 450 nm can be applied in the Huggins bands for comparison purposes with Bouguer's plot. A more detailed description of the instrumentation is given elsewhere (Simon et al., 1987).

OBSERVATIONS AND DATA ANALYSIS

The balloons were successfully launched by the "Centre National d'Etudes Spatiales" (CNES) from the french balloon launching facility situated at Aire sur l'Adour (France, 43°42' N, 0°15' W). Table 1 gives the dates, time of observation, type of measurements and wavelength ranges used for ozone profile determinations.

a) In the case of planet occultations, the reference spectrum (called extra-atmospheric spectrum of Venus) is recorded at zenith angles smaller than 80°. For the float altitude of this flight, namely 39 km, the optical depth due to absorption of atmospheric constituents is negligible for such zenith angles in the wavelength range of the observations (647-672nm), and cannot be detected.

The wavelength scale of each spectrum is adjusted to the H α line at 656.3 nm. A synthetic attenuated spectrum is fitted to each measured transmission during occultation, by using the least square method. The ozone absorption cross-section are those compiled by Ackerman (1971). Attenuation due to Rayleigh scattering is calculated using the extinction cross section formula published by Nicolet (1984) and a density profile taken from the U.S. Standard Atmosphere, 1976. For the lower altitudes of the line of sight (tangent height below 20 km), the residual optical depth is attributed to stratospheric aerosol assuming no wavelength dependence in the observed wavelength range (647-672 nm). A more complete description of the experimental procedure and data analysis has been published by Rigaud et al. (1983).

b) For solar absorption measurements in the Huggins and the Chappuis bands of ozone, the Bouguer law was applied to determine the total content of ozone along the line of sight. The so-called reference spectrum has been recorded for a solar zenith angle of 82° at about 29 km of altitude. Consequently, this measured spectrum has been corrected for residual ozone absorption in the wavelength ranges of interest and also

for Rayleigh scattering extinction according to the formula published by Nicolet (1984). The representative ozone vertical profile defined for the MAP/Globus campaign (de La Noë et al., 1987) given in Table 2 and model published by Krueger and Minzner (1976) have been taken for altitudes above the balloon. The absorption cross-sections measured by Bass (1984) have been used for absorption correction computation up to 340 nm and those published by Griggs (1968) for the Chappuis bands. Ratio between corrected reference spectrum and spectra recorded for zenith angles larger than 90° have been calculated. Their natural logarithms have been plotted versus absorption cross-section of ozone. Intensity values have been corrected for Rayleigh scattering extinction along the optical path corresponding to each recorded spectra. The integrated content of ozone is given by the slope of the linear regression in equation :

$$\ln \frac{I_o(\lambda)}{I(\lambda)} - \tau_{RS}(\lambda) = N_{O_3} \sigma_{O_3}(\lambda) \quad (1)$$

where

λ is the wavelength

$I_o(\lambda)$ is the solar irradiance above the atmosphere

$I(\lambda)$ is the solar irradiance after its absorption in the atmosphere

τ_{RS} is the optical depth due to Rayleigh scattering extinction

N_{O_3} is the total content of ozone along the line of sight (cm^{-2})

σ_{O_3} is the absorption cross section of ozone.

On the other hand, the three wavelength differential absorption method has been used for analysing the same data in the Huggins bands, according to publications of Brewer et al. (1973) and Gillis et al. (1982). Table 3 defines the maxima and minima in the absorption cross-sections selected between 320 and 340 nm.

c) The infrared emission measurements at the limb were performed only at

TABLE 2. MAP/GLOBUS representative atmosphere
Average 9-28 September 1983.

Altitude km	Pressure mb	Temperature K	Ozone Conc. (cm ⁻³)
14	150	208	1.11 x 10 ¹²
16	107	211	1.72
18	77	215	2.70
20	56	218	3.86
22	41.5	220	4.61
24	30.2	222	4.67
26	22.5	224	4.43
28	16.3	228	3.50
30	12.0	233	2.65
32	8.7	238	2.03 ⁺
34	6.5	243 [*]	1.58 ⁺

* Extrapolated values

+ Krueger and Minzner (1976)

36.5 km of altitude because of the failure of the spectrometer cryogenic system. Consequently, the observing time was reduced to 10 minutes. A more extensive description of the experimental procedure and data evaluation is published by Rippel et al. (1986).

DISCUSSION OF RESULTS

Figure 1 shows the integrated concentration along the line of sight for flight 29 and 37 performed respectively on 14 September and 28 September 1983. The comparison between these two observations is not straightforward because of, first, the differences in balloon altitudes during the occultation measurements and, second, observations were made eastward by pointing the planet Venus and, obviously, westward for sunset. The lowest tangent altitude is respectively situated at about 500 km away from the balloon, north of the "Observatoire de Haute Provence" (OHP), and at about 200 km, above Bordeaux. Differences between the two curves on figure 1 are explained by longer optical depth for flight 29 for which the balloon altitude during occultation is 38 km instead of 29 km for flight 37. When optical depths become comparable, the two curves are merging.

It should be mentioned that the Bouguer law as described in the previous section is not verified over the wavelength interval 540-660 nm used for solar occultation measurements on 28 September 1983. Deviation from this law increases for lower tangent altitudes of the line of sight and cannot be explained either by errors on ozone cross-sections or Rayleigh scattering corrections. According to the 2 years of observations made after the eruption of El Chichon in April 1982, by means of an airborne lidar at various latitudes and reported by McCormick and Swisler (1986), the aerosol loading in the stratosphere is still very important in September 1983, at latitudes corresponding to the balloon flights (44°N). This fact is also supported by the Solar Mesosphere Explorer (SME) satellite observations showing an important increase of limb radiances at various wavelengths after April 1982, between 20 and 38




Figure 1. Comparison between integrated concentration of ozone along the line of sight deduced from absorption measurements in the Chappuis bands performed on September 1983 (Venus occultation at 38 km of altitude) and on 28 September 1983 (solar occultation at 29 km of altitude).

km of altitude (Thomas et al., 1986). The residual optical thicknesses have been evaluated for both balloon observations in the Chappuis bands. The data inversion yields to a peak value of $8 \times 10^{-3} \text{ km}^{-1}$ at 16 km of altitude. The vertical profiles are systematically higher than those deduced from previous balloon flights performed in 1980 and 1981 (Naudet, 1983). In addition, for solar occultation measurements, the residual optical thickness strongly increases for wavelength lower than 620 nm (Simon et al., 1986). This fact could explain the higher values for the ozone integrated concentrations deduced from the data around 620 nm and below. A more detailed analysis is presented in Simon et al. (1986).

Figure 2 gives detailed results deduced from the data measured during flight 37 on 28th September 1983, by using the Bouguer's plots in the Huggins and Chappuis bands and also the differential absorption method applied in the Huggins bands.

Differential absorption method gives systematically higher values with an average divergence of 20 percent. This difference is difficult to explain except that when using the Bouguer technique, the integrated ozone concentration is deduced by a linear regression through about 80 data points scattered in the Huggins bands instead of five triplets, namely only 11 different wavelengths, for the differential absorption (see table 3). This could introduce additional but not systematic error.

Vertical profiles of ozone were derived by using a non-linear iterative inversion method (Naudet, 1983) for flight 29 and a layer-by-layer linear inversion method for flight 37. Results are summarized on figure 3 with the data obtained at 36.5 km from infrared limb emission measurement performed on 23 September 1983. Corresponding numerical values are given in table 4, 5 and 6. The uncertainties on inferred ozone concentrations from planet and solar occultation measurements are ± 10 percent.




Figure 2. Comparison between integrated concentration of ozone along the line of sight deduced from absorption measurement in the Huggins and Chappuis bands performed on 28 September 1983 (solar occultation at 29 km of altitude).




Figure 3. Comparison between ozone profiles deduced from remote sensing measurements performed during the MAP/Globus campaign in September 1983. In situ average profile deduced from measurements performed on 24 September 1983 is also shown for discussion purposes.

The flight dates refer to the following institutions :

14 Sept. 83 LPCE, Orléans (P. Rigaud)

23 Sept. 83 U. of Wuppertal (D. Offermann)

24 Sept. 83 in situ average profile (Airmedieu *et al.*, 1987)

28 Sept. 83 IASB, Bruxelles (P.C. Simon).

TABLE 3. Wavelength triplets

Triplet	Wavelengths (nm)		
	Min	Max	Min
1	321.250	322.250	323.850
2	323.850	324.950	326.700
3	326.700	327.900	329.475
4	329.475	330.925	332.575
5	332.575	333.750	335.600

TABLE 4. Gondola G5 14 September 1983 night

Pressure (mb)	Altitude (km)	O ₃ Conc. (cm ⁻³ x10 ⁻¹²)	Δ O ₃ [*] x 10 ⁻¹²	O ₃ mixing ratio ppmV
75.7	18	4.94	+ 0.50	1.95
55.3	20	6.33	+ 0.42	3.43
40.5	22	5.97	+ 0.33	4.45
29.7	24	5.34	+ 0.27	5.47
21.9	26	4.92	+ 0.22	6.91
16.2	28	4.28	+ 0.14	8.21
12.0	30	3.25	+ 0.11	8.50
8.9	32	2.31	+ 0.08	8.20
6.6	34	1.75	+ 0.07	8.50
5.0	36	1.37	+ 0.06	9.10
3.8	38	0.95	+ 0.09	8.56

* Errors related to altitude uncertainties not included

TABLE 5. Gondola G1 23 September 1983 4h 30 U.T.

Pressure (mb)	Altitude (km)	O ₃ Conc. (cm ³ x 10 ⁻¹²)	Accuracy	O ₃ mixing ratio ppmV
4.85	36.5	1.25	± 15%	8.50

TABLE 6. Gondola G2b 28 September 1983 sunset

Pressure (mb)	Altitude (km)		O ₃ Conc. (cm ⁻³ × 10 ⁻¹²)		Mixing ratio ppmV	
	Huggins	Chappuis	Huggins	Chappuis	Huggins	Chappuis
13.8	29.1		2.73 ₋ 10%		6.31	
14.2		28.9		3.17-3.87		7.10-8.67
14.7	28.7		3.13 ₋ 10%		6.76	
16.1		28.1		3.00-3.96		5.88-7.76
17.2	27.7		4.28 ₋ 10%		7.81	
20.7		26.5		4.42-5.40		6.63-8.10
22.2	26.1		4.27 ₋ 10%		5.96	
29.5		24.2		4.82-6.50		5.01-6.76

The error sources for these observations come from signal-to-noise ratio, absorption cross section values, ozone concentration above the balloon, air masses along the line of sight, altitude determination and resolution and refraction for altitudes below 20 km. According to Rigaud et al. (1985), uncertainties due to the signal-to-noise ratio is less than ± 2 percent. The absorption cross section values are generally accurate within ± 5 percent. Possible errors due to their temperature dependence could be excluded (Humphrey and Badger, 1947). The balloon altitude and the geometry of the line of sight are based on pressure measurements performed by sensor integrated in the payload. An altitude accuracy of the order of 500 m can only be estimated and affects the vertical profiles by approximatively the same amount. The altitude resolution is very good in the case of planet occultations but decreases from 230 m to 1.4 km during solar occultation for flight 37. The ozone concentration profile above the balloon altitude is based on the aforementioned models, introducing an error of about ± 6 percent mainly for the first ozone concentration value nearest to the balloon altitude. Calculated air mass values for atmospheric layers below the balloon altitude could introduced errors of ± 3 percent. The errors bars on figure 3 illustrates the total uncertainty of ± 10 percent except for the solar absorption data in the Chappuis bands which also include the uncertainties coming from the Bouguer law departures.

At 36.5 km of altitude, the only ozone concentration value deduced from infrared remote sensing is in agreement with the ozone profile obtained by planet occultation measurement. This latter method gives systematically higher values than the representative model of ozone based on in situ soundings made from OHP and Biscarosse (de La Noë et al., 1987). Differences are between 10 and 20 percent, except below 24 km of altitude where day-to-day variations due dynamical changes in the lower stratosphere should be taken into account. In addition, aerosol absorption becomes more significantly important, leading also to larger discrepancies. Solar occultation measurements in the Huggins bands give ozone concentrations in agreement over 3 km in altitude with in situ

soundings except at one altitude, namely 27.7 km. The same observations in the Chappuis bands at 25.5 and 24.2 km of altitude yield to values very close to those obtained from planet occultation. This fact supports the assumption of an important contribution of aerosol in September 1983, even around 660 nm, on the optical depth measured below 28 km of altitude.

Systematic discrepancies with in situ measurements at higher altitudes reported by Aumedieu et al. (1987) are difficult to explain in terms of aerosol loading of the stratosphere. On the other hand, direct influence of transport is weak in such region (above 30 km and at 44° N) where ozone budget is chemically controlled according to the 2-D model of Garcia and Solomon (1985). However, the comparison of other trace species observed during the same campaign shows similar discrepancies. These have been tentatively interpreted in terms of wavelike dynamical disturbances (Offermann et al., 1987).

In conclusion, ozone measurements by occultation techniques are very precise but the final accuracy depends upon a good knowledge of all atmospheric absorptions along the line of sight. Such a knowledge can be reached by extending the wavelength range of observation out of the absorption bands of ozone in order to accurately determine optical depths of the stratosphere at various wavelengths and to reveal atmospheric extinction by other species like aerosol. A good altitude resolution is needed in order to resolve possible layer structure in aerosol profiles.

ACKNOWLEDGMENTS

This work was partly supported by "Bundesministerium für Forschung und Technologie", (FRG), by "Centre National de la Recherche Scientifique" and "Centre National d'Etudes Spatiales" (France). The support given by the CNES balloon center at Aire sur l'Adour and the MAP/GLOBUS 1983 coordinator Prof. D. OFFERMANN are gratefully acknowledged.

REFERENCES

- Ackerman, M. (1971). Ultraviolet solar radiation related to mesospheric processes in Mesospheric Models and Related Experiments (ed. G. Fiocco), D. Reidel, p. 149.
- Aimedieu, P., Krueger, A.J., Robbins, D.E. and Simon, P.C. (1983). Ozone profile intercomparison based on simultaneous observations between 20 and 40 km. Planet. Space Sci., 31, 801.
- Aimedieu, P., Matthews, W.A., Attmannspacher, W., Cisneros, J., Hartmannsgruber, R., Komhyr, W. and Robbins, D.E. (1987). Comparison of in situ stratospheric ozone measurements obtained during the MAP/Globus 83 campaign. Planet. Space Sci., same issue.
- Bass, A. (1984). Private communication.
- Brasseur, G. and Simon, P.C. (1981). Stratospheric chemical and thermal response to long-term variability in solar UV. J. Geophys. Res., 86, 7343.
- Brewer, A.W., McElroy, C.T. and Kerr, J.B. (1973). Nitrogen dioxide concentrations in the atmosphere. Nature, 246, 129.
- de La Noë, J., Brillet, J., Turaci, C., Mégie, G., Pelon, J., Marché, P., Barbe, A., Gibbins, C.J. and Dawkins, A.W.J. (1987). Remote ground based measurements during the MAP/GLOBUS 1983 campaign. Planet. Space Sci., same issue.
- Garcia, R.R. and Solomon, S. (1985). The effect of breaking gravity waves on the dynamics and chemical composition of the mesosphere and lower thermosphere, J. Geophys. Res., 90, 3850.
- Garcia, R.R., Solomon, S., Roble, R.C. and Rusch, D.W. (1984). A numerical response of the middle atmosphere to the 11-year solar cycle. Planet. Space Sci., 32, 411.
- Gillis, J.R., Goldman, A., Williams, W.I. and Murcray, D.G. (1981). Atmospheric ozone profiles from high resolution UV spectra obtained with a balloon-borne spectrometer. Appl. Optics, 21, 413.

- Griggs, M.J. (1968). Absorption coefficients of ozone in the ultraviolet and visible regions. *J. Chem. Phys.*, 49, 857.
- Hilsenrath, E., Ainsworth, J., Holland, A., Mentall, J., Torres, A., Attmanspacher, W., Bass, A., Evans, W., Komhyr, W., Mauersberger, K., Miller, A.J., Profitt, M., Robbins, D., Taylor, S. and Weinstock, E. (1985). Results from the balloon ozone intercomparison campaign (BOIC). in Atmospheric Ozone (Eds. C.S. Zerefos and A. Ghazi), D. Reidel, p.454.
- Huguenin, D. and Magnan, A. (1978). Nacelle astronomique stabilisée avec détecteur U.V à galette de microcanaux. Proceedings of Esrange Symposium. Rep. ESA SP-135.
- Humphrey, G. and Badger, R. (1947). The absorption spectrum of ozone in the visible. *J. Chem. Phys.* 15, 794.
- McCormick, M.P. and Swissler, T.J. (1986). Temporal variation and decay of the El Chichon stratospheric aerosol. *J. Geophys. Res.*, to be published.
- Krueger, A.J. and Minzner, R.A. (1976). A mid-latitude ozone model for the 1976 U.S. standard atmosphere. *J. Geophys. Res.*, 81, 1477.
- Naudet, J.-P. (1983). Mesures de constituants minoritaires de la stratosphère, Thèse de doctorat.
- Nicolet, M. (1984). On the molecular scattering in the terrestrial atmosphere : an empirical formula for its calculation in the homosphere. *Planet. Space Sci.*, 32, 1467.
- Offermann, D. (1983). MAP/GLOBUS Campaign 1983. Handbook for MAP (Ed. C.F. Sechrist Jr.) 8, 17.
- Offermann, D., Rippel, H., Aïmediu, P., Matthews, W.A., Mégie, G., Arijs, E., Ingels, J., Nevejans, D., Arnold, F., Schlager, H., Attmanspacher, W., Cisneros, J.M., Dawkins, A.W., De Muer, D., Fabian, P., Karcher, F., Froment, G., Langematz, U., Reiter, R., Rothe, K.W., Schmidt, U. and Thomas, R.J. (1987). Disturbance of stratospheric trace gas mixing ratios during the MAP/Globus 1983 campaign. *Planet. Space Sci.*, same issue.

- Rigaud, P., Naudet, J.P. and Huguenin, D. (1983). Simultaneous measurements of vertical distribution of stratospheric NO_3 and O_3 at different periods of the night. *J. Geophys. Res.* 88, 1463.
- Rigaud, P., Naudet, J.P. and Huguenin, D. (1985). Mesure de la répartition verticale de l'ozone atmosphérique par spectrophotométrie d'absorption dans le visible. in Atmospheric Ozone (Eds. C.S. Zerefos and A. Ghazi), D. Reidel, p. 509.
- Rippel, H., Hilbert, L., Jarisch, M., Kampf, D., Offermann, D. and Schneider, R. (1986). A helium cooled balloon borne infrared experiment for measurement of stratospheric trace gas. Submitted to *Appl. Optics*.
- Simon, P.C., Peetermans, W. and Plateau, E. (1986). Stratospheric ozone and aerosol measurements by solar occultation with a balloon-borne spectrometer. *EOS, Transactions, American Geophysical Union*, 67, 876, 1986.
- Simon, P.C., Peetermans, W., Vandeneede, P. and Plateau, E. (1987). Stratospheric ozone and nitric dioxide measurements from UV and visible solar absorption observations by balloon. To be published.
- Thomas, G.E., Rusch, D.W., Thomas, R.J. and Todd Clancy, R. (1986). Long-term changes in the stratosphere due to the 1982 eruption of El Chichon. Workshop on ozone variation on climatological time scales : observations and theory, and perturbations by UV solar flux variations and the eruption of El Chichon, Salzburg, Austria, August 19-22, 1985.