

issn 0065-3713

I N S T I T U T D A E R O N O M I E S P A T I A L E D E B E L G I Q U E

3 - Avenue Circulaire

B - 1180 BRUXELLES

AERONOMICA ACTA

A - N° 371 - 1993

**Infrared Cross Sections and Global Warming Potentials of
10 Alternative Hydrohalocarbons**

by

C.CLERBAUX, R.COLIN, P.C.SIMON and C.GRANIER

B E L G I S C H I N S T I T U U T V O O R R U I M T E - A E R O N O M I E

3 - Ringlaan

B - 1180 BRUSSEL

Foreword

This work is an expanded version of a publication accepted in J.Geophys.Res. as a brief report.

Avant-Propos

Ce travail est une version d'une publication acceptée dans J.Geophys.Res., avec un tableau et des figures supplémentaires.

Voorwoord

Dit werk is een uitgebreide versie van een publikatie aanvaard in het tijdschrift J.Geophys.Res. Het bevat een tabel en aanvullende figuren.

Vorwort

Diese Arbeit ist eine Version, mit einer zusätzlichen Tafel und zusätzlichen Figuren versehen, von einer Veröffentlichung in J.Geophys.Res.

Infrared Cross Sections and Global Warming Potentials of 10 Alternative Hydrohalocarbons

C. CLERBAUX, R. COLIN

*Laboratoire de Chimie Physique Moléculaire, CPI 160/09,
Université Libre de Bruxelles, 50 av. Roosevelt, 1050 Brussels, Belgium*

P.C. SIMON

*Institut d'Aéronomie Spatiale de Belgique,
3 av. Circulaire, 1180 Brussels, Belgium*

C. GRANIER

*National Center for Atmospheric Research, P.O. Box 3000,
Boulder, CO 80307, USA*

Abstract.

Absorption cross sections have been obtained in the infrared atmospheric window, between 600 and 1500 cm^{-1} , for 10 alternative hydrohalocarbons : HCFC-22, HCFC-123, HCFC-124, HCFC-141b, HCFC-142b, HCFC-225ca, HCFC-225cb, HFC-125, HFC-134a and HFC-152a. The measurements were made at three temperatures (287K, 270K and 253K) with a Fourier transform spectrometer operating at 0.03 cm^{-1} apodized resolution. Integrated cross sections have been introduced into a two dimensional radiative-chemical-dynamical model in order to calculate the Global Warming Potential of each gas.

Résumé

Les sections efficaces d'absorption de 10 hydrohalocarbures alternatifs (HCFC-22, HCFC-123, HCFC-124, HCFC-141b, HCFC-142b, HCFC-225ca, HCFC-225cb, HFC-125, HFC-134a and HFC-152a) ont été obtenues dans la fenêtre atmosphérique IR, entre 600 et 1500 cm^{-1} . Les spectres ont été enregistrés à 3 températures (287K, 270K et 253K) à l'aide d'un spectromètre à transformée de Fourier opérant à la résolution de 0.03 cm^{-1} . L'introduction des sections efficaces intégrées dans un modèle 2D radiatif-chimique-dynamique a conduit à la détermination des 'Global Warming Potentials' des différents gaz.

Samenvatting

De werkzame absorptiedoorsneden van 10 alternatieve zachte chloorfluorkoolwaterstoffen (HCFC-22, HCFC-123, HCFC-124, HCFC-141b, HCFC-142b, HCFC-225ca, HCFC-225cb, HFC-125, HFC-134a and HFC-152a) werden bekomen in het atmosferisch venster tussen 600 en 1500 cm^{-1} . De spectra werden by 3 temperaturen (287K, 270K et 253K) gemeten door een spectrometer met Fourier transformatie en werkend met een resolutie van 0.03 cm^{-1} . Door de invoering van de geïntegreerde werkzame absorptiedoorsneden in een radiatief, chemisch, dynamisch 2D model heeft men de 'Global Warming Potentials' van verschillende gassen kunnen bepalen.

Zusammenfassung

Die effektiven Absorptionsquerschnitten von 10 alternativen Hydrohalokarbidien (HCFC-22, HCFC-123, HCFC-124, HCFC-141b, HCFC-142b, HCFC-225ca, HCFC-225cb, HFC-125, HFC-134a and HFC-152a) wurden gemessen in dem atmosphärischen IR Fenster zwischen 600 und 1500 cm^{-1} . Die Spektren wurden bei 3 Temperaturen (287K, 270K et 253K) mit der Hilfe eines Fourier-Spectrometers mit 0.03 cm^{-1} Auflösung registriert. Die Einführung von integrierten effektiven Querschnitten in einem 2D radiativen-chemischen-dynamischen Modell hat die Bestimmung von den 'Global Warming Potentials' der verschiedenen Gasen ermöglicht.

1. INTRODUCTION

Substitutes for the currently used chlorofluorocarbons (CFCs) have been introduced worldwide since the latter were recognized to be responsible for the depletion of stratospheric ozone (Molina and Rowland, 1974; Stolarski and Cicerone, 1974). The international agreements (Montreal Protocol, 1987; London Amendment, 1990) have established the phaseout of CFCs by the year 2000 for industrialized nations and 10 years later for developing countries. Alternative chemicals and technologies are being developed by industries to meet the future needs in refrigeration, foam-blowing, insulation, electronic cleaning and drying processes (Fisher *et al.*, 1991). The partially halogenated hydrochlorofluorocarbons (HCFCs) and hydrofluorocarbons (HFCs) were found to have the same physical properties as CFCs and to be environmentally more acceptable compounds. Unlike the long-lived CFCs, which are essentially removed through UV photolysis processes in the stratosphere, hydrohalocarbons contain at least one hydrogen atom and hence react with hydroxyl radicals in the troposphere. According to their reactivity with OH, their lifetimes range from 1.5 to 35 years, resulting in a lower Ozone Depletion Potential (Wuebbles, 1983; Fisher *et al.*, 1990a; WMO, 1992).

The potential contribution of HCFCs and HFCs as greenhouse gases has to be considered owing to their strong absorption bands located in the IR atmospheric window. The ability of one gas to participate to the radiative forcing of the atmosphere is characterizable by a parameter, the Global Warming Potential, which is calculated from the infrared absorption spectra and the amount of gas present in the atmosphere over a specified period of time (Fisher *et al.*, 1990b; WMO, 1992).

Although hydrohalocarbons are present in very small amounts in the present-day atmosphere, some have reached the minimum concentration levels of a few ppbv required to be measured in the ambient air (Pollock *et al.*, 1992). As these concentrations are going to increase in the forthcoming years, spectroscopic infrared data are necessary in order to determine atmospheric abundances of these trace gases using infrared spectroscopy.

This paper reports the accurate measurements of infrared absorption cross sections, as a function of the temperature, for HCFC-22 (CHClF_2), HCFC-123 (CHCl_2CF_3), HCFC-124 ($\text{CHClF}_2\text{CF}_3$), HCFC-141b ($\text{CH}_3\text{CCl}_2\text{F}$), HCFC-142b (CH_3CClF_2), HCFC-225ca ($\text{CHCl}_2\text{CF}_2\text{CF}_3$), HCFC-225cb ($\text{CClF}_2\text{CF}_2\text{CHClF}$), HFC-125 (CHF_2CF_3), HFC-134a (CFH_2CF_3) and HFC-152a (CH_3CHF_2). This data set has been introduced into the NCAR coupled radiative-chemical model in order to obtain Global Warming Potentials for the substitutes.

2. EXPERIMENTAL

The spectra have been recorded in the infrared atmospheric window, between 600 and 1500 cm^{-1} , using a Bruker IFS 120HR Fourier Transform Spectrometer. A globar source and a mercury-cadmium-telluride (MCT) detector, maintained at liquid N_2 temperature, were employed. With an apodized resolution of 0.03 cm^{-1} , a hundred scans were necessary to obtain a good signal to noise ratio; the baseline noise does not exceed 0.01 in absorbance units.

The samples were provided by Solvay S.A. (HCFC-22, HCFC-123, HCFC-141b, HCFC-142b, HFC-134a), E.I. du Pont de Nemours (HCFC-124, HFC-125, HFC-152a) and Asahi Glass Company (HCFC-225ca, HCFC-225cb). Each of these compounds was provided with a purity analysis : 8 of them were guaranteed to have a volumic purity higher than 99.98% and 2 were analyzed to be purer than 99.0% (HCFC-124, HFC-125). The gas to be studied was introduced into a laboratory built thermostatic cell (Hurtmans *et al*, 1992). The 5 cm long cell was constructed of anodised aluminium and closed by two pairs of KBr windows. Its exact length was measured by interferometry with an accuracy of 0.2%. The cell was placed inside the sample chamber of the spectrometer which was then evacuated in order to avoid interferences with ambient CO_2 and H_2O . The pressure of the gas to be analyzed was measured using a 100 torr full scale absolute MKS Baratron capacitance gauge (model 390 HA), stabilized at 45°C. According to the gauge manufacturer, the error associated with the measured pressure is lower than 0.15% in the range of pressure used (1-4 torr). The temperature of the cell can be stabilized anywhere between -30° and +60°C by liquid circulating in a double wall within the body of the cell . The temperature stabilization was provided by a cryostat (Neslab ULT80) and methanol was used as a circulating fluid. The temperature of the gas is measured inside the cell by a transducer accurate to $\pm 0.2^\circ\text{C}$. The nominal temperatures (287K, 270K, 253K) were reproduced in each individual measurement to within 0.5°C.

For each species eighteen spectra have been recorded, at six different pressures and three different temperatures. In addition, similar measurements were made for CFC-12, in order to allow a comparison with the numerous data available in the literature for this compound.

3. RESULTS

3.1 Temperature Dependent Cross Sections

According to the Beer-Lambert law, the transmission of radiation through a

homogeneous gas sample is described by the relation

$$I(\nu) = I_o(\nu) \exp(-\sigma(\nu)nl) \quad (1)$$

where I_o and I are the intensities of the incident and transmitted radiation at wavenumber ν , n is the concentration of the gas (molec/cm³) and l is the optical path (cm). σ is the absorption cross section (cm²/molec) and can be written

$$\sigma(\nu) = \frac{1}{nl} \ln \frac{I_o(\nu)}{I} \quad (2)$$

For the relatively heavy molecules studied here, the individual rotational lines are merged into a continuum (Q branches) or grouped in peaks (P and R branches), as can be observed in figure 1. The 0.03 cm⁻¹ resolution is however insufficient to perform a complete rotational analysis of the bands. All the absorption features are convolved with a shape function which includes both the line profile (Doppler, Lorentz or Voigt according to the experimental conditions) and the instrumental lineshape due to the finite resolving power of the instrument. To avoid the problem of the finite spectral resolution, different techniques have been used. Pressure broadening techniques by a foreign nonabsorbing gas are described by Wilson and Wells (1946), Penner and Weber (1951) and Kagann *et al* (1983). The curve-of-growth method, which requires the measurement of an equivalent width, is discussed by Benedict *et al* (1956). Mc Daniel *et al* (1991) have used a method of data reduction which eliminates the need for addition of inert gas into the cell. In our experiment, which uses a Fourier spectrometer, the finite resolution is associated with the limited displacement δ of the scanning mirror. This truncation corresponds formally to a multiplication of the interferogram with a boxcar function. After Fourier transformation the spectrum is thus the convolution of the "true" spectrum with a $\sin x/x$ function ($x=1/\delta$), which is the Fourier transform of a boxcar. By multiplying the interferogram by a well chosen smoothing function, the strong sidelobes in the spectrum are damped. We have thus simply apodized our data with a Norton-Beer strong function (Griffiths and de Haseth, 1986) in order to minimize the instrumental distortion and to preserve the true line surface.

The cross sections were determined using equ.(2) for each spectral data point, 0.0085 cm⁻¹ apart, between 600 and 1500 cm⁻¹. Some care must be taken in obtaining the base e absorbance $\ln I_o/I$: a small baseline shift, due to the very slow heating of the source and of the detector, often occurs. To avoid this, two empty cell background spectra were taken, one before and one after each recorded spectrum, and the mean value of these has been used as I_o . The quality of the baseline was checked each time by observing zero absorbance in areas where no




Figure 1. Temperature dependence of the absorption cross sections, between 964 and 971 cm⁻¹, for HCFC-142b.

absorption occurs. Secondly, all the pressures were chosen to prevent saturation of the spectra : a maximum of 75% absorption was allowed. Neglecting these two precautions would lead to large errors in the determined cross sections.

In order to increase the accuracy, a linear least squares fit was applied for each point using all 6 pressures. For each of the three temperatures the cross sections were obtained by this procedure and zero values have been set in the regions where no absorption occurs and only noise was recorded. It should be noted that no dependence of these cross sections on pressure was found, in contrast to the results reported in Mc Daniel *et al* (1991) for eight CFCs measured at six different temperatures. The uncertainties associated with these cross sections, at 0.03 cm⁻¹ resolution, range from 1-2% for strong absorptions to 3-4% for weak absorptions. They were obtained by adding the uncertainties due to the following sources of error, described in chapter 2: pressure and temperature measurements, purity determination, pathlength and standard deviation in the least squares fit. The cross sections values obtained for each compound at 287K, between 600 and 1500 cm⁻¹, are plotted in figure 2 to 11. The complete set of data is available on PC diskettes upon request.

Atmospheric abundances of CFCs have been determined by IR Fourier transform spectroscopy using laboratory spectroscopic data (Zander *et al*, 1987). As the emissions of hydrohalocarbons are expected to increase in the future, the temperature dependent cross section data obtained in this work can be used as reference spectra for atmospheric detection of these species by satellite, aircraft or balloon-borne infrared spectrometers. As reported in Varanasi (1992), it is important to know the temperature and the pressure dependence of the spectral features in order to derive correct concentrations. As can be seen in figure 1, the intensity of a strong Q branch may increase by as much as 33% with a 35K decrease in temperature. No systematic studies of the pressure-broadening effects have been undertaken.

3.2 Integrated Band Strengths

Integrated band strengths have been calculated using the formula :

$$\sigma_{int} = \int_{\nu_1}^{\nu_2} \sigma(\nu) d\nu = \sum_{\nu_1}^{\nu_2} \sigma(\nu) \Delta\nu \quad (3)$$

Two different sets of data have been calculated. To obtain the results presented in Table 1 and 2, the integration limits ν_1 and ν_2 were chosen in order to cover the main absorption features. Some integration regions may contain several vibrational bands. In Table 1, the integrated cross sections σ_{int} obtained for CFC-12 and

HCFC-22




Figure 2. Absorption cross sections (287K), between 700 and 1500 cm^{-1} , for HCFC-22.

HCFC-123




Figure 3. Absorption cross sections (287K), between 700 and 1500 cm⁻¹, for HCFC-123.

HCFC-124




Figure 4. Absorption cross sections (287K), between 600 and 1500 cm^{-1} , for HCFC-124.




Figure 5. Absorption cross sections (287K), between 700 and 1500 cm^{-1} , for HCFC-141b.

HCFC-142b




Figure 6. Absorption cross sections (287K), between 600 and 1500 cm^{-1} , for HCFC-142b.




Figure 7. Absorption cross sections (287K), between 600 and 1500 cm^{-1} , for HCFC-225ca.




Figure 8. Absorption cross sections (287K), between 600 and 1500 cm^{-1} , for HCFC-225cb.




Figure 9. Absorption cross sections (287K), between 700 and 1500 cm⁻¹, for HFC-125.

HFC-134a




Figure 10. Absorption cross sections (287K), between 700 and 1500 cm⁻¹, for HFC-134a.




Figure 11. Absorption cross sections (287K), between 700 and 1500 cm^{-1} , for HFC-152a.

HCFC-22 are compared with the numerous results published in the literature. The uncertainties on the last given digit are indicated in parentheses. Some results, originally reported in $\text{cm}^{-2}\text{atm}^{-1}$, have been transformed into cm molec^{-1} units. Our results for these two molecules agree well with the results reported by most of these authors. In Table 2, the integrated band strengths determined for all 10 HCFCs and HFCs are given for the three temperatures investigated. A small temperature dependence appears to exist for some of these bands.

A second set of data has been derived (Table 3) in order to be used in atmospheric radiative models (see section 3.3), constant intervals of 100 cm^{-1} have been chosen to divide up the spectrum.

A set of experiments was carried out in order to test the effect of the instrument resolution on the measured values of the integrated cross sections, at a resolution 5 times higher (0.006 cm^{-1}) and 5 times lower (0.15 cm^{-1}) than measured previously. Peak cross sections were strongly affected by resolution changes, but no significant differences were found for the integrated values.

3.3 Global Warming Potentials

In order to compare the effect on the climate caused by the emission of radiatively active gases, a relative index has been introduced : The Global Warming Potential (IPCC, 1990). The GWP of a well-mixed gas is defined as the time integrated commitment to radiative forcing from the instantaneous release of 1kg of a trace gas expressed relative to that from the release of 1kg of a reference gas :

$$GWP = \frac{\int_0^n a_i c_i dt}{\int_0^n a_{ref} c_{ref} dt} \quad (4)$$

where a_i and a_{ref} are the instantaneous radiative forcings due to a mass unit increase in the concentration of the gas, c_i and c_{ref} are the concentrations of the trace gas i and of the reference gas remaining at time t after its release and n the number of years over which the calculation is performed. The gas chosen as reference is generally CO_2 for climate studies (IPCC, 1990) or CFC-11 for halocarbons or ozone related studies (Fisher *et al*, 1990b). The radiative forcing due to a perturbation in the concentration of a greenhouse gas is given by the net radiative flux change (in W/m^2) induced by this perturbation at the tropopause. Due to an incomplete understanding of all the interaction feedback mechanisms on clouds, oceans and polar ice, a characterization in terms of direct heat trapping or radiative forcing is generally preferred to a characterization by a change in the Earth's surface temperature. Because CFCs and alternatives are present in very small quantities and their absorption bands do not overlap with those of

Table 1 : Integrated cross sections ($\text{cm molec}^{-1} \times 10^{17}$) for CFC-12 and HCFC-22

Molecule	Integration limits (cm^{-1}) ^a	Varanasi and Ko (1977) ^b 296K	Kagann <i>et al</i> (1983) ^b 296K	Massie <i>et al</i> (1985) ^b 296K	Varanasi <i>et al</i> (1988) ^b 300K	Fisher <i>et al</i> (1990b) 296K	McDaniel <i>et al</i> (1991) 293K	This work 287K
CFC12	850 - 950	5.322(161)	5.832(186)	6.131(615)	5.809(49)	5.742(?)	5.455(545)	5.880(124)
(CF ₂ Cl ₂)	1070 - 1130	4.644(89)	4.602(157)	{ 7.139(715)	4.615(45)	{ 7.142(?)	4.206(421)	4.497(76)
	1130 - 1190	2.935(56)	3.134(105)		2.931(57)		2.925(292)	3.083(71)
HCFC22	765-855	-	-	-	2.233(8)	2.284(?)	2.132(213)	2.357(90)
(CHClF ₂)	1060-1210	-	-	-	7.179(93)	6.751(?)	6.007(601)	6.825(171)
	1275-1380	-	-	-	-	1.269(?)	0.882(88)	1.081(89)

^a The limits of integration are those used in the present work

^b Reported in Mc Daniel *et al* (1991)

Table 2 : Integrated cross sections ($\text{cm molec}^{-1} \times 10^{17}$) for HCFCs and HFCs.

Molecule	Integration limits (cm^{-1})	Integrated cross sections			Uncertainties
		T = 287K	270K	253K	
HCFC-22 (CHClF_2)	765-855	2.36	2.32	2.25	(.09)
	1060-1210	6.83	6.73	6.64	(.17)
	1275-1380	1.08	1.11	1.09	(.09)
HCFC-123 (CHCl_2CF_3)	740-900	2.39	2.30	2.08	(.14)
	1080-1450	10.49	10.26	10.09	(.37)
HCFC-124 (CHClFCF_3)	675-715	.53	-	-	(.04)
	790-920	1.81	-	-	(.14)
	1035-1430	12.09	-	-	(.52)
HCFC-141b ($\text{CH}_3\text{CCl}_2\text{F}$)	710-790	2.37	2.22	2.16	(.08)
	895-955	1.09	1.02	1.00	(.06)
	990-1210	3.95	3.72	3.33	(.20)
	1325-1470	.34	.33	.34	(.12)
HCFC-142b (CH_3CClF_2)	650-705	.75	.69	.68	(.05)
	875-1030	2.58	2.51	2.40	(.14)
	1075-1265	7.19	7.11	6.96	(.20)
	1360-1475	.61	.64	.78	(.09)
HCFC-225ca ($\text{CHCl}_2\text{CF}_2\text{CF}_3$)	695-865	3.02	3.02	3.00	(.08)
	1010-1420	14.69	14.62	14.49	(.42)
HCFC-225cb ($\text{CClF}_2\text{CF}_2\text{CHClF}$)	715-880	3.19	3.45	3.48	(.16)
	890-990	1.65	1.73	1.68	(.09)
	1025-1375	10.74	11.27	11.36	(.36)
HFC-125 (CHF_2CF_3)	700-745	.53	-	-	(.04)
	840-890	.67	-	-	(.05)
	1060-1465	14.91	-	-	(.51)
HFC-134a (CFH_2CF_3)	815-865	.25	.21	.24	(.04)
	935-1010	.79	.81	.84	(.07)
	1035-1130	1.41	1.43	1.47	(.09)
	1135-1340	9.71	9.76	9.68	(.23)
	1345-1485	.45	.46	.44	(.11)
HFC-152a (CH_3CHF_2)	840-995	1.15	1.17	1.15	(.14)
	1050-1205	4.40	4.44	4.51	(.16)
	1320-1490	1.34	1.40	1.41	(.15)

Table 3 : Integrated cross sections (cm/molec $\times 10^{17}$) over 100 cm^{-1} intervals

Integration limits (cm^{-1})	Integrated cross sections (T=287K)									
	HCFC22	HCFC123	HCFC124	HCFC141b	HCFC142b	HCFC225ca	HCFC225cb	HFC125	HFC134a	HFC152a
600-700	0.0	0.0	0.336	0.0	0.743	0.005	0.0	0.0	0.0	0.0
700-800	0.555	0.265	0.210	2.373	0.012	1.609	1.965	0.530	0.0	0.0
800-900	1.803	2.125	1.746	0.002	0.560	1.402	1.233	0.675	0.247	0.117
900-1000	0.0	0.0	0.050	1.089	1.996	0.0	1.645	0.0	0.795	1.030
1000-1100	0.891	0.024	0.609	1.081	0.199	1.756	1.429	0.088	0.686	0.069
1100-1200	5.933	5.208	4.726	2.860	4.854	2.500	7.450	5.206	5.429	4.329
1200-1300	0.137	3.800	5.594	0.002	2.156	9.251	1.661	7.703	3.561	0.001
1300-1400	0.944	1.409	1.099	0.254	0.398	1.166	0.199	1.840	1.507	0.490
1400-1500	0.0	0.046	0.062	0.085	0.211	0.012	0.0	0.069	0.388	0.850
600-1500	10.263	12.877	14.432	7.746	11.129	17.701	15.582	16.111	12.613	6.886

the main greenhouse gases, the radiative forcing associated with such compounds increases linearly with the amount of absorber. For the more abundant gases where saturation effects occur, the dependence is better approximated by a square-root or a logarithmic function.

The cross section data we have measured for 10 hydrohalocarbons have been introduced in the NCAR radiative-chemical-dynamical 2D interactive model (Brasseur *et al*, 1990) and GWP values have been obtained, relative to CFC-11, for time horizons corresponding to 5, 10, 20, 50, 100, 200 and 500 years.

Radiative forcings of HCFCs and HFCs have been determined with a new version of the radiative code used in the NCAR Community Climate Model version 1 (CCM1) (Kiehl *et al*, 1987) : the new longwave scheme takes into account the thermal radiative effects of H₂O, CO₂, CH₄, O₃, N₂O, CFC-11 and CFC-12 (Briegleb, 1992). In this wideband model, the averaged integrated cross sections over intervals of 100 cm⁻¹ have been introduced for each of the HCFCs and HFCs. The small temperature dependences of these data have been neglected due to the larger uncertainties present in other parameters. The forcings calculated by the model, relative to CFC-11, are reported in Table 4 on a molecule-per-molecule basis and on a kg-per-kg basis. As the location and the intensity of the absorbing bands differ for each species, the relative forcings can be smaller or greater than unity. Our values are generally in agreement (better than 10%) with the radiative forcings reported by IPCC (1990) (from Fisher *et al*, 1990b), except for HCFC-123 and HFC-125 for which our values are higher by 12% and lower by 15% respectively. The results of Fisher *et al* (1990b) are based on unpublished cross section values; it is therefore difficult to comment further on the observed discrepancies. Radiative forcings for HCFC-225ca and HCFC-225cb are reported, to our knowledge, for the first time.

For all the greenhouse gases included in the model, the vertical concentration profiles from 0 to 85 km and the atmospheric lifetimes have been determined with an updated version of the 2D model described by Brasseur *et al* (1990). Current ground level concentrations have been used as a boundary condition for all the major gases (including HCFC-22) and set at 2.5ppt for all the other hydrohalocarbons. Due to the very small quantities of hydrohalocarbons introduced in the model no indirect effects, such as changes in the distribution of stratospheric or tropospheric O₃, were included. Atmospheric lifetimes were obtained using the rate coefficients for reaction with OH from WMO (1992) for tropospheric removal and the photodissociation rates were computed by using the UV absorption cross sections from Gillotay *et al* (1991a, 1991b). These lifetimes are reported in Table 4 and are in general agreement with the other published results (IPCC, 1990 ; WMO, 1992). All the GWPs relative to CFC-11 obtained for the different timescales are summarized in Table 4. The differences between our results and the GWP values

Table 4 : Lifetimes (years), radiative forcings and GWPs relative to CFC-11

Halocarbon	Lifetime	Radiative forcing		Global Warming Potential						
		per molec.	per kg	5	10	20	50	100	200	500 (years)
CFC-11 ^a	57.0	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
HCFC-22	14.3	.85	1.35	1.19	1.06	.86	.56	.41	.35	.34
HCFC-123	1.5	.90	.81	.25	.13	.07	.04	.03	.02	.02
HCFC-124	6.0	.94	.95	.67	.50	.32	.17	.12	.10	.10
HCFC-141b	9.7	.66	.77	.63	.53	.39	.22	.16	.14	.13
HCFC-142b	21.1	.82	1.12	1.04	.97	.86	.64	.50	.43	.41
HCFC-225ca	2.4	1.08	.73	.32	.19	.10	.05	.04	.03	.03
HCFC-225cb	6.8	1.31	.88	.65	.50	.34	.18	.13	.11	.11
HFC-125	33.9	.91	1.04	1.01	.98	.93	.82	.71	.64	.62
HFC-134a	13.1	.78	1.06	.92	.81	.64	.41	.29	.25	.24
HFC-152a	1.5	.50	1.03	.31	.17	.09	.05	.03	.03	.03

^aused as reference gas.

reported, relative to CO₂, in WMO (1992) are lower than 10% and are mostly due to the different lifetimes used, except for HFC-125. For this substitute, GWP values lower by 25% were found and are mainly due to the discrepancy in the radiative forcings.

CO₂ is often preferred as a reference gas to compare the impact on global warming of all the radiatively active gas (CH₄, N₂O...). In this work, model calculations have also been performed relative to CO₂ but were found to be less representative and are not presented here. The uncertainties associated with the lifetime of CO₂ and the strong non linear dependence of the radiative forcing on the concentration changes make CFC-11 a better choice.

4. DISCUSSION AND CONCLUSION

In this paper, experimental infrared cross section data at relatively high resolution and at three atmospheric temperatures have been obtained for hydrohalocarbons. Global Warming Potentials have been determined for different time horizons, using integrated cross sections into a 2D radiative-chemical model.

Fisher *et al* (1990b) were the first to calculate the GWPs of CFCs and substitutes using CFC-11 as a reference gas. They conclude that all substitutes are environmentally preferable to CFC-11, but in their work it was assumed that the integration time was set to infinity. We agree with their general conclusion. The integration over different time horizons obtained in this work shows that, as expected, the warming effect decreases exponentially with a time constant equal to the atmospheric lifetime of the species. But these calculations also show that for three compounds (HCFC-22, HCFC-142b, HFC-125) the GWPs (relative to CFC-11) are higher than unity for a five to ten year period. Moreover, these compounds maintain a relatively strong GWP for a very long time. Therefore, taking into account the fact that for some industrial applications larger quantities of CFC alternatives are sometimes needed to replace the present CFCs in order to obtain an equivalent efficiency, their contribution to the global warming may remain important. We hope that the present results will help in the definition of future regulations.

Acknowledgments. We thank Guy Brasseur, Bruce Briegleb, John Orlando and Steve Massie (NCAR) for helpful discussions and comments in the preparation of this paper. We are grateful to Solvay S.A. (Belgium)(in particular James Franklin), E.I. du Pont de Nemours (USA) and Asahi Glass Company (Japan) for supplying the samples through AFEAS (USA). Didier Lemaître is acknowledged for his contribution in the experimental work .

This work has been supported by the Belgian State- Prime Minister's Service- Science Policy Office (*Global Change* and *Eurotrac/Topas* programs). Claire Granier is supported by the Gas Research Institute (USA) under contract n° 5090-254-1993.

REFERENCES

- Benedict, W.S., R. Herman, G.E. Moore, and S. Silverman, The strengths, widths, and shapes of infrared lines. I. General considerations, *Can.J.Phys.* **34**, 830-849, 1956.
- Brasseur, G., M.H. Hitchman, S. Walters, M. Dymek, E. Falise, and M. Pirre, An interactive chemical dynamical radiative two-dimensional model of the middle atmosphere, *J.Geophys.Res.* **95**, 5639-5655, 1990.
- Briegleb, B.P., Longwave band model for thermal radiation in climate studies, private communication (accepted for publication in *J.Geophys.Res.*).
- Fisher, D.A., C.H. Hales, D.L. Filkin, M.K. Ko, N.D. Sze, P.S. Connell, D.J. Wuebbles, I.S. Isaksen, and F. Stordal, Model calculations of the relative effects of CFCs and their replacements on stratospheric ozone, *Nature* **344**, 508-512, 1990a.
- Fisher, D.A., C.H. Hales, W.C. Wang, M.K. Ko, and N.D. Sze, Model calculations of the relative effects of CFCs and their replacements on global warming, *Nature* **344**, 513-516, 1990b.
- Fisher, S.K., P.J. Hughes, P.D. Fairchild, C.L. Kusik, J.T. Dieckmann, E.M. McMahon, and N. Hobday, Energy and global warming impacts of CFC Alternative Technologies, Executive Summary, DOE/AFEAS, 1991.
- Gillotay D., and P.C. Simon, Temperature dependence of ultraviolet absorption cross sections of alternative chlorofluoroethanes, *J. Atmos.Chem.* **12**, 269-285, 1991a.
- Gillotay D., and P.C. Simon, Temperature dependence of ultraviolet absorption cross sections of alternative chlorofluoroethanes : 2. The 2-chloro 1,1,1,2-tetrafluoroethane - HCFC124, *J. Atmos.Chem.* **13**, 289-299, 1991b.
- Griffiths, P.R. , and J.A. de Haseth, Fourier transform infrared spectroscopy, *Chemical analysis*, **83**, Wiley-interscience, 1986.
- Hurtmans, D., R. Petrisse, D. Lemaître, C. Clerbaux, and J. Vander Auwera, A thermostatic absorption cell for high accuracy measurement of absolute intensities, *EPA Newsletter* **44**, 40-42, 1992.

- IPCC, Climate change - The IPCC Scientific Assessment, Chap.2, Report prepared for the WMO and UNEP Intergovernmental Panel on Climate Change, by working group 1, J.T. Houghton, G.J. Jenkins, and J.J. Ephraums eds., Cambridge Univ. Press, New York, NY, 1990.
- Kagann, R.H., J.W. Elkins, and R.L. Sams, Absolute band strengths of halocarbons F-11 and F-12 in the 8 to 16- μ m region, *J.Geophys.Res.* **88**, 1427-1432, 1983.
- Kiehl J.T., R.J. Wolsky, B.P. Briegleb, and V. Ramanathan, Documentation of radiation and cloud routines in the NCAR Community Climate Model (CCM1), NCAR Tech. Note, NCAR/TN-288+IA, Boulder, Colorado, 1987.
- Massie, S.T., A. Goldman, D.G. Murcray, and J.C. Gille, Approximate absorption cross sections of F12, F11, ClONO₂, N₂O₅, HNO₃, CCl₄, CF₄, F21, F113, F114, and HNO₄, *Appl.Opt.*, **24**, 3426-3427, 1985.
- Mc Daniel, A.H., C.A. Cantrell, J.A. Davidson, R.E. Shetter, and J.G. Calvert, The temperature dependent, infrared absorption cross-sections for chlorofluorocarbons : CFC-11, CFC-12, CFC-13, CFC-14, CFC-22, CFC-113, CFC-114, and CFC-115, *J. Atmos.Chem.* **12**, 211-227, 1991.
- Molina, M.J., and F.S. Rowland, Stratospheric sink for chlorofluoromethanes : chlorine atom-catalysed destruction of ozone, *Nature* **249**, 810-814, 1974.
- Penner, S.S. and D. Weber, Quantitative infrared intensity measurements : I. Carbon monoxide pressurized with infrared-inactive gases, *J.Chem.Phys.*, **19**, 807-816, 1951.
- Pollock, W.H., L.E. Heidt, R.A. Lueb, J.F. Vedder, M.J. Mills, and S.Solomon, On the age of stratospheric air and ozone depletion potentials in polar regions, private communication (accepted for publication in *J.Geophys.Res.*).
- Stolarski, R.S., and R.J. Cicerone, Stratospheric chlorine : A possible sink for ozone, *Can.J.Chem.* **52**, 1610-1615, 1974.
- Varanasi, P., and F. Ko, Intensity measurements in freon bands of atmospheric interest, *J.Quant.Spectrosc.Radiat.Transfer* **17**, 385-388, 1977.
- Varanasi, P., and S. Chudamani, Infrared intensities of some chlorofluorocarbons capable of perturbing the global climate, *J.Geophys.Res.* **93**, 1666-1668, 1988.
- Varanasi, P., Absorption spectra of HCFC-22 around 829 cm⁻¹ at atmospheric conditions, *J.Quant.Spectrosc.Radiat.Transfer* **47**, 251-255, 1992.

- Wilson, E.B., Jr, and A.J. Wells, The experimental determination of the intensities of infra-red absorption bands, I. Theory of the method, *J.Chem.Phys.*, **14**, 578-580, 1946.
- WMO, World Meteorological Organization, Scientific Assessment of Ozone Depletion: 1991, *Global Ozone Research and Monitoring Project*, Report n° 25, Geneva, Switzerland, 1992.
- Wuebbles, D.J., Chlorocarbon emission scenarios : potential impact on stratospheric ozone, *J.Geophys.Res.* **88**, 1433-1443, 1983.
- Zander, R., C.P. Rinsland, C.B. Farmer, and R.H. Norton, Infrared spectroscopic measurements of halogenated source gases in the stratosphere with the ATMOS instrument, *J.Geophys.Res.* **92**, 9836-9850, 1987.