

Preliminary analysis of PFS/MEx observations of Comet Siding Spring

M. Giuranna (1), O. Witasse (2), A. Aronica (1), S. Aoki (3,4,5), P. Wolkenberg (1,6), M. Costa (7), B. Sánchez-Cano (8)

(1) Istituto di Astrofisica e Planetologia Spaziali (IAPS), Istituto Nazionale di Astrofisica (INAF), Via del Fosso del Cavaliere 100, 00133 Roma, Italy (marco.giuranna@iaps.inaf.it) (2) European Space Agency, Directorate of Science, Noordwijk, The Netherlands (3) Planetary Aeronomy, Belgian Institute for Space Aeronomy, 3 av. Circulaire, B-1180 Brussels, Belgium (4) Fonds National de la Recherche Scientifique, rue d'Egmont 5, B-1000 Brussels, Belgium (5) Department of Geophysics, Tohoku University, Sendai, Miyagi 980-8578, Japan (6) Space Research Centre of Polish Academy of Sciences, Bartycka 18A, Warsaw, Poland (7) European Space Agency, ESAC, Spain (8) Department of Physics and Astronomy, University of Leicester, Leicester, UK

Abstract

On October 19th 2014, Mars experienced a close encounter with Comet C/2013 A1 (Siding Spring), at a distance of 138,000 km. We analyze observations by the Planetary Fourier Spectrometer (PFS) [1] onboard Mars Express performed between October 13th and October 21st 2014 to search for spectral signatures of the comet and to investigate possible effects of its passage on the suspended dust and ice content in the Martian atmosphere.

1. Measurements and datasets

PFS Observations

PFS observations dedicated to the Siding Spring comet flyby were performed between October 13th and October 21st 2014, and in particular around the closest approach (19 October 2014 at 18:28 UT, Ls 217). Given the large FoV of PFS (1.52° FWHM for the Short-wavelength channel, SWC, and 2.69° FWHM for the Long-wavelength channel, LWC; [2,3]), and the large distance of the target, the spectral signatures of the comet, if any, are expected to be very weak in the PFS spectra, and most likely to be observed only around the closest approach. We used the SPICE-enhanced Cosmographia Mission Visualization Tool v3.0 [4,9] for a 3D visualization of Mars Express, Mars and the comet at closest approach (Figure 1). As the two PFS channels have slightly different boresight directions, we used the WebGeocalc online tool [5] with the latest kernel of the comet to investigate when the comet is expected to be within the PFS SWC and LWC FoV. As a

result, for the day of the closest approach, we found four time windows (Table 1).




Figure 1: 3D visualization of Mars Express, Mars and the comet at closest approach with the SPICE enhanced Cosmographia Mission Visualization Tool v3.0. The comet is defined with a sphere of 15,000 km (inner coma). The blue vector represents the PFS SWC nadir boresight.

	Start Time	Stop Time	Duration (secs)
1	2014-10-19 15:31:29.430102 UTC	2014-10-19 15:51:53.160819 UTC	1223.73071754
2	2014-10-19 19:20:26.581536 UTC	2014-10-19 19:21:02.702504 UTC	36.12096786
3	2014-10-20 03:26:38.262334 UTC	2014-10-20 03:57:42.493487 UTC	1864.23115331
4	2014-10-20 20:26:33.985109 UTC	2014-10-20 21:01:19.595671 UTC	2085.61056215

	Start Time	Stop Time	Duration (secs)
1	2014-10-19 15:30:05.761012 UTC	2014-10-19 15:56:53.407226 UTC	1607.64621437
2	2014-10-19 19:20:20.371114 UTC	2014-10-19 19:21:09.958563 UTC	49.58744907
3	2014-10-20 03:26:24.454111 UTC	2014-10-20 03:57:59.363859 UTC	1894.90974790
4	2014-10-20 20:26:24.136971 UTC	2014-10-20 21:02:19.138034 UTC	2155.00106317

Table 1: Time windows, for the day of the closest approach, when the Siding Spring comet is expected to be within the PFS SWC (top panel) and LWC (bottom panel) FoVs.

PFS Retrievals

We use PFS retrievals of suspended dust and water ice opacity [6,7,8] to investigate possible effects of

Comet Siding Spring's passage on the suspended dust and ice content in the Martian atmosphere.

2. Preliminary analysis

Clear spectral signals in the PFS data could only be detected for the MEX orbit 13709, which is exactly when PFS observed Siding Spring around its closest approach. Three spectra can be easily distinguished from the others in both the SWC and the LWC, being well above the instrumental noise level. We can also exclude a random/unusual fluctuation of the PFS spectra, because the three spectra correspond in the two channels (same spectrum number in both channels). We confirm that these spectra were acquired within the time window when the Siding Spring comet was inside the PFS SWC and LWC FoV (Table 1). Given the large distance, the comet is only a small percentage of the PFS FoV, but this seems to be enough to measure some signal in the thermal range. Proper calibration and interpretation of these PFS spectra will be challenging. We performed PFS retrievals for the observations recorded during 126 MEX orbits. We separated all observations performed before 19 October 2014 from those performed after this date. We binned data in 1° of latitude. The results are shown in Figure 2.




Figure 2: PFS dust opacities retrieved in October 2014 before (black crosses) and after (red crosses) the closest approach

Dust Opacities are normalized to the mean Martian surface pressure (6.1 mbar). Black crosses in Figure 2 are for observations performed before the closest approach, while Red crosses are for subsequent observations. These preliminary results indicate an

increase of dust in the Martian atmosphere after the comet's closest approach, especially in the latitude range 40°S - 10°N .

Acknowledgements

The PFS experiment has been built at the Istituto di Astrofisica e Planetologia Spaziali (IAPS) of Istituto Nazionale di Astrofisica (INAF), and has been founded by the Italian Space Agency (ASI) in the context of the Italian participation to the ESA's Mars Express Mission. PFS data are publicly available via the ESA Planetary Science Archive.

References

- [1] Formisano, V., et al., (2005), The Planetary Fourier Spectrometer (PFS) onboard the European Mars Express mission, *Planet. Space Sci.*, 53, 963-974.
- [2] Giuranna, M., et al., (2005a), Calibration of the Planetary Fourier Spectrometer Short Wavelength Channel, *Planet. Space Sci.*, 53, 975-991.
- [3] Giuranna, M., et al., (2005b), Calibration of the Planetary Fourier Spectrometer Long Wavelength Channel, *Planet. Space Sci.*, 53, 993-1007.
- [4] SPICE-Enhanced Cosmographia Mission Visualization Software, <https://naif.jpl.nasa.gov/naif/cosmographia.html>
- [5] <http://spice.esac.esa.int/webgeocalc>
- [6] Grassi, D., et al. (2005), Methods for the analysis of data from the Planetary Fourier Spectrometer on the Mars Express mission. *Planet. Space Sci.* 53 (10), 1017-1034.
- [7] Giuranna, M., et al. (2017), 12 Years of Atmospheric Monitoring by the Planetary Fourier Spectrometer onboard Mars Express. 6th MAMO, January 17-20 2017, Granada, Spain.
- [8] Wolkenberg, P., et al. (2017), Characterization of dust activity on Mars from MY27 to MY32 by PFS-MEX observations, submitted to *Icarus*.
- [9] Costa, M. et al., EPSC 2017.