

Spectral inversion of OMEGA/MEx limb observations considering multiple scattering

Arnaud Mahieux (1,2,3), Shohei Aoki (2,3,4), Masashi Toyooka (4), Hiromu Nakagawa (4), Hironobu Iwabuchi (5), Yasumasa Kasaba (5), Giuseppe Sindoni (6), Anna Geminale (6), Fabrizio Oliva (6), Emiliano D'Aversa (6), Francesca Altieri (6), Giancarlo Bellucci (6), Ann Carine Vandaele (2), Brigitte Gondet (7), Jean-Pierre Bibring (7), Mathieu Vincendon (7), Michael Wolff (7,8)

(1) University of Texas at Austin, Austin, USA (arnaud.mahieux@utexas.edu), (2) Royal Belgian Institute for Space Aeronomy, Belgium, (3) Fonds National de la Recherche Scientifique, Belgium, (4) Department of Geophysics, Tohoku University, Japan, (5) Center for Atmospheric and Oceanic Studies, Tohoku University, Japan, (6) Institute for Space Astrophysics and Planetology, Italy, (7) Institut d'Astrophysique Spatiale, France, (8) Space Science Institute, USA

Abstract

This work presents the results of the OMEGA/Mars Express limb spectral inversion in the infrared, using a Bayesian inversion scheme. We used a multiple scattering radiative transfer code to simulate the light transmission and scattering in the Mars atmosphere. We will present the vertical profiles of the detected species and aerosols, i.e. CO₂, CO, H₂O, dust and ice.

1. Introduction

The OMEGA instrument on board Mars Express (MEx) is an infrared spectrometer taking observations of the Mars atmosphere since 2004 in the 0.35 to 5.1 μm wavelength region [1]. Amongst all the observations, OMEGA observed the Mars atmosphere in limb geometry, during which radiances were measured with a spectral sampling of 7 nm (0.35-1.0 μm), 13 nm (0.93-2.65 μm), and 20 nm (2.51-5.1 μm) and an instantaneous field of view of 1.2 mrad. These measurements show a good coverage of the planet in terms of latitude, longitude and solar longitude (Ls).

In this work, we present an algorithm used to invert the limb observations using a Bayesian approach [2]. In this scheme, the forward model, JACOSPAR, is a full radiative transfer code which accounts for multiple scattering of the Sun light by the atmospheric aerosols, in order to model the radiances with a high precision [3,4,5].

We intend to retrieve CO₂, H₂O and CO vertical profiles for the gas species, and water ice and dust vertical profiles for the aerosol species, which all show clear absorption and/or scattering structure in the considered wavenumber region.

2. The method

2.1 Forward model

JACOSPAR is a multiple scattering radiative transfer code that uses the backward-propagating Monte Carlo method, and the dependent sampling approach in order to reduce the computational time [6]. It calculates the scattering for a given number of wavenumber values and interpolates the radiance for the other wavenumbers [3,4,5]. JACOSPAR accounts for the instrumental field of view in its calculations.

JACOSPAR also computes precise analytical Jacobians relative to the radiances with respect to the absorption and scattering extinction profiles. They are used to derive the Jacobians to volume mixing ratios (VMR) of the different atmospheric gases and aerosols, as well as to the aerosols mean radius, which are used in the Bayesian algorithm.

We compute the aerosols single scattering albedo, phase function and extinction coefficients using the Mie theory [7], for altitude constant modified-gamma size distributions taken from [8], using refractive index of dust and water ice from [9] and [10], respectively.

2.2 Bayesian algorithm

We implemented the Bayesian algorithm approach developed by [2] using the Gauss-Newton method. Based on an a-priori atmosphere obtained from the GEM-MARS [11], we fit the logarithm of the different species VMR and the aerosols mean radius, assuming temperature and pressure conditions obtained from MCD for the latitude, longitude, time and Ls observation mean value.

2.3 Sensitivity study

We present a sensitivity study, conducted in different illumination geometries, to the different retrieved VMR profiles, which shows that the precision of the retrieved profiles is correlated with the distance of the MEx spacecraft to the impact point of the observations due to the field of view of OMEGA, as well as with the vertical sampling (vertical distance between two consecutive impact points). We show that the sensitivity to CO₂, H₂O and dust is reasonable, while is weaker for water ice and CO, due to their respective absorption strength and wavelength dependencies.

3. Results and discussion

We searched for the best covariance values to the a-priori atmosphere in order to best fit the OMEGA measured spectra, and we obtained good fits for the fraction of the observations that have already been inverted.

The vertical profiles derived from a few observations will be presented and discussed in different illumination geometries, geographical position and seasons, and compared to previous observations. We discuss variations of the different VMR vertical profiles as a function of time, latitude and Ls within the retrieved uncertainties, and compare with profiles from models, such as GEM-MARS. In particular, we focus on water and dust vertical profiles.

4. Future applications

The code is intended to be applied to the inversion of the NOMAD [12] on board ExoMars limb observations, that will be carried out from end of 2017. One of the channels of NOMAD, LNO, is a high resolution echelle grating spectrometer using the AOTF technology to select the wavenumber ranges to be measured, working in the infrared from 2.2 to 3.8 μm .

Acknowledgements

The authors would like to thank the UPWARDS Horizon 2020 project for funding. This research was supported by the FNRS CRAMIC project under grant number T.0171.16. MJW acknowledges support of the Paris-Saclay D'Alembert Fellowship Program.

References

- [1] J.-P. Bibring et al., Perennial water ice identified in the south polar cap of Mars, *Nature* 428, 627, 2004.
- [2] C. Rodger, *Inverse method for atmosphere sounding: Theory and Practice*, Oxford University press, 2000
- [3] H. Iwabuchi, *Efficient Monte Carlo Methods for Radiative Transfer Modeling*, *J. Atm. Sc.*, 63, 2006.
- [4] H. Iwabuchi et al., Fast and accurate radiance calculations using truncation approximation for anisotropic scattering phase functions, *JQSRT*, 110, 2009
- [5] A. Mahieux et al., Inversion of Mars atmosphere limb observations considering multiple scattering: Application to OMEGA/MEx observations, (in preparation), 2018
- [6] G. Marchuk et al., *The Monte Carlo Methods in Atmospheric Optics*, Springer-Verlag, 1980
- [7] W. Wiscombe et al., A model for the spectral albedo of snow, *J. Atm. Sc.*, 37, 1980
- [8] A. Kleinbohl et al., Mars Climate Sounder limb profile retrieval of atmospheric temperature, pressure, and dust and water ice opacity, *JGR*, 114, 2009
- [9] S. Warren et al., Optical constants of ice from the ultraviolet to the microwave, *Applied Optics*, 23, 8, 1984.
- [10] M. J. Wolff, et al., "Wavelength dependence of dust aerosol single scattering albedo as observed by the Compact Reconnaissance Imaging Spectrometer.", *Journal of Geophysical Research (Planets)*, 114, E00D04, 2009
- [11] L. Neary and F. Daerden, *The GEM-Mars General Circulation Model for Mars: Description and Evaluation*, Icarus, (accepted), 2017.
- [12] A. C. Vandaele, Science objectives and performances of NOMAD, a spectrometer suite for the ExoMars TGO mission, *PSS*, 119, 2015