

The NOMAD spectrometer suite on ExoMars Trace Gas Orbiter: Data processing status after one year of observing the Red Planet

Ian R. Thomas (1), Ann Carine Vandaele (1), Shohei Aoki (1), Cédric Depiesse (1), Yannick Willame (1), Justin Erwin (1), Loïc Trompet (1), Geronimo Villanueva (2), Giuliano Liuzzi (2), Matteo Crismani (2), Miguel Angel Lopez-Valverde (3), Giancarlo Bellucci (4), Fabrizio Oliva (4), Séverine Robert (1), Arianna Piccialli (1), Bojan Ristic (1), Claudio Queirolo (1), Frank Daerden (1), Sophie Berkenbosch (1), Roland Clairquin (1), Bram Beeckman (1), Eddy Neefs (1), Jon Mason (5), Manish Patel (5), José Juan Lopez Moreno (3) and the NOMAD Team.

(1) BIRA-IASB, Brussels, Belgium; (2) NASA Goddard Space Flight Center, USA; (3) IAA/CSIC, Granada, Spain; (4) IAPS/INAF, Rome, Italy; (5) Open University, U.K. (email: ian.thomas@aeronomie.be)

Abstract

NOMAD is one of four scientific instruments on the ExoMars Trace Gas Orbiter. It consists of three spectrometers operating in the ultraviolet-visible and infrared [2]. The nominal science phase began in April 2018, almost exactly one year ago at the time of writing this abstract; and recently the first results of the TGO mission were published [1, 5].

Here we will describe the first results from the first year of NOMAD observations; with an emphasis on calibration and data processing, detailing the work that has been done so far, and improvements that will be made in future.

1. Introduction

There are two main types of observations performed by NOMAD [4]:

Solar occultations, where the Sun is continually observed as the instrument field of view passes through the atmosphere, have extremely high signal to noise ratios, but can only be performed when the orbital position of the spacecraft allows such observations to be made. Solar occultations are routinely measured in both the infrared (2.2-4.3 μm) and ultraviolet-visible (200-650nm) spectral regions [2, 3].

Dayside nadir observations, where the instrument field of view is pointed towards the surface directly below the spacecraft, and reflected sunlight is

observed. These observations can be measured on almost all orbits, but the trade-off is a lower signal to noise ratio than for solar occultations and therefore higher detection limit for trace gases. Nadirs are typically measured in both the infrared (2.2-3.8 μm) and ultraviolet-visible (200-650nm) spectral regions; although IR observations are usually only made on half of all orbits to keep the instrument at a cooler temperature [4].

In addition to these observing modes, night-side nadirs and limbs are also routinely measured. Until recently, limb measurements have been limited to the infrared only, however it is now possible to measure the Martian limb in both the infrared and ultraviolet-visible at the same time.

2. Results

The spectrometers operate at very high spectral resolution, up to $\sim 0.15\text{cm}^{-1}$ in the infrared and $\sim 1.2\text{nm}$ in the ultraviolet-visible regions [2, 3]; permitting both routine monitoring of gases for climatological studies and high-sensitivity detections of trace gases. Currently CO_2 , CO , H_2O , HDO and O_3 are regularly measured, plus dust and aerosol optical properties; with continued calibration improvements and further studies, new detections could be possible [1, 5].




Figure 1: Attenuation of the SO channel signal as a function of altitude, as measured in solar occultation mode.

In April 2019 the first results from NOMAD and ACS (another spectrometer onboard TGO) were published, showing that (1) no methane has been detected in the Martian atmosphere during the first six months of mapping [1]; (2) D/H ratios and dust vertical profiles can be measured and were affected by the recent global dust storm [5]. As NOMAD continues to routinely observe the atmosphere of Mars, and as the dataset becomes more complete and instrument calibration improves, studies of the composition and ongoing processes in the Martian atmosphere will continue to yield new results. This presentation will highlight the current status of these efforts.

Acknowledgements

The authors would like to thank everyone who has contributed to the design, construction, testing, operations and management of NOMAD and the TGO. ExoMars is a space mission of the European Space Agency (ESA) and Roscosmos. The NOMAD experiment is led by the Royal Belgian Institute for Space Aeronomy (IASB-BIRA), assisted by Co-PI teams from Spain (IAA-CSIC), Italy (INAF-IAPS), and the United Kingdom (Open University). This project acknowledges funding by the Belgian Science Policy Office (BELSPO), with the financial and contractual coordination by the ESA Prodex Office (PEA 4000103401, 4000121493), by the Spanish MICINN through its Plan Nacional and by European funds under grants PGC2018-101836-B-I00 and ESP2017-87143-R (MINECO/FEDER), as well as by UK Space Agency through grant ST/P000886/1 and Italian Space Agency through grant 2018-2-HH.0.

The IAA/CSIC team acknowledges financial support from the State Agency for Research of the Spanish MCIU through the ‘Center of Excellence Severo Ochoa’ award for the Instituto de Astrofísica de Andalucía (SEV-2017-0709). This work was supported by the Belgian Fonds de la Recherche Scientifique – FNRS under grant numbers 30442502 (ET_HOME) and T.0171.16 (CRAMIC) and BELSPO BrainBe SCOOP Project. US investigators were supported by the National Aeronautics and Space Administration.

References

- [1] Korabiev et al.: No detection of methane on Mars from early ExoMars Trace Gas Orbiter observations, *Nature* 568. 2019.
- [2] Neefs et al.: NOMAD spectrometer on the ExoMars trace gas orbiter mission: design, manufacturing and testing of the infrared channels. *Appl. Opt.* 54:28. 2015.
- [3] Patel et al.: NOMAD spectrometer on the ExoMars trace gas orbiter mission: part 2 - design, manufacturing and testing of the UVIS channel. *Appl. Opt.* 56:10. 2017.
- [4] Vandaele et al.: Science objectives and performances of NOMAD, a spectrometer suite for the ExoMars TGO mission. *PSS* 119:15. 2015.
- [5] Vandaele et al.: Martian dust storm impact on atmospheric H₂O and D/H observed by ExoMars Trace Gas Orbiter. *Nature* 568. 2019.