

# Carbon Dioxide and Temperature Profile Retrievals from NOMAD SO Channel on board ExoMars TGO

**Loïc Trompet** (1,2,3), Ann Carine Vandaele (1), Shohei Aoki (1), Justin T. Erwin (1), Ian R. Thomas (1), Geronimo L. Villanueva (4), Giuliano Liuzzi (4), Arnaud Mahieux (1,5,6), Séverine Robert (1), Arianna Piccialli (1), Miguel Angel Lopez-Valverde (7), Brittany Hill (7), Valérie Wilquet (1), Frank Daerden (1), Bojan Ristic (1), Jose Juan Lopez-Moreno (7), Giancarlo Bellucci (8), Manish R. Patel (9) and the NOMAD Team.

(1) Royal Belgian Institute for Space Aeronomy, Belgium, (2) University of Namur, Belgium, (3) Institute of Life, Earth and Environment, Belgium, (4) NASA Goddard Space Flight Center, USA, (5) Fond National de la Recherche Scientifique, Belgium, (6) The University of Texas at Austin, USA, (7) Instituto de Astrofísica de Andalucía, CSIC, Spain, (8) Institute for Space Astrophysics and Planetology, Italy, (9) Open University, UK. ([loic.trompet@aeronomie.be](mailto:loic.trompet@aeronomie.be))

## Introduction

NOMAD (Nadir and Occultation for MArS Discovery) [1, 2] is one of the four instruments on-board the ExoMars Trace Gas Orbiter (TGO). It consists of three spectrometers, SO (Solar Occultation), LNO (Limb, Nadir and Occultation) and UVIS. The SO channel started to make solar occultation measurements on April 21, 2018. It has regularly performed solar occultation measurements of the atmosphere of Mars.

## The NOMAD SO Channel

The SO channel is dedicated to solar occultation measurements: SO sounds both night side and day side of the terminator of Mars. SO is a copy of the SOIR instrument [3] that operated successfully during the entire Venus Express mission. It is an infrared spectrometer working in the 2.2 to 4.3  $\mu\text{m}$  spectral range ( $2325\text{--}4545\text{ cm}^{-1}$ ) with a spectral resolution around  $0.15\text{ cm}^{-1}$ . The instrument is composed of an echelle grating in a near Littrow configuration, and an Acousto-Optical Tunable Filter (AOTF) for the diffraction order selection [4]. TGO is on a quasi-circular orbit at around 400 km of altitude. During a solar occultation, SO scans six diffraction orders each second. Most of these diffraction orders span a spectral range containing carbon dioxide lines. The tangent altitudes where carbon dioxide profiles are retrieved depend on the lines intensity and aerosols presence. The highest altitude bounds are limited by overly weak lines with respect to the signal to noise ratio. The lowest altitudes are limited by either line saturation or dust presence. The later varies mainly with respect to the season and latitude. The different intensities of

carbon dioxide lines amongst the diffraction orders allows a coverage of all altitudes below 200 km.

## Profile Retrievals

ASIMUT-ALVL [5], the radiative transfer program developed at BIRA-IASB, is the program used to retrieve profiles of species from NOMAD spectra. ASIMUT-ALVL is based on the Optimal Estimation Method [6] and includes the analytical calculation of the Jacobians [6]. From carbon dioxide vertical profiles, temperature profiles are derived using the hydrostatic equation and the ideal gas law, similar to the work that was accomplished for the SOIR instrument spectra [7].

The preliminary results of retrieved carbon dioxide and temperature vertical profiles from the SO channel will be presented, discussed, and compared to the results obtained by other instruments as well as models.

## Acknowledgements

ExoMars is a space mission of the European Space Agency (ESA) and Roscosmos. The NOMAD experiment is led by the Royal Belgian Institute for Space Aeronomy (IASB-BIRA), assisted by Co-PI teams from Spain (IAA-CSIC), Italy (INAF-IAPS), and the United Kingdom (Open University). This project acknowledges funding by the Belgian Science Policy Office (BELSPO), with the financial and contractual coordination by the ESA Prodex Office (PEA 4000103401, 4000121493), by the Spanish Ministry of Science and Innovation (MCIU) and by European funds under grants PGC2018-101836-B-I00 and ESP2017-87143-R (MINECO/FEDER), as well as by UK Space Agency through grant

ST/R005761/1 and Italian Space Agency through grant 2018-2-HH.0. The IAA/CSIC team acknowledges financial support from the State Agency for Research of the Spanish MCIU through the ‘Center of Excellence Severo Ochoa’ award for the Instituto de Astrofísica de Andalucía (SEV-2017-0709). This work was supported by the Belgian Fonds de la Recherche Scientifique – FNRS under grant numbers 30442502 (ET\_HOME) and T.0171.16 (CRAMIC) and BELSPO BrainBe SCOOP Project. US investigators were supported by the National Aeronautics and Space Administration.

## The NOMAD Team

*Scientific team:* Vandaele, Ann Carine; Lopez Moreno, Jose Juan; Bellucci, Giancarlo; Patel, Manish; Allen, Mark; Alonso-Rodrigo, Gustavo; Altieri, Francesca; Aoki, Shohei; Bauduin, Sophie; Bolsée, David; Carrozzo, Giacomo; Clancy, Todd; Cloutis, Edward; Crismani, Matteo; Daerden, Frank; Da Pieve, Fabiana; D’Aversa, Emiliano; Depiessé, Cédric; Erwin, Justin; Etiopa, Giuseppe; Fedorova, Anna; Formisano, Vittorio; Funke, Bernd; Fussen, Didier; Garcia-Comas, Maia; Geminale, Anna; Gérard, Jean-Claude; Giuranna, Marco; Gkouvelis, Leo; Gonzalez-Galindo, Francisco; Hewson, Will; Holmes, James; Hubert, Benoît; Ignatiev, Nicolai; Kaminski, Jacek; Karatekin, Ozgur; Kasaba, Yasumasa; Kass, David; Kleinböhl, Armin; Lanciano, Orietta; Lefèvre, Franck; Lewis, Stephen; Liuzzi, Giuliano; López-Puertas, Manuel; López-Valverde, Miguel; Mahieux, Arnaud; Mason, Jon; Mc Connell, Jack; Mège, Daniel; Mumma, Mike; Nakagawa, Hiromu; Neary, Lori; Neefs, Eddy; Novak, Rob; Oliva, Fabrizio; Piccialli, Arianna; Renotte, Etienne; Ritter, Birgit; Robert, Severine; Schmidt, Frédéric; Schneider, Nick; Sindoni, Giuseppe; Smith, Michael D.; Teanby, Nicolas A.; Thiemann, Ed; Thomas, Ian; Trokhimovskiy, Alexander; Trompet, Loïc; Vander Auwera, Jean; Villanueva, Geronimo; Viscardy, Sébastien; Whiteway, Jim; Willame, Yannick; Wilquet, Valérie; Wolff, Michael; Wolkenberg, Paulina, Yelle, Roger – *Tech team:* Alonso-Rodrigo, Gustavo; Aparicio del Moral, Beatriz; Barzin, Pascal; Beeckman, Bram; BenMoussa, Ali; Berkenbosch, Sophie; Biondi, David; Bonnewijn, Sabrina; Candini, Gian Paolo; Clairquin, Roland; Cubas, Javier; Giordanengo, Boris; Gissot, Samuel; Gomez, Alejandro; Hathi, Brijen; Jeronimo Zafra, Jose; Leese, Mark; Maes, Jeroen; Mazy, Emmanuel; Mazzoli, Alexandra; Meseguer, Jose; Morales, Rafael; ; Orban, Anne; Pastor-Morales, M; Perez-grande, Isabel; Queirolo, Claudio; Ristic, Bojan; Rodriguez Gomez, Julio; Saggin, Bortolino; Samain, Valérie; Sanz Andres, Angel; Sanz, Rosario; Simar, Juan-Felipe; Thibert, Tanguy

## References

- [1] Vandaele, A.C., et al.: Science objectives and performances of NOMAD, a spectrometer suite for the ExoMars TGO mission, *Planet. Space Sci.*, Vol. 119, pp 233-249, 2015.
- [2] Neefs, E., et al.: NOMAD spectrometer on the ExoMars trace gas orbiter mission: part I – design, manufacturing and testing of the infrared channels, *Appl. Opt.*, Vol. 54(28), pp 8494-8520, 2015.
- [3] Nevejans, D., et al.: Compact, high resolution space borne echelle grating spectrometer with AOTF based on order sorting for the infrared domain from 2.2 to 4.3 micrometer, *Appl. Opt.*, Vol. 45, pp 5191-5206, 2006.
- [4] Thomas, I.R., et al.: Optical and radiometric models of the NOMAD instrument part II: the infrared channels – SO and LNO, *Opt. Express*, Vol. 24(4), pp 3790-3805, 2016.
- [5] Vandaele, A.C., Kruglanski, M., De Mazière, M., Modeling and retrieval of atmospheric spectra using ASIMUT. Proceedings of the First Atmospheric Science Conference, ESRIN, Frascati, Italy, 2006.
- [6] Rodgers, C.D., Inverse method for atmospheric sounding: theory and practice. Hackensack, N.J. (Ed.), World Scientific University of Oxford, Oxford, 2000.
- [7] Mahieux, A. et al.: Densities and temperatures in the Venus mesosphere and lower thermosphere retrieved from SOIR on board Venus Express: Carbon dioxide measurements at the Venus terminator, *J. Geophys. Res.*, Vol 117, E07001, 2012.