

THE UV CLIMATOLOGY IN BELGIUM FROM UV FIELD MONITORING.

Didier Gillotay, and David Bolsée
Institut d'Aéronomie Spatiale de Belgique
Avenue Circulaire, 3, B-1180 Brussels, Belgium

ABSTRACT

Since the end of the 80's, the Belgium Institute for Space Aeronomy (IASB) has developed an automatic station to measure continuously the UV (UV-B & UV-A) – Visible Solar irradiance (280-600 nm) at the Earth's surface.

The IASB monitoring station consists in a combination of instruments including spectro-radiometer, filter-radiometers and broadband radiometers providing absolute values of the total, direct and diffuse components of the Solar irradiance. The IASB data set is completed by ancillary measurements e.g; Ozone and SO₂ total column, Ozone concentration profiles, meteorological conditions...mainly provided by the Royal Meteorological Institute of Belgium (KMI/IRM). From the available 11-years period of continuous measurements, it is possible to define the major characteristics of the UV climatology in Belgium and by extension in the 50° - latitude area.

1. INTRODUCTION

The penetration of solar UV radiation through the atmosphere depends on the solar zenithal angle (SZA), the ozone overhead column and other atmospheric absorbers and scatterers such as clouds and aerosols. In particular, clouds are responsible for a great deal of the observed irradiance variability. The interpretation of observed UV-B time series, and e.g. the detection of possible trends due to human activity, requires the correct understanding of the effects of these different 'factor of influence' and a detailed study of their evolution with time.

The instrumentation is described in the next section. Total ozone is measured at Uccle by the Royal Meteorological Institute (KMI/IRM) using a Dobson and a Brewer spectroradiometer, [De Muer and De Backer, 1992]. Ozone, temperature and relative humidity profiles are obtained by balloon soundings, also provided by KMI/IRM. The cloud fraction and type as well as the ground meteorological parameters (pressure, temperature, horizontal visibility,...) are monitored routinely by KMI/IRM.

UV measurements in Uccle (Brussels) Belgium are available since April 1989 by combining the IASB measurements and data from KMI/IRM.

The major results are presented and discussed in terms of correlation between the UV-B irradiance and the main atmospheric parameters like Ozone, SO₂, Clouds

cover, Aerosols... Some preliminary results on potential trends are also presented and discussed.

2. EXPERIMENTAL

2.1 Ground based monitoring station

The IASB-automated station is located at Uccle, a residential area in the Brussels suburbs (lat:50°47'54''N, long:4°21'29''E, Alt.:105m asl). It is operational since mid-march 1993 [Gillotay, 1996]. The core instrument of the main station is a double monochromator (modified HD10, Jobin-Yvon).

It includes also three filter radiometers (SPUV-10, UVMFR-7 from Yankee Environmental System, YES and GUV 511C from Biospherical Instruments) and four pyranometers (YES), two in the UV-B range (UVB-1), one in the UV-A (UVA-1) and the last covering the wavelength range from the UV-A up to the near IR (TSP-700).

The spectro-radiometer, with its optical axis pointing the zenith direction, is fitted with a Lambertian Teflon diffuser with a 2πsr field of view, measures the total solar irradiance (diffuse + direct), and a nearly perfect cosine response. One scan is performed every 15 minutes for SZA smaller than 100°.

The 10-channels filter radiometer (SPUV-10, YES) measure the direct solar irradiance from 300 nm to 1040 nm. It is mounted on a sun tracking system (INTRA, Brusag). This radiometer is designed to provide direct solar irradiance measurements from which the ozone total column and the atmospheric turbidity (the optical depth of aerosols in clear sky conditions) can be deduced.

GUV 511C and UVMFR-7 are respectively 5 and 7 channels 2πsr filter radiometers. The UVMFR-7 equipped with a shadowing band, is designed to perform direct and diffuse quasi simultaneous measurements of the solar irradiance, from which complementary informations on ozone and aerosols can be deduced.

The pyranometers cover the full range of the solar spectra scanned by the monochromators, with a much higher time sampling (1 mean integrated measurement every minute). One of the UV-B meters is shadowed in order to measure the diffuse component of the solar irradiance.

Finally, KMI/IRM perform UV-B measurements with a Brewer (Mk II) single monochromator from 280

to 325 nm, initially a noon (from April 1989 to December 1990) and more recently on an hourly base (January 1991 – today).

TABLE 1: Main Characteristics of the instruments.

SPECTRORADIOMETER

INSTRUMENT	SPECTRORADIOMETER
COMPANY	Jobin-Yvon.
TYPE	Modified HD10
FOCAL LENGTH (mm)	100
GRATINGS	Holographic Concave
Lines/mm	1200
FWHM at 300 nm (nm)	0.488
USUAL STEP (nm)	0.5
USUAL RANGE (nm)	From 280 to 600 nm
SCAN DURATION (s)	400
FIELD OF VIEW	$2\pi sr$
DIFFUSER	Teflon
DETECTOR	PMT
type	Hamamatsu R292

YES PYRANOMETERS

	UV-B METER	UV-A METER	TOTAL
	UVB-1	UVA-1	TSP-1
RANGE (nm)			
from	280	300	300
to	330	380	3000
maximum	294	330	

FILTER RADIOMETERS

SPUV-10		GUV 511C		UVMFR-7	
WL (nm)	FWHM (nm)	WL (nm)	FWHM (nm)	WL (nm)	FWHM (nm)
300	2	305	2	300	2
310	2	320	4	305.5	2
318	2	340	4	311.5	2
368	2	380	4	317.5	2
501	10	PAR		325	2
615	10			332.5	2
675	10			368	2
781	10				
871	10				
1039	20				

2.2 Calibration and quality control of the data

Periodical absolute calibration is performed in a dark room using five different NIST-FEL 1000 W standard lamps. Furthermore, stability is periodically checked by means of a Transportable Lamp System (TLS) developed specifically in our laboratory. It consists of five 200 W quartz-halogen lamps and a

Mercury low pressure source, mounted on a carousel inside a movable container. In the field, the different lamps are successively placed and automatically aligned with the entrance optics of the instruments. With both 'standards' the uncertainties can be estimated to be less than $\pm 5\%$ on all the wavelength range. This estimation was confirmed during the previous European Inter-comparison Campaign. [Gardiner *et al.*, 1993]. Moreover; the coherency of the data set is verified by comparing the filter radiometer and broadband measurements with the corresponding convoluted spectral measurements.

2.3. Time series of measurements

Erythemal doses at noon in Uccle are evaluated from both sets of spectral UV-Visible measurements, by weighting each spectrum by the CIE action spectrum. [McKinlay and Diffey, 1987]. The KMI/IRM data set is corrected to take into account the lack of spectral measurements between 325 and 400 nm. The comparison of the two data sets gives a good agreement (within 5%) for most of the cases over the overlap period (1993-1999). Nevertheless, in some occasions, the discrepancy can reach 20-25%. This is probably due to 1) the unperfected synchronism between the measurements and 2) the correction of the Brewer measurements which does not take into account the modification of the cloud cover during one scan duration.

Figure 1 illustrates the available time series and shows their seasonal variation. The peak values are achieved in June, corresponding to the smallest SZA of the year and relatively low ozone columns.

The scatter within the seasonal fluctuation can be ascribed to changes in cloud coverage.




Figure 1. Time series of the erythemal doses at Uccle.

3. FACTORS OF INFLUENCE

The two most important factors limiting the penetration and explaining the day to day variations of the UV-B radiation to the Earth's surface are the ozone and the cloud coverage. These two 'factors of influence' will be detailed in the next sections. Other factors like aerosols, which are relatively constant in Brussels, will be neglected in this paper.

3.1 Ozone.

Figure 2 illustrates the anti-correlation between ozone total column and UV-B integrated irradiance corrected for the effect of cloud cover. The applied correction is simply the ratio UV-B/UV-A that takes into account, as a first approximation, the effect of clouds as a neutral filter.




Figure 2. Anti-correlation between ozone and UV-B

A discrete ordinates radiative model [Stamnes *et al.*, 1988] has been used to simulate the experimental data and verify the anti-correlation between ozone and UV-B.

The extraterrestrial flux is a combination of the SUSIM spectrum below 350 nm [Van Hoosier *et al.*, 1984] and the Neckel and Labs spectrum [Neckel and Labs, 1984] up to 600 nm.

The wavelength dependence of the aerosol optical properties follows the parametrisation of WCP [WCP, 1986] for typical continental mixtures. This choice is motivated by a air pollution lower in Uccle than in typical urban centres. The weak dependence of cloud extinction and asymmetry factor is parametrised following [Slingo, 1989].

A good agreement (better than 5%) between experimental data and the simulation has been established for SZA between 30° and 70° in clear sky condition. The discrepancies between modelled and experimental data increase generally with the SZA and can exceed 10% at high SZA in the visible range.

Figure 2 shows clearly that the anti-correlation observed experimentally is well reproduced by modelling and that practically all the experimental conditions are included

in $\pm 10\%$ limit compared to the predicted anti-correlation. This 10 % variation can easily be explained by considering the error in the ozone measurements (5%) and the unsophisticated correction of the cloud layer effects.

3.2 Clouds

In order to investigate the role of clouds as a function of wavelength, average spectra for well-defined conditions (complete overcast, similar zenith angles) have been derived from the observations, and compared with a corresponding clear sky spectrum. The average cloud transmission ratios for SZA=30° are displayed on figure 3, and compared to a modelled transmission ratio. A 1-km low cloud with an optical depth equal to 50 has been assumed.




Figure 3. Ratio of cloudy (8 Octas) to clear sky irradiance..

Despite the large variability of the cloud impact, a consistent picture is found. The attenuation is lowest in the UV-A, and highest in the ozone absorption bands (UV-B) because of the increased multiple scattering and tropospheric ozone absorption caused by cloud.

The attenuation increases into a lesser extent in the visible range, reflecting the lesser importance of Rayleigh diffusion at higher wavelengths.

Finally, the average attenuation of sunlight by different type of clouds can be also directly estimate from the pyranometers data. As expected, the attenuation by cirrus clouds (high altitude) is found to be very small. In contrast, low clouds (mainly stratocumulus) reduce solar irradiance by about a factor 5 on average.

This attenuation is found to increase monotonously with the solar zenith angle in the UV-A and UV-B ranges, but not for the total integrated irradiances (300-3000 nm). These last results have to be examined in the more detailed future modelling studies.

4. TRENDS

The bring to light of potential trends of UV-B radiation at the Earth's surface due to human activity is a high interest for the public health medical community as well as for all the scientifics interested in the effects of UV-B on biology and material sciences.

The aim of this chapter is just to illustrate what can be deduced from a 11-years period of UV-B monitoring. Figures 4 illustrates the high variability of UV-B effective doses on a monthly bases mainly due to the variability of meteorological conditions.




Figure 4. UV-B effective monthly averaged doses in Uccle (Brussels) Belgium.

Figure 5 and 6 give a first idea of the potential trends of UV-B and ozone in Brussels.




Figure 5. UV-B trends in Brussels 1989-2000.

UV-B trends shows an increase of 0.6% a year that looks coherent with the ozone trends of 0.3%.

4. CONCLUSION

These results show the consistency of both our model and experimental data. They provide a first understanding of the UV-B climatology in Belgium that could be extrapolate to the 50° latitude area. An extended period of measurements is necessary to improve the preliminary trends given above. Nevertheless the increase of UV-B radiation seems to be real and needs to be explored in more detail.




Figure 6. Ozone trends in Brussels (1978-2000) by combining TOMS and KMI/IRM data.

ACKNOWLEDGMENTS

The authors are very grateful to Drs De Muer and De Backer for the providing ozone and UVB data.

REFERENCES

De Muer, D., and H. De Backer, 1992, Revision of 20 years of Dobson total ozone data at Uccle (Belgium): Fictitious Dobson total ozone trends induced by sulfur dioxide trends, *J. Geophys. Res.*, 97, 5921-5937.

Gardiner B.G., Webb A.R., Bais A. F., Blumthaler M., Dirmhirn I., Forster P., Gillotay D., Henriksen K., Huber M., Kirsch P.J., Simon P.C., Svenoe T., Weihs P. and Zerefos C.S., 1993, European intercomparison of ultraviolet spectroradiometers, *Environ. Technol.*, 14, 25-43.

Gillotay D., 1996, UV monitoring in Belgium: Past, present and future. in *Measurements and Trends of Terrestrial UVB Radiation in Europe*, (Ed. B. L. Diffey), pp. 41-53.

Neckel, H., and D. Labs, 1984, The Solar Radiation Between 3300 and 12500 A, *Solar Physics*, 90, 205-258.

McKinlay, A;F; and B.L. Diffey 1987, A reference action spectrum for Ultraviolet induced erythema in human skin. *CIE J.* 6 p; 17-22

Stamnes, K., S. C. Tsay, W. J. Wiscombe, and K. Jayaweera, 1988, Numerically stable algorithm for discrete-ordinate-method radiative transfer in multiple scattering and emitting layered media, *Appl. Opt.*, 27, 2502-2509.

Van Hoosier, M., J.-D. Bartoe, G. Brueckner, and D. Prinz, 1988, Absolute solar spectral irradiance 120 nm - 400 nm (Results from the Solar Ultraviolet Spectral Irradiance Monitor - SUSIM- Experiment on board Spacelab 2), *Astro. Lett. and Communications*, 27, 163-168.

World Climate Programme (WCP), 1986, A preliminary cloudless standard atmosphere for radiation computation, *WMO Report WMO/TD-No. 24*, World Meteorological Organisation.