



Validation of the CAMS regional services: concentrations above the surface

Status update for the period
December 2019 – February 2020

Issued by: KNMI

Date: 4 June 2020

Ref: CAMS84_2018SC2_D4.1.1-DJF2020.pdf

This document has been produced in the context of the Copernicus Atmosphere Monitoring Service (CAMS). The activities leading to these results have been contracted by the European Centre for Medium-Range Weather Forecasts, operator of CAMS on behalf of the European Union (Delegation Agreement signed on 11/11/2014). All information in this document is provided "as is" and no guarantee or warranty is given that the information is fit for any particular purpose. The user thereof uses the information at its sole risk and liability. For the avoidance of all doubts, the European Commission and the European Centre for Medium-Range Weather Forecasts has no liability in respect of this document, which is merely representing the authors view.



Validation of the CAMS regional services: concentrations above the surface

Status update for the period
December 2019 – February 2020

AUTHORS:

D. Akritidis (AUTH), T. Antonakaki (AA), Y. Bennouna (CNRS-LA),
A.-M. Blechschmidt (IUP-UB), T. Bösch (IUP-UB), H. Clark (CNRS-LA),
C. Gielen (BIRA-IASB), F. Hendrick (BIRA-IASB), J. Kapsomenakis (AA),
S. Kartsios (AUTH), E. Katragkou (AUTH), D. Melas (AUTH), A. Mortier (MetNo),
E. Peters (IUP-UB), K. Petersen (MPI), A. Piters (KNMI), A. Richter (IUP-UB),
M. van Roozendaal (BIRA-IASB), M. Schulz (MetNo), N. Sudarchikova (MPI),
A. Wagner (MPI), P. Zanis (AUTH), C. Zerefos (AA)

EDITORS:

J. Douros (KNMI), H.J. Eskes (KNMI)

**REPORT OF THE COPERNICUS ATMOSPHERE MONITORING SERVICE,
VALIDATION SUBPROJECT (CAMS-84).**

CITATION:

Douros, J., H.J. Eskes, D. Akritidis, T. Antonakaki, Y. Bennouna, A.-M. Blechschmidt, T. Bösch, H. Clark, C. Gielen, F. Hendrick, J. Kapsomenakis, S. Kartsios, E. Katragkou, D. Melas, A. Mortier, E. Peters, K. Petersen, A. Piters, A. Richter, M. van Roozendaal, M. Schulz, N. Sudarchikova, A. Wagner, P. Zanis, C. Zerefos, Validation of CAMS regional services: concentrations above the surface, Status update for December 2019 - February 2020, Copernicus Atmosphere Monitoring Service (CAMS) report, CAMS84_2018SC2_D4.1.1-DJF2020, 4 June 2020, doi:10.24380/2510-0365.

STATUS:

Version 1, final

DATE:

4 June 2020

REF:

CAMS84_2018SC2_D4.1.1-DJF2020



Executive Summary

The Copernicus Atmosphere Monitoring Service (CAMS, <http://atmosphere.copernicus.eu>) is a component of the European Earth Observation programme Copernicus. The CAMS service consists of two major forecast and analysis systems. First, the CAMS global near-real time (NRT) service provides daily analyses and forecasts of reactive trace gases, greenhouse gases and aerosol concentrations, and is based on the ECMWF Integrated Forecast System (called CAMS-global in this document). Secondly, seven regional models (9 models starting from 16 October 2019) in Europe perform air quality forecasts and analyses on a daily basis, nested within CAMS-global. Based on these individual forecasts and analyses, an ensemble forecast of air quality over Europe is produced and disseminated by Météo-France (called ENSEMBLE or CAMS-regional below). The regional members use the global forecasting results as boundary conditions at the sides and top of the domain.

This document reports on two validation activities, namely

- an evaluation of the consistency between the global and regional modelling components of CAMS, focussing on the boundaries of the regional domain, and
- an evaluation of the regional ensemble and the seven individual models contributing to the ensemble with independent observations, focusing on the concentrations above the surface.

The current analysis includes ozone (O₃), nitrogen dioxide (NO₂), aerosol (PM₁₀/PM_{2.5}/AOD) and carbon monoxide (CO) forecasts covering the period up to February 2020.

The forecasts from the regional models were compared with the following set of observations:

- aerosol lidar observations from the EARLINET network;
- aerosol AOD observations from the AERONET network;
- IAGOS routine aircraft measurements of ozone and CO;
- ozone sonde profiles;
- MaxDOAS NO₂ tropospheric columns;
- GOME-2/MetOp-A NO₂ satellite tropospheric column retrievals (IUP-UB v1.0 product);
- high-altitude ozone surface stations;
- CO and O₃ from GAW mountain stations; and
- CO observations from the MOPITT satellite instrument.

These observations are available to CAMS within one month after the observations were made.

This report is based on regional model data available for the months February 2016 to February 2020, with a focus on December 2019 – February 2020 (DJF2020). The report is updated every 3 months. The main results are summarised below. This summary is focusing on the performance of the regional Ensemble. Detailed results, also for the individual models, are presented in sections 3 to 11, and each of these sections starts with a summary of the main results. Model specific findings are summarised in section 12. For the period up to February 2020 only data from the 7 original ensemble members was available to us.




Figure S.1. CAMS global ozone forecast for day 1 (left), CAMS regional ENSEMBLE ozone forecast for day 1 (middle), CAMS regional ENSEMBLE ozone analysis (right). From top to bottom: 0, 250, 2000, 5000m altitude level. The results are averaged over the December 2019 - February 2020 period.

During 2019 there was an upgrade implemented for CAMS-global (on 9 July 2019, moving from 60 to 137 vertical levels), while on 4 February 2020 there was an upgrade for the CAMS regional models, implementing new anthropogenic emissions and making available three new aerosol species.

General conclusions for the regional ENSEMBLE forecasts

The comparison of the European regional CAMS ENSEMBLE air quality forecasts and analyses against above-surface observations of O₃, NO₂, CO for the period up to 1st of March 2020 demonstrates that overall, the biases observed are small, often within the uncertainty of the validation approach while temporal correlations for ozone and CO are reasonable. Performance of the ENSEMBLE analysis product is found to be generally superior to that of the ENSEMBLE forecasts. Regional models - and thus the ENSEMBLE - benefit from the use of the global CAMS boundary conditions, which are implemented efficiently. The ENSEMBLE performs generally better than any of the individual models for ozone, NO₂ and CO, showing the strength of the ensemble approach adopted in CAMS.




Figure S.2. Normalised bias (left) and correlation coefficient (right) for ozone for the high-altitude (above 1 km) EEA Air Quality e-reporting stations (top) and the 5 high-altitude European GAW stations (bottom). Lines represent ENSEMBLE forecast (solid red), ENSEMBLE analysis (solid green) and CAMS-global system (blue) for December 2019 - February 2020. The horizontal axis is the station identifier referring to Hohenpeissenberg (HPB), Jungfrauoch (JFJ), Monte Cimone (CMN), Sonnblick (SNB) and Zugspitze (ZUG).




Figure S.3. Modified normalised mean bias (MNMB) against ozone sondes for the regional ENSEMBLE forecasts (red) and analyses (green) from February 2019 to February 2020 (horizontal axis). Ozone was averaged over the lower-middle free troposphere region, 500 hPa < p < 850 hPa.




Figure S.4. Left: Daily time series of ozone at Frankfurt for December 2019 - February 2020. The regional ENSEMBLE forecast is shown in red, regional analyses in green, CAMS-global in blue and the IAGOS aircraft observations in black. The black thin line is the monthly mean of the observations (MOZAIC) over the period 2003-2019, the black dashed lines show 1 standard deviation from the monthly mean and the black dotted-dashed line shows 3 standard deviations from the monthly mean. Right: Mean IAGOS aircraft profile of ozone at Frankfurt for the period December 2019 - February 2020. The dashed black line is the standard deviation of the observations during December-February from the years 2003-2019.

Ozone

The differences between the global and regional systems reveal themselves in the boundary layer and at the surface over land, as expected. A comparison of the regional analysis product with the regional day 1 forecast (Fig. S.1) shows some differences between the regional ENSEMBLE forecast and analysis, with the analysis generally having lower concentrations near the surface. At the boundaries of the regional domain, the ENSEMBLE agrees well with CAMS-global, indicating that the implementation of the boundary conditions was done properly.

For high altitude stations, Fig. S.2, mostly an overestimation between -5% and 35% is observed for the ENSEMBLE first day forecast, while comparison against ozonesondes shows a milder overestimation in the 5-20% range (also Fig. S.3). Time correlations at high-altitude and GAW stations range between 0.25 and 0.8 during this period, with noticeably improved figures for the ENSEMBLE analyses as compared to the ENSEMBLE forecast in the case of the high-altitude stations. The CAMS regional forecast and analysis perform equally when compared with the GAW station observations.




Figure S.5. Maps of satellite retrieved and model simulated tropospheric NO₂ columns [molecules cm⁻²] for February 2020 for GOME-2 (top left), regional ENSEMBLE forecasts (top right), CAMS-global forecasts (middle left) and regional ENSEMBLE analyses (middle right) The panel at the bottom shows corresponding time series of average tropospheric NO₂ columns [10¹⁵ molecules cm⁻²] from GOME-2 (black), regional ENSEMBLE forecasts (red), CAMS-global forecasts (blue) and regional ENSEMBLE analyses (green). GOME-2 data were gridded to regional model resolution (i.e. 0.1° x 0.1°). Model data were treated with the same reference sector (25°W - 20°E) subtraction approach as the satellite data and linearly interpolated to the satellite overpass time (9:30 LT).

Comparisons with IAGOS aircraft observations, see Fig. S.4, reveal that ozone is overestimated by both the regional ensemble and CAMS-global, consistent with the sonde and high-altitude station observations. The comparison with the climatological variability shows that no major episode was observed in the low troposphere at Frankfurt during DJF2020. However, several ozone values in the surface and boundary layer are close to, or slightly greater than, one standard deviation from climatology.




Figure S.6: Diurnal cycles (averages over hourly bins) of tropospheric NO₂ VCDs [10¹⁵ molec. cm⁻²] from MAX-DOAS and models for De Bilt (Netherlands) (top left), Uccle (Belgium) (top right) and Bremen (Germany) (bottom). The coloured lines show (black) MAX-DOAS retrievals, (red) regional ENSEMBLE forecasts and (blue) CAMS-global. Period: June 2016 – February 2020.

The averaged model profiles (Fig. S.4, right) reveal overestimations across the whole vertical extent of the measurements, more so above 500m. ENSEMBLE forecasts and analyses produce improved estimates compared to CAMS-global between 500-2000m.

Nitrogen dioxide (NO₂)

The overall spatial distribution of tropospheric NO₂ as observed from space by GOME-2 is reproduced by the ensemble during DJF 2020 (see Fig. S.5), given known uncertainties in the satellite observations over Central Europe in winter and in January in particular. As described in previous reports, winter values over European emission hotspots simulated by the regional ensemble analysis, forecasts and CAMS-global show significantly smaller values than GOME-2. Values over European emission hotspots are better represented by the regional ensemble analysis and forecasts than by CAMS-global. As a result regional model upgrades in June 2019 and February 2020, which include the use of updated European emissions inventory with improved estimates for anthropogenic emissions in North Africa and the Middle East, enhanced tropospheric columns of NO₂ are now reproduced by the ensemble model runs over these regions, revealing most large urban areas in Egypt, Lebanon and Israel.




Figure S.7. Extinction profiles December 2019 - February 2020 derived from the ENSEMBLE forecast mass concentration profiles (red envelope) and from EARLINET (climatology) backscatter profiles (grey envelope: lidar ratio uncertainty, light grey: including sampling error). “n: XX means number of individual EARLINET profiles assembled (December-February 2006-2020). The EMC used for the calculation of the extinction from the concentration profiles is indicated for each station below the number of EARLINET profiles “n” used for the calculation of the climatology.

Systematic uncertainties in the retrievals (on average on the order of 20% – 30% over polluted regions) depend on the season, with winter values in mid and high latitudes normally associated with larger error margins. Conclusions may differ for comparisons to other satellite NO₂ products (e.g. TEMIS GOME-2, <http://www.temis.nl>). We note that since the CAMS-global upgrade of 26 June 2018, GOME-2 observations are assimilated by the global system. This is, however, a different retrieval product than what is used in the validation reported here (University of Bremen retrieval).




Figure S.8. CAMS global PM10 forecast for day 1 (left), CAMS regional ENSEMBLE PM10 forecast for day 1 (middle), CAMS regional PM10 analysis (right). From top to bottom: 0, 250, 2000, 5000m altitude level. Period: December 2019 to February 2020.

Comparisons to ground based remote sensing MAX-DOAS retrievals at three different European stations (see Figure S.6) show that regional ENSEMBLE forecasts are closer to the urban station observations than CAMS-global, mainly attributed to the difference in spatial resolution. The performance of simulations for diurnal cycles of tropospheric NO₂ columns depends on the location, but generally shows a good performance for the ENSEMBLE products.

Aerosol / PM

The regional models are compared with EARLINET climatological lidar profiles for the same season (data from 2006-2019), Fig. S.7. The standard dissemination of CAMS-regional forecasts does not include information on composition, size and humidity growth of the aerosol in the models. This introduces considerable uncertainty to the PM derived extinction, which conservatively spans a factor 10 for absolute extinction values. Relative differences among nearby stations and the form of extinction profiles are more certain. The order of magnitude in extinction is similar between the models and the lidar profiles, with Ispra, Italy, close to the Alps, and Granada, Spain, being the most notable exceptions. The decrease in extinction with height seems to be generally steeper in the observations, such that upper level extinction in the models is higher in some locations.




Figure S.9. MNMBs [%] (top) and correlation coefficients (bottom) for CO regional ENSEMBLE forecasts (red), analyses (green) and CAMS-global (blue) compared to observations at GAW stations. Period: December 2019 - February 2020.

The PM₁₀ concentrations in the regional ENSEMBLE forecasts and analyses are similar, but larger PM values are observed for the analysis over some areas of Central and Western Europe at the surface level, see Figure S.8. CAMS-global continues to show higher dust loads over the Mediterranean. An increased high bias in PM₁₀-dust over the Mediterranean appeared with the upgrade of CAMS-global to 46R1 in July 2019 (Wagner et al., 2020). Also, notable hotspots are seen over Spain, Sardinia, Sicily and Turkey, similar to what was found during the previous quarters. The agreement between CAMS-global and the ENSEMBLE products is better for PM₁₀ and PM_{2.5} in the upper layers compared to the lower layers, as expected. For PM_{2.5} the difference between the CAMS-global and CAMS-regional is more pronounced than for PM₁₀ within the PBL especially for dust near the southern boundary and over the Middle East (exceeding 50 µg/m³).

Carbon monoxide (CO)

Comparison at the GAW stations reveal biases between 8% and -9% for the analysis and between 1% and -12% for the forecast, while temporal correlations coefficients are between 0.4 and 0.66 for the analysis and between 0.31 and 0.86 for the forecast (Fig. S.9). The comparison against IAGOS aircraft observations shows that, in the lower layers, CO is mostly underestimated by all models with, in some cases, a better performance from the regional ENSEMBLE. In the free troposphere CAMS-global performs well, while the regional ensemble underestimates.

Comparisons with MOPITT CO satellite observations (version 8, Fig. S.10) data show slight underestimation of CO values (~10%) with some regional and temporal variability. The models capture well both the relatively high CO values over the northern part of the domain as well as the relatively low values over the southern part. The ENSEMBLE analysis shows similar patterns compared to the ENSEMBLE forecast only with slightly stronger biases.




Figure S.10. CO total column for MOPITT v8 satellite retrievals (top row, in molecules/cm²), relative difference between the regional ENSEMBLE forecasts and MOPITT (middle row) and regional ENSEMBLE analyses and MOPITT (bottom row) for December 2019 (left column), January (middle column) and February 2020 (right column). Grey colour indicates missing values.



Table of Contents

Executive Summary	4
1 Introduction	16
2 Regional and global CAMS forecasting systems	17
2.1 Regional models	17
2.2 Global CAMS system based on the ECMWF IFS model	20
3 Consistency between the global and regional modelling components of CAMS	22
3.1 Summary	22
3.2 Methodology for the comparisons of CAMS-global and CAMS-regional	22
3.3 Consistency between the global and regional forecasts	27
3.4 Regional variability	28
3.5 Time series	33
3.6 Diurnal cycles	35
3.7 Regional domain boundary cross sections	37
3.8 Regional analysis vs. regional forecasts	42
4 Vertical profile and column aerosol comparisons	47
4.1 Summary for the EARLINET lidar and Aeronet comparisons	47
Introduction	47
4.2 Methodology	48
4.3 Results	51
4.3.1 Comparison of extinction profiles	51
4.3.2 Seasonal variability	54
5 IAGOS aircraft CO and O₃ profile comparisons	55
5.1 Summary	55
5.2 IAGOS Ozone	57
5.3 IAGOS Carbon Monoxide	65
6 Validation of regional model tropospheric NO₂ using MAX-DOAS	70
6.1 Summary	70
6.2 Introduction	70
6.3 Inter-comparison method	71
6.4 Results	71
7 Validation of tropospheric NO₂ columns against satellite retrievals	78
7.1 Summary	78
7.2 Comparison with GOME-2 NO ₂	78



8	Comparison with high-altitude EEA Air Quality e-reporting surface stations	81
8.1	Summary	81
8.2	Introduction	81
8.3	Regional ensemble results	83
8.4	Results for the seven regional models	86
9	Comparison with ozone sonde observations	87
9.1	Summary	87
9.2	Comparison approach	87
9.3	Results for the ENSEMBLE	87
9.4	Results for individual regional models	90
10	Comparison with GAW stations	92
10.1	Summary	92
10.2	Comparison method	92
10.3	Ozone	94
10.4	Carbon monoxide	97
11	Comparisons with MOPITT CO	99
11.1	Summary	99
11.2	Method	99
12	Summary of findings for individual models	104
13	Acknowledgements	105
14	References	106



1 Introduction

The Copernicus Atmosphere Monitoring Service (CAMS, <https://atmosphere.copernicus.eu>) is a component of the European Earth Observation programme Copernicus. The CAMS near-real time services consist of daily analysis and forecasts with the IFS system with data assimilation of trace gas concentrations and aerosol properties. The global modelling system is also used to provide the boundary conditions for an ensemble of more detailed regional air quality models that are used to zoom in on the European domain and produce 4-day forecasts of air quality. The regional forecasting service provides daily 4-days forecasts of the main air quality species and analyses of the day before, based on the results from 7 state-of-the-art atmospheric chemistry models. The ensemble represents the median of the 7 model forecasts.

Routine validation of the regional models against surface observations from the European member states (EEA Air Quality e-reporting) is provided for each model individually as well as the ensemble in separate quarterly validation reports. Validation reports of the CAMS regional products are available in the following portal:

<https://atmosphere.copernicus.eu/regional-air-quality-production-systems>.

This web page provides access to the quarterly reports on the daily analyses and forecast activities and verification of the regional ensemble. An overview of the regional air quality forecasting system is provided by Marécal et al (2015).

Validation reports (e.g. Wagner et al., 2020) for the CAMS global products are available at <https://atmosphere.copernicus.eu/node/325>, including the evaluation on Earth's troposphere, stratosphere, aerosols and greenhouse gases, with state-of-the-art observational datasets (GAW, IAGOS, MOPPIT, EMEP, GOME-2, OMPS-LP, BASCOE, AERONET etc.). A published overview on the validation of reactive gases and aerosols in the global analysis and forecast system can be found in Eskes et al (2015). A validation study of the global surface ozone reanalysis for Europe is provided by Katragkou et al (2015).

Details of the various observational datasets can be found in Eskes et al. (2019), "Observations characterisation and validation methods document", also available at:

https://atmosphere.copernicus.eu/sites/default/files/2020-02/CAMS84_2018SC1_D6.1.1-2019_observations_v4_0.pdf.

This document presents an evaluation of the concentrations above the surface as modelled by the set of 7 (9) regional models and the ensemble forecasts derived from these individual forecasts and analyses, and the consistency between the global and regional modelling systems of CAMS.



2 Regional and global CAMS forecasting systems

2.1 Regional models

The European Air Quality products are provided from the Copernicus Atmosphere Monitoring Service (<http://atmosphere.copernicus.eu/>). These data are available in NetCDF or Grib-Edition2 format. The files are available each day through ftp protocol from the Météo-France server (<ftp.cnrm-game-meteo.fr>). Since the beginning of February 2020, the data are also available through the Atmosphere Data store (<https://ads.atmosphere.copernicus.eu>). The products are available in Near Real Time (NRT) for four forecast days, following the protocol below:

- Each day 96h model forecasts and 24h analyses for the previous day are provided with hourly resolution. Consistent provision of the analysis product started on the 5th of July 2017.
- Products are available at eight vertical height levels: surface, 50, 250, 500, 1000, 2000, 3000, 5000 meters.
- The pollutants are O₃, CO, NO₂, SO₂, PM_{2.5}, PM₁₀, NO, NH₃, NMVOC, PANs, dust aerosols (fraction below 10µm), secondary inorganic aerosols (fraction below 2.5µm), birch, grass, olive, and ragweed pollen. Since the upgrade of February 4th 2020, three new aerosol species (PM from wildfires, EC from fossil fuels and EC from wood burning) are also available.
- The regional datasets cover the longitudes 335.05°E to 44.95°E every 0.1°, and latitudes 69.95°N – 30.05°N also at 0.1° resolution (~10km). Since June 12th, 2019, the northern boundary of the domain extends to 71.95°N.
- The forecasts until the 48th hour are available before 7:30 UTC
- The forecasts 49-96th hour are available before 9:30 UTC
- The analyses are provided before 12:00 UTC
- Since the June 2019 upgrade, the regional models make use of the CAMS-REG-AP_v2_2_1 emissions (reference year: 2015) and since February 4th, the updated CAMS-REG-AP_v3_1 emissions dataset (reference year: 2016).

The NRT forecast and analysis regional air quality data are available for the seven air quality models and their ensemble median (CAMS-regional or ENSEMBLE):

- MOCAGE model (MFM)
- LOTOS-EUROS model (KNM)
- EMEP MSC-W model (EMP)
- MATCH model (SMH)
- EURAD-IM model (RIU)
- CHIMERE model (CHI)
- SILAM model (FMI)

After October 16th, 2019 the regional ensemble is calculated based on two additional new regional models, namely DEHM and GEM-AQ. Separate data for those models were, however, not available to us at the time of writing of this report, using the dissemination channels mentioned above.



Every evening, a full download of the 96h forecasts and 24h analyses fields to KNMI at full resolution is performed. These fields are co-located to the set of surface stations used, and this largely reduced datasets is shared with all validation partners.

Documentation about the regional models may be found at the address <https://atmosphere.copernicus.eu/regional-air-quality-production-systems>. For the purposes of this report however, it's useful to indicate what kinds of observations are actively assimilated in each model (Table 2.1).

Table 2.1: Surface and other observations that are actively assimilated in regional models.

Model	Method	Surface	Other
CHIMERE	Kriging	O ₃ , NO ₂ , PM10, PM2.5	
EMEP	Intermittent 3D-var	O ₃ , SO ₂ , NO ₂ , PM10*, PM2.5*	OMI NO ₂
EURAD	Intermittent 3D-var	O ₃ , SO ₂ , NO ₂ , PM10, PM2.5	OMI and MetOp/GOME-2 NO ₂ and SO ₂ , MOPITT and IASI CO
LOTOS-EUROS	Ensemble Kalman filter	O ₃ , NO ₂ , PM10, PM2.5	OMI NO ₂
MATCH	Intermittent 3D-var	O ₃ , NO ₂ , CO, PM10, PM2.5	
MOCAGE	3D-var	O ₃ , NO ₂ , PM10	
SILAM	Intermittent 3D-var	O ₃ , NO ₂ , CO, PM10, PM2.5	













* PM2.5 and PM10 were not assimilated by EMEP in December and January of 2020

Validation reports of the CAMS regional products are available in the following portal: <https://atmosphere.copernicus.eu/regional-services>.

Whenever possible, in this report, models follow the naming and colour scheme of Table 2.2.



Table 2.2: Naming and colour scheme followed throughout this report. Since October 2019 the ensemble is computed using also the two new models, DEHM and GEM-AQ. These additional models have already been included in the table below.

<i>Model</i>	<i>Short model name</i>	<i>Colour name</i>	<i>Colour</i>
CAMS-global	CAMSG	Blue	
ENSEMBLE forecast	ENS-fc	Red	
ENSEMBLE analysis	ENS-an	Green	
CHIMERE	CHIM	Yellow	
EMEP	EMEP	Brown	
EURAD	EURAD	Cyan	
LOTOS-EUROS	LOTOS	Purple	
MATCH	MATCH	Grey	
MOCAGE	MOCA	Pink	
SILAM	SILAM	Orange	
DEHM	DEHM	Fuchsia	
GEM-AQ	GEMAQ	Light Green	



2.2 Global CAMS system based on the ECMWF IFS model

The CAMS-global operational assimilation/forecast system consists of the IFS-CB05 chemistry combined with the MACC aerosol model. The chemistry is described in Flemming et al. (2015); aerosol is described by the bulk aerosol scheme (Morcrette et al., 2009). Dissemination of CAMS-global forecasts is twice a day, at about 10:00 and 22:00UTC. The forecast length is 120 h. Users can get access at <https://atmosphere.copernicus.eu/catalogue>. Table 2.3 provides information on the satellite data used in CAMS-global.

Table 2.3: Satellite retrievals of reactive gases and aerosol optical depth that are actively assimilated in CAMS-global.

Instrument	Satellite	Provider	Version	Type	Status
MLS	AURA	NASA	V4	O ₃ Profiles	20130107 -
OMI	AURA	NASA	V883	O ₃ Total column	20090901 -
GOME-2A	Metop-A	Eumetsat	GDP 4.8	O ₃ Total column	20131007 - 20181231
GOME-2B	Metop-B	Eumetsat	GDP 4.8	O ₃ Total column	20140512 -
SBUV-2	NOAA-19	NOAA	V8	O ₃ 21 layer profiles	20121007 -
OMPS	Suomi-NPP	NOAA / EUMETSAT		O ₃ Profiles	20170124 - 20190409
IASI	MetOp-A	LATMOS/ULB EUMETSAT	-	CO Total column	20090901 - 20180621 20180622 - 20191118
IASI	MetOp-B	LATMOS/ULB EUMETSAT	-	CO Total column	20140918 - 20180621 20180622 -
IASI	MetOp-C	EUMETSAT		CO Total column	20191119-
MOPITT	TERRA	NCAR	V5-TIR V7-TIR V7-TIR Lance V8-TIR	CO Total column	20130129 - 20160124 - 20180626 20180626 20190702
OMI	AURA	KNMI	DOMINO V2.0	NO ₂ Tropospheric column	20120705 -
OMI	AURA	NASA	v003	SO ₂ Tropospheric column	20120705 - 20150901
GOME-2A/2B	METOP A/B	Eumetsat	GDP 4.8	SO ₂ Tropospheric column	20150902-
MODIS	AQUA / TERRA	NASA	Col. 6 Deep Blue	Aerosol total optical depth, fire radiative power	20090901 - 20150902 -
PMAp	METOP-A METOP-B	EUMETSAT		AOD	20170124 - 20170926 -
GOME-2A/2B	METOP A/B	Eumetsat	GDP 4.8	NO ₂ Tropospheric column	20180624-



The most recent upgrade to cycle 46R1, relevant to this report, took place on 9 July 2019 and involves a change from 60 to 137 vertical levels. A detailed changelog and the corresponding validation reports for this last upgrade can be found on the following page:

<https://atmosphere.copernicus.eu/node/472>.

Upgrade and version information is available here:

<https://atmosphere.copernicus.eu/changes-cams-global-production-system>.

Documentation on the global system can be found here:

<https://atmosphere.copernicus.eu/global-production-system>.



3 Consistency between the global and regional modelling components of CAMS

3.1 Summary

This chapter reports on the consistency between the global and regional modelling components of CAMS, and the impact of global CAMS boundary conditions on regional forecasts. The current evaluation includes ozone (O₃) carbon monoxide (CO) and aerosol (PM₁₀/PM_{2.5}) forecasts covering the period from December 2019 to February 2020.

Global and regional ensemble forecasts: The two forecast products compare well, surface O₃ appears lower and surface CO higher in CAMS-global compared to the ENSEMBLE.

Regional forecast variability: The regional ENSEMBLE members exhibit the expected regional variability for O₃, CO and particulate matter. Minor deviations (compared to CAMS-global and the majority of regional models) are generally noted. Lower CO concentrations are seen for LOTOS-EUROS at the level of 250 m. MOCAGE has distinct western, southern and northern PM₁₀ boundaries.

Diurnal cycles: Comparison of the diurnal cycles between the global and the regional forecasts over different European subregions indicate a very good agreement for surface O₃ and more pronounced diurnal variability for the surface CO global forecast. The agreement in the PM₁₀ is not as good in the south as with PM_{2.5}.

Daily time series. There is a very good temporal comparison between the global and the regional ensemble forecast for surface O₃. A good agreement is seen for surface CO. PM_{2.5} compare better than PM₁₀. LOTOS-EUROS seems to overestimate some PM episodes.

Regional forecast and analysis: The comparison of regional analyses and forecasts shows strong model and species dependence.

3.2 Methodology for the comparisons of CAMS-global and CAMS-regional

Operational download

The daily regional CAMS forecasts are retrieved on a daily basis. This includes the 3-hourly (0,3,6,9 etc.) regional forecast data (ensemble members and regional ensemble) for all provided species at all vertical layers for the 5 forecast days extracted from the Météo-France ftp server, and the 3-hourly (0, 3, 6, 9 etc.) global forecast data for 5 forecast days extracted from the ECMWF CAMS ftp server.



Methodology of global-regional comparison

The following methodology is used to a) convert CAMS-global species from mass mixing ratio (kg/kg) to concentration ($\mu\text{g}/\text{m}^3$) and, b) extract CAMS-global species concentrations from the vertical levels that lie closest to the regional height levels.

The following parameters are used from the CAMS-global model: hybrid layer coefficients; temperature, surface pressure; "GEMS" ozone; carbon monoxide; "aermr01" to "aermr11" (aerosol species, kg/kg). Data from the first 28 vertical layers (from the surface) are used.

The thickness of each vertical layer Δz (m) is calculated:

$$\Delta z_k = \frac{R * T_k}{M_{\text{air}} * g} * \ln\left(\frac{p_{i_{k+1}}}{p_{i_k}}\right) \quad (\text{E. 1})$$

where $R=8.314 \text{ J/mol}\cdot\text{K}$ the gas constant, T the temperature at vertical layer midpoint, $M_{\text{air}}=28.97\cdot 10^{-3} \text{ kg/mol}$ the molecular weight of air and $g=9.8 \text{ m/s}^2$ the gravity acceleration.

The mass-mixing ratio (kg/kg) for ozone (go_3) and carbon monoxide (CO) is initially provided. Conversion from mass mixing ratio (kg/kg) to concentration ($\mu\text{g}/\text{m}^3$) is performed using the following approach:

$$\rho_{\text{O}_3} = \text{mmr}_{\text{O}_3} * \left(\frac{p_m * M_{\text{air}}}{R * T}\right) \quad (\text{E. 2})$$

where ρ_{O_3} is the ozone concentration (kg/m^3) and mmr_{O_3} the ozone mass mixing ratio (kg/kg). The expression inside the parentheses in E.2 corresponds to the air concentration (kg/m^3). The same approach is also used for the CO unit conversion from kg/kg to $\mu\text{g}/\text{m}^3$.

The mass mixing ratio for all aerosol species (see Table 3.1) is initially provided. The PM10 and PM2P5 species are converted to $\mu\text{g}/\text{m}^3$ as follows:

$$\rho_{\text{PM}_{10}} = \left[\frac{\text{aermr01}}{4.3} + \frac{\text{aermr02}}{4.3} + \text{aermr04} + \text{aermr05} + 0.4 * \text{aermr06} + \text{aermr07} + \text{aermr08} + \text{aermr11} + \text{aermr09} + \text{aermr10} + \text{aermr16} + \text{aermr17} + \text{aermr18} \right] * \left(\frac{p_m}{R_{\text{spec}} * T}\right) \quad (\text{E. 3})$$

$$\rho_{\text{PM}_{2.5}} = \left[\frac{\text{aermr01}}{4.3} + \frac{\text{aermr02} * 0.5}{4.3} + \text{aermr04} + \text{aermr05} + 0.7 * \text{aermr07} + 0.7 * \text{aermr08} + 0.7 * \text{aermr11} + \text{aermr09} + \text{aermr10} + 0.7 * \text{aermr16} + 0.25 * \text{aermr17} + 0.7 * \text{aermr18} \right] * \left(\frac{p_m}{R_{\text{spec}} * T}\right) \quad (\text{E. 4})$$

where $R_{\text{spec}}=287.058 \text{ J}/(\text{kg}\cdot\text{K})$ is the specific gas constant for dry air. The expression inside the parentheses in E.3 and E.4 corresponds to the dry air concentration (kg/m^3).




Figure 3.1. Left: Mean global and regional ensemble forecast ozone fields for four different vertical layers (0, 250, 2000, 5000 m) for DJF2020. Right: Cross sections for the same period of the global and regional ensemble ozone boundaries (south, west, north, east).

Regional model products are provided at the height levels of 0, 50, 250, 500, 1000, 2000, 3000 and 5000m. For every grid point and time step of the CAMS-global model, the differences between the height of each vertical layer midpoint z_m and the regional model height (e.g. 5000m) is calculated. The layer midpoint that exhibits the minimum height difference is the one that lies closest to the regional height level and is therefore selected for extraction of both chemical and aerosol species concentrations. The above procedure is performed for every regional height level. The final global product contains the O₃, CO, PM10 and PM2.5 concentrations in eight height levels that correspond to the CAMS-global vertical levels that lie closest to the regional height levels.




Figure 3.2. Left: Mean global and regional ensemble forecast CO fields for four different vertical layers (0, 250, 2000, 5000 m) for DJF2020. Right: Cross sections for the same period of the global and regional ensemble ozone boundaries (south, west, north, east).

Table 3.1: Aerosol species description.

Label		Name	Size (µm)
aermr01	SS1	Sea Salt Aerosol	0.03-0.5
aermr02	SS2	Sea Salt Aerosol	0.5-5
aermr03	SS3	Sea Salt Aerosol	5-20
aermr04	DD1	Dust Aerosol	0.03-0.55
aermr05	DD2	Dust Aerosol	0.55-0.9
aermr06	DD3	Dust Aerosol	0.9-20
aermr07	OM1	Hydrophobic Organic Matter Aerosol	
aermr08	OM2	Hydrophilic Organic Matter Aerosol	
aermr09	BC1	Hydrophobic Black Carbon Aerosol	
aermr10	BC2	Hydrophilic Black Carbon Aerosol	
aermr11	SU1	Sulphate Aerosol	
aermr16	NI1	Nitrate fine mode aerosol	
aermr17	NI2	Nitrate fine mode aerosol	
aermr18	NH3	Ammonium aerosol	




Figure 3.3. Left: Mean global and regional ensemble forecast PM10 fields for four different vertical layers (0, 250, 2000, 5000 m) for DJF2020. Right: Cross sections for the same period of the global and regional ensemble ozone boundaries (south, west, north, east).




Figure 3.4 Left: Mean global and regional ensemble forecast PM2.5 fields for four different vertical layers (0, 250, 2000, 5000 m) for DJF2020. Right: Cross sections for the same period of the global and regional ensemble CAMS ozone boundaries (south, west, north, east).

3.3 Consistency between the global and regional forecasts

This section reports on the consistency of the global and regional ensemble forecast. The analysis is performed for O₃, CO, PM10 and PM2.5 at four levels (0, 250, 2000 and 5000 m) for the time period from December 2019 to February 2020 (DJF).

Ozone (O₃)

Figure 3.1 shows the average spatial distribution of O₃ for different vertical layers (left) and the cross sections of the lateral boundaries (right) for the CAMS-global forecast and the regional ENSEMBLE. CAMS-global exhibits lower ozone near the surface and in the boundary layer over continental Europe.

Carbon monoxide (CO)

Figure 3.2 illustrates the seasonal mean fields of CO and the cross sections of lateral boundaries (right) for ENSEMBLE and CAMS-global. CAMS-global and the ENSEMBLE exhibit some regional differences in CO, surface concentrations are somewhat higher in CAMS-global, but occasionally lower in higher altitudes (e.g. Fig3.2 northern boundary).

Aerosols (PM10 and PM2.5)



Figures 3.3 and 3.4 illustrate the CAMS-global and ENSEMBLE spatial distributions (left) and lateral boundary cross sections (right) of PM10 and PM2.5 mean fields, respectively. The main inconsistencies for particulate matter are summarized below:

- Differences between surface and PBL PM10 are seen over the southern boundary, indicating different dust aerosol loadings. The global forecast system has higher and spatially more extended dust concentrations than the regional ensemble below < 2 Km.

3.4 Regional variability

Ozone (O₃)

Figure 3.5 illustrates ozone mean fields for DJF2020 of the individual regional ensemble members and CAMS-global (bottom panel) for selected altitudes (0, 250, 2000, 5000 m). No particular inconsistencies identified, there is an expected variability between the regional ensemble members. A boundary issue is obvious in at the 5 Km in the EURAD model, obviously the regional model domain is smaller than the standard CAMS domain.

Carbon monoxide (CO)

The DJF2020 mean fields of carbon monoxide for the regional ensemble members as well as for CAMS-global are illustrated in Figure 3.6 for different vertical layers. There are generally no discrepancies, we mostly see the expected regional variability. LOTOS-EUROS has generally lower CO concentrations, especially seen at the 250 m level.

Aerosols (PM10 and PM2.5)

The mean PM10 and PM2.5 fields are illustrated in Figures 3.7 and 3.8, respectively. The greatest difference among regional models and with the CAMS-global are seen over the southern boundary. The model with the most distinct behavior is MOCAGE with different western, southern northern PM boundaries compared to other regional ensemble members and the CAMS-global. The agreement is better for the PM2.5 forecast products.




Figure 3.5 Mean regional ozone forecasts for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global).




Figure 3.6. Mean regional CO forecasts for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global).




Figure 3.7. Mean regional PM10 forecasts for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global).




Figure 3.8. Mean regional PM_{2.5} forecasts for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global).



3.5 Time series

Figure 3.9 shows the mean daily time series from December 2019 to February 2020 for the four species, namely O₃, CO, PM₁₀, PM_{2.5} (from left to right) for different European sub-regions (from top to bottom): Alps (AL), British Isles (BI), East Europe (EA), France (FR), Iberian Peninsula (IP), Mediterranean (MD), Mid-Europe (ME), Scandinavia (SC).

Each subregion is defined with the following latitude/longitude boundaries:

Name = (BI, IP, FR, ME, SC, AL, MD, EA)

West = (-10, -10, -5, 2, 5, 5, 3, 16)

East = (2, 3, 5, 16, 30, 15, 25, 30)

South = (50, 36, 44, 48, 55, 44, 36, 44)

North = (59, 44, 50, 55, 70, 48, 44, 55)

The letter R denotes the temporal correlation between the two products. Only concentrations over land are used.

For O₃ the temporal correlation ranges are higher than 0.9 over all regions. The agreement is surface O₃ magnitude are very good. The EURAD model exhibits only toward the end of the DJF period lower surface O₃, especially over SC.

High temporal correlations are found for surface CO. The higher surface CO concentrations are evident in specific areas (AL, MD, and also some forecasted spikes in FR).

The PM correlations are generally lower but, overall, they indicate a relatively good agreement between the regional and global forecasts. LOTOS-EUROS seems to overestimate some PM events over southern Europe for PM₁₀. The agreement with PM_{2.5} is better.




Figure 3.9. Mean daily time series for surface O₃, CO, PM₁₀ and PM_{2.5} for DJF2020 (unit $\mu\text{g}/\text{m}^3$). The blue line is CAMS-global and the red line the ENSEMBLE forecasts. Each line in the composite plot denotes a different European subregion.



3.6 Diurnal cycles

Figure 3.10 shows the diurnal cycles for surface O₃, CO, PM₁₀ and PM_{2.5} averaged over the period DJF2020 for different European sub-regions (from top to bottom): Alps (AL), British Isles (BI), East Europe (EA), France (FR), Iberian Peninsula (IP), Mediterranean (MD), Mid-Europe (ME), Scandinavia (SC). The red colour is used for the regional ENSEMBLE and the blue for CAMS-global.

There is a good agreement between the O₃ CAMS-global and the regional diurnal cycles.

The diurnal range for surface CO is less pronounced for the regional products over some regions, mostly over the Alps and over the Mediterranean.

The agreement in the PM₁₀ diurnal cycle is not a good one in the south (IP, MD) and France (out of phase). The agreement for PM_{2.5} is generally better.




Figure 3.10. Diurnal cycles for surface O_3 , CO, PM_{10} and $PM_{2.5}$ for the period DJF2020 (unit $\mu\text{g}/\text{m}^3$). The blue line is CAMS-global and the red line the ENSEMBLE forecast. Each line in the composite plot denotes a different European sub-region.



3.7 Regional domain boundary cross sections

Ozone (O₃)

Figure 3.11 shows the regional variability in the lateral domain boundary cross sections of O₃ (from left to right): south, west, north, east and the different ensemble members and CAMS-global (from top to bottom) averaged over the period DJF2020. A clear inconsistency is seen for CHIMERE. LOTO-EUROS also does not show the expected boundary variability.

Carbon monoxide (CO)

Figure 3.12 shows the regional variability in the lateral cross sections of CO (from left to right): south, west, north, east and the different ensemble members and CAMS-global (from top to bottom). We identify only in CHIMERE an inconsistency in boundaries.

Aerosols (PM₁₀ and PM_{2.5})

Figure 3.13/3.14 shows the regional variability in the lateral cross sections of PM₁₀/PM_{2.5} respectively (left to right): south, west, north, east and the different ensemble members and CAMS-global (top to bottom) averaged over the period DJF2020. The boundaries exhibit an expected higher regional variability especially over the southern boundary. MOCAGE is the model with the greatest differences compared to CAMS-global. The agreement is better for PM_{2.5}.




Figure 3.11. Ozone cross sections for DJF2020 for the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global) and the lateral boundaries (left to right: south, west, north, east).




Figure 3.12. Carbon Monoxide cross sections for DJF2020 for the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global) and the lateral boundaries (left to right: south, west, north, east).




Figure 3.13. Aerosol PM10 cross sections for DJF2020 for the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global) and the lateral boundaries (left to right: south, west, north, east).




Figure 3.14. Aerosol PM_{2.5} cross sections for DJF2020 for the seven ensemble members and CAMS-global (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and CAMS-global) and the lateral boundaries (left to right: south, west, north, east).



3.8 Regional analysis vs. regional forecasts

In the following sections we compare the regional analysis products with the regional forecasts (Day1). The four following figures (3.15-3.18) show the mean regional differences between analysis and forecasts for the time period DJF2020 at four different vertical layers (0, 250, 2000, 5000 m, left to right) including O₃, CO, PM10 and PM2.5, respectively.

Regional models with the largest differences between the analyses and the 1st day forecasts are for:

Ozone

- CHIMERE in the surface layer.
- EMEP mostly within the PBL
- LOTOS-EUROS, MATCH and MOCAGE over the total vertical extent.

Carbon monoxide

- EMEP up to 250 m
- SILAM up to 2km altitude.
- MATCH over the total vertical extent.

PM10 and PM2.5

- Most pronounced differences in LOTOS-EUROS and MATCH over the Mediterranean, EMEP over continental Europe.

It is important to note that the differences observed between the analysis and the forecast do not only reflect the impact of the assimilation but may also result from differences in the model setup between the analysis systems and the forecast systems.




Figure 3.15. Mean regional O₃ differences between analysis and forecast for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from regional ensemble and individual ensemble members (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and ENSEMBLE).




Figure 3.16. Mean regional CO differences between analysis and forecast for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from regional ensemble and individual ensemble members (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and ENSEMBLE).




Figure 3.17. Mean regional PM10 differences between analysis and forecast for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from regional ensemble and individual ensemble members (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and ENSEMBLE).




Figure 3.18. Mean regional PM_{2.5} differences between analysis and forecast for DJF2020 for four different vertical layers (0, 250, 2000, 5000 m) from regional ensemble and individual ensemble members (top to bottom: CHIMERE, EMEP, SILAM, LOTOS-EUROS, MOCAGE, EURAD-IM, MATCH and ENSEMBLE).



4 Vertical profile and column aerosol comparisons

4.1 Summary for the EARLINET lidar and Aeronet comparisons

The regional models are compared with climatological lidar profiles for each season (EARLINET/ACTRIS data from 2006-2018). Missing information on composition, size and humidity growth of the aerosol in the models introduces considerable uncertainty to the PM derived extinction, which conservatively spans up to a factor 10 for absolute extinction values. Aeronet data are used to calibrate the conversion from modelled mass to optical property aerosol extinction. This way the order of magnitude in extinction is similar between the models and the lidar profiles, but also significant differences appear at some stations in the lowest layers (Granada, Athens). Relative differences in the form of extinction profiles are more certain. We choose the most representative five stations to compare in retrospective the seasonal average aerosol profiles since 2016. The retrospective of the seasonal comparisons since 2016 shows very similar profiles during this season. The respective overestimation or underestimation of the extinction found in 2016 are usually also found in following years with the ENSEMBLE.

Introduction

The vertical distribution of aerosol reflects processes like atmospheric mixing, removal, and aerosol transport from outside of the domain or formation of secondary aerosol. The vertical mixing processes determine ground concentrations in polluted areas. Long-range transported aerosol, often carried aloft, may contribute to pollution in clean regions. Evaluation of the simulated aerosol column and vertical profiles are thus valuable for the performance characterisation of air quality models.

The 7 regional models provide mass concentration vertical profiles (PM_{2.5} and PM₁₀) over Europe and may thus be evaluated for their aerosol vertical distribution. However, only very few aircraft campaigns and mountain sites are available to validate aerosol mass at altitude. In contrast frequent measurements of vertical profiles of aerosol backscatter, extinction or its integral, aerosol optical depth, exist. Deriving aerosol optical properties from the mass concentrations is thus needed, assuming lidar ratios and mass extinction coefficients, at least until the models provide more specific output on aerosol composition and optical properties.

In order to assess mass extinction coefficients chosen, the aerosol optical depth derived from the model mass profiles is first compared to AERONET AOD measurements. Secondly, we document a comparison of the extinction profiles derived from modelled mass concentration with climatological extinction profiles derived from European EARLINET/ACTRIS lidars.




Figure 4.1. Comparison of mean profiles of two aerosol optical properties derived from the same Earlinet Raman lidar at the same time: a) aerosol extinction @532nm (light blue line) and b) extinction @532nm derived from the aerosol backscatter coefficient (dark blue) using a lidar ratio of 50 sr. The profiles use data taken at simultaneous times at each station; number of profiles given as n: x. Based on measurements in the period 2006-2014.

4.2 Methodology

AERONET data

The AERONET sun-photometers measure in non-cloudy conditions the aerosol optical depth at several wavelength and in near real time. The spatial distribution of the instruments allows a good coverage of aerosol observation over Europe. The version 3 level 1.5 data has been used in the reporting period presented here. This version and quality level ensures an efficient filtering of the residual clouds (mainly cirrus) for data in near real time. Daily AERONET aerosol optical depth, measured at 550 nm, has been averaged over summer for the European sites available in CAMS model output.

Lidar data

The EARLINET/ACTRIS Lidars are distributed over several locations in Europe and allow comparison across different climates in Europe (Pappalardo et al., 2014). Yet, up to now, there are no near-real-time data available. Regular measurements in EARLINET are sparse and often acquired once per week, with gaps due to maintenance or funding restrictions. A climatology has been computed per station and per season with all measurements available between 2006 and 2018.




Figure 4.2. Mass extinction coefficient estimated at AERONET location sites (blue dots) and EARLINET stations (triangles) for D2019-JF2020 (left). A European map of the Mass Extinction Coefficient has been constructed with cubic interpolation in the inner part of the region covered by AERONET on the model grid (right), and with the nearest neighbours in the outer part of this region (not shown).

The backscatter coefficient and extinction profiles at 532 nm have been extracted from the EARLINET database. The more frequently measured backscatter profiles are considered here with priority. An aerosol extinction coefficient profile is computed from the backscatter coefficient using a range of plausible lidar ratios. This latter parameter depends on the aerosol type and is more likely decreasing with the size of the aerosol. Minimum values are observed for sea salt aerosol (below 30 sr at 550 nm, [Ackermann et al., 1998, Omar et al., 2009]), while larger values are related to urban particles (55 sr in [Muller et al., 2007], 70 sr in [Cattrall et al., 2005]). Desert dust is associated with intermediate lidar ratios, ranging from 30 sr to 60 sr depending on the sources and the transport regime. A climatology of aerosols in West Africa published in Mortier et al. [2016] revealed an average lidar ratio (over 9 years) of about 30 ± 15 sr. Due to the location of the stations involved in this study, both dust and urban aerosols and any a mix of them might occur. In order to represent the uncertainty on the nature of aerosols, we show the range in likely mean extinction using a lidar ratio extending from 30 to 70 sr.

As a test, the conversion of lidar backscatter to extinction coefficient is performed assuming a constant average lidar ratio of 50 sr at locations where both backscatter and extinction coefficients are measured in EARLINET. This allows consistent comparison and visualization of some of the error associated to our simplified constant lidar ratio assumption. We have excluded from this comparison cases where local extinction coefficient was above 0.5 km^{-1} in order to avoid outliers. The profiles shown in figure 4.1 from 4 stations reveal an error in the mean profile of 0–30% in extinction, which is small compared to the model spread documented below. The extinction profiles derived from the backscatter coefficients look vertically smoother. We have excluded from this comparison cases where local extinction coefficient was above 0.5 km^{-1} in order to avoid outliers.



In addition to this aerosol typing uncertainty, a sampling error should be accounted for. The observations are sporadic, while the models predict the aerosol concentration continuously. Therefore, seasonal averages are not computed with the same coverage in model and observation. Our earlier model-based bootstrap studies revealed, that, depending on the station, a set of ca. 30 daily observations allows reproducing the seasonal average with an error of about 10% [ACTRIS Deliverable WP6/D6.21]. In our case, this error might be larger since the synoptic situation is very different between our EARLINET climatological dataset, covering 2006-2018, and the season covered in this report. An overall uncertainty of about 20% has been chosen to represent the sampling error.

Model data

The ENSEMBLE and the underlying 7 regional CAMS models are investigated. For each of these models, the hourly PM₁₀ and PM_{2.5} vertical profiles are extracted at the EARLINET station locations from the first day of each daily forecast at levels 0, 50, 250, 500, 1000, 2000, 3000, 5000 m.

The conversion of PM₁₀ mass concentration to extinction requires a mass extinction coefficient (MEC). MEC depends on the size distribution, refractive index and density of the particles. This information is not yet available from the models. For different kind of aerosols, MEC values can vary from about 0.5 m²g⁻¹ in the case of desert dust aerosols up to 8 m²g⁻¹ for urban particles [Chin et al., 2002]. No variation with height or aerosol type is taken into account, mainly because the models provide no further info on aerosol speciation. We derive the MEC value to convert the model profile data to extinction profiles from a combination of the modelled mass column load and consistent Aeronet AOD data.

For the Aeronet based computation of the MEC, the model data are picked at the location and on the day when sun photometer observations were available. The CAMS-regional mass concentrations have been averaged for coincident days (with the measurements) and averages are converted, with a seasonal and site dependent mass extinction coefficient estimated with AERONET retrievals, into extinction profiles. A seasonal and site dependent mass extinction coefficient is obtained when combining it with AERONET AOD retrievals. Values of MEC are ranging from 1 m²g⁻¹ in South-West of Europe to more than 10 m²g⁻¹ in the North-East. Since some of the EARLINET stations are not co-located with Sun photometers, a European map of MEC has been constructed, for each season, by interpolating (cubic interpolation) and extrapolating (nearest neighbour) the available AERONET based MEC calculations on the grid of the model (figure 4.2). One can notice a gradient with the longitude with lower values found in the Western part while the highest values are observed in the Eastern part of Europe. Also, the values are generally lower as compared to last year, which might reveal a higher concentration in coarse particles (dust). The seasonal AERONET-based MEC is then used at each EARLINET station to calculate the model extinction from concentration profiles. The uncertainty on the MEC being removed allows more accurate comparisons with the observed vertical profiles than using an average MEC over whole Europe.





Fig 4.3. Extinction profiles December 2019 - February 2020 derived from the ENSEMBLE forecast mass concentration profiles (red envelope) and from EARLINET (climatology) backscatter profiles (grey envelope: lidar ratio uncertainty, light grey: including sampling error). “n: XX means number of individual EARLINET profiles assembled (December 2018 - February 2006-2018). The EMC used for the calculation of the extinction from the concentration profiles is indicated for each station below the number of EARLINET profiles “n” used for the calculation of the climatology.

4.3 Results

4.3.1 Comparison of extinction profiles

The extinction profiles estimated from the 7 CAMS models and the ENSEMBLE and EARLINET measurements are compared for DJF2020 (figure 4.3).

One observes generally a good agreement between the two datasets at the exception of Ispra and Granada while Ispra is associated with the highest number of measurements. In Cabauw, Leipzig and Bucharest, the models are in perfect agreement with the observed climatology above 1 km of altitude. In Leipzig, where the observations are also available close to the ground, the






Fig 4.4: Seasonal extinction profiles derived from ENSEMBLE forecast mass concentration profiles for 2016 (ENS16), 2017 (ENS17), 2018 (ENS18), 2019 (ENS19) and EARLINET climatology. The parenthesis indicates for the CAMS profiles, the MEC used for the extinction estimation, and for the EARLINET, the number of profiles used for constructing the climatology.

ENSEMBLE underestimates the extinction, while one of the models (EMEP) shows a similar pattern than in the observation dataset. In the Southern stations, a good agreement is in Evora, and Potenza above 2 km of altitude. In Barcelona, models are overestimating the extinction coefficient above 1 km of altitude and underestimate the extinction below this level. In Potenza and Athens, the models underestimate the extinction below 2 km of altitude.



4.3.2 Seasonal variability

In order to investigate the performance of the model in reproducing the vertical profiles, it is interesting to observe the inter-annual variability for the different seasons. This will be of use for the development of a score providing an assessment of the models skills, and is also useful to investigate the models synoptic variability. The seasonal profiles have been reported since 2016 at 5 stations (Leipzig, Barcelona, Potenza, Evora and Bucharest) in Figure 4.4, using the new EARLINET climatology. In autumn, the models seem to have the same biases as compared to the observations in between the different years. Performances of the models in 2019 look similar to the performances observed in winter 2018. The biggest difference is found in Barcelona where the models overestimate the extinction coefficient above the altitude of 1 km while the agreement seemed better, as compared to observations, last year.



5 IAGOS aircraft CO and O₃ profile comparisons

5.1 Summary

Routine observations of ozone and CO over European airports are available from the IAGOS fleet. Take-off and landing profiles were sampled from the hourly model 3D forecasts along the flight tracks.

Ozone in-situ: Both the regional ENSEMBLE and CAMS-global overestimate ozone from the surface to the free troposphere with very similar behaviour in all layers. In the lowest layers, ozone is well represented by the models with slightly better performance from the regional model, while in the bias in free troposphere is larger for all models.

Carbon monoxide: In the lowest layers, CO is mostly underestimated by the regional ENSEMBLE and CAMS global which present similar behaviour. In the free troposphere the agreement is better and CAMS-global performs better than the regional ENSEMBLE which underestimates.

IAGOS Validation Method

Validation is possible at the European airports visited by the IAGOS fleet. For the European-based carriers, there are regular profiles at the home airports. There are two aircraft operated by Lufthansa, one operated by Air France. Thus, when the fleet is fully operational, there are daily profiles Frankfurt and Paris (CDG). IAGOS is also installed on two aircraft operated by the Asian-based carrier China Airlines. Aircraft fly regularly from Taipei to Amsterdam or Vienna and sometimes to Rome. Other airports may be visited depending on the operational schedules of the airlines. During this period, in Europe only the airport of Frankfurt has been visited as shown in Figure 5.1.

We download the daily latitude-longitude datasets for the 7 regional models and the ENSEMBLE for two species (carbon monoxide and ozone) on 8 vertical levels (surface, 50m, 250m, 500m, 1000m, 2000m, 3000m, 5000m). The aircraft takes about 10 minutes to climb or descend the 5000m vertical extent covered by the regional models. During this time and travelling at up to 166 m s^{-1} , it covers about 120km and therefore traverses many grid-boxes of resolution 10km. We perform a spatial interpolation from the grid of the regional models to the aircraft's trajectory. The IAGOS measurements in ppbv are converted to $\mu\text{g m}^{-3}$ using the temperatures measured by IAGOS. The data are validated by the PI but are not yet calibrated. Calibration takes place after an operational period of about 6 months.




Figure 5.1. Map showing the number of profiles available in the period December 2019 - February 2020.




Figure 5.2: Daily time series of ozone at Frankfurt for the period DJF2020. The regional ENSEMBLE and associated analysis are shown in red and green respectively, and CAMS-global is shown in blue. The black dotted line is the monthly mean of the observations (IAGOS/MOZAIC) over the period 2003-2016 (Level 2 data), the black dashed line shows 1 standard deviation from the monthly mean and the black dot-dashed line shows 3 standard deviations from the monthly mean.




Figure 5.3: Mean profile of ozone at Frankfurt for the period DJF2020. The regional ENSEMBLE and associated analysis are shown in red and green respectively, and the global o-suite is shown in blue. The shaded area indicates the range of the mean climatology of the observations plus/minus one standard deviation during the same period for all years between 2003 and 2016 (level 2 data).

5.2 IAGOS Ozone

For the period December 2019 - February 2020, observations are available only at Frankfurt, the daily time series are presented in Figure 5.2 and associated averaged profiles over the whole period are presented in Fig. 5.3. In all plots, the forecast and analysis of the regional ENSEMBLE is shown in red and green respectively, and CAMS-global is shown in blue. Fig. 5.2 and 5.3 show that the regional and global models differ mostly at around 1 km. In the lowest layers, the ENSEMBLE performs better than CAMS which present larger overestimations. In the upper layers, the regional and global models behave more similarly.

During DJF2020 no major episode is observed in the low troposphere at Frankfurt, however ozone values in the surface are close to three standard deviation in two cases, and more often ozone values close to or slightly greater than one standard deviation from climatology are found (Fig. 5.2). On the time series, in the surface layer a number of clear peaks are found around 9 December, 15 December, 28 January, 11 February and 26 February, which also correspond to high boundary layer values often close to one standard deviation. The largest boundary layer values are found around 22 January, reaching nearly three standard deviations while surface values are below one standard deviation for the corresponding dates.




Figure 5.4.a. Selection of ozone profiles at Frankfurt during the period DJF2020. The ENSEMBLE is shown at 4 forecast times (blue: 1-day; cyan: 2-day; green: 3-day; red: 4-day). (Units: $\mu\text{g m}^{-3}$).




Figure 5.4.b: Selection of ozone profiles at Frankfurt during the period DJF 2020. The ENSEMBLE is shown at 4 forecast times (blue: 1-day; cyan: 2-day; green: 3-day; red: 4-day). (Units: $\mu\text{g m}^{-3}$).




Figure 5.4.c: Selection of ozone profiles at Frankfurt during the period DJF 2020. The ENSEMBLE is shown at 4 forecast times (blue: 1-day; cyan: 2-day; green: 3-day; red: 4-day). (Units: $\mu\text{g m}^{-3}$).




Figure 5.4.d: Selection of ozone profiles at Frankfurt during the period DJF 2020. The ENSEMBLE is shown at 4 forecast times (blue: 1-day; cyan: 2-day; green: 3-day; red: 4-day). (Units: $\mu\text{g m}^{-3}$).

The individual profiles related to the aforementioned cases are presented in Fig. 5.4.a-d. For most of these profiles, ozone concentrations in the surface layer are around $50 \mu\text{g}/\text{m}^3$, with maximum values between 70 and $80 \mu\text{g}/\text{m}^3$ reached on 28 January and 11 February. These profiles present a nearly constant shape from the boundary layer to the free troposphere with values for most cases in the range of 70 - $80 \mu\text{g}/\text{m}^3$. Here, the regional ENSEMBLE performs generally well with the exception of some cases (1207 14:10, 1208 11:48) where overestimations are found. The results for the different forecast times are similar.




Figure 5.5.a: Ozone profiles at Frankfurt during the period DJF 2020. Each colour/Line style corresponds to one of the 7 models for the ensemble. In the legend, the models are CHIM=CHIMERE, EMEP=EMEP, SILAM=SILAM, LOTOS=LOTOS-EUROS, MOCA=MOCAGE, EURAD=EURAD, MATCH=MATCH. (Units: $\mu\text{g m}^{-3}$).




Figure 5.5.b: Ozone profiles at Frankfurt during the period DJF 2020. Each colour/Line style corresponds to one of the 7 models for the ensemble. In the legend, the models are CHIM=CHIMERE, EMEP=EMEP, SILAM=SILAM, LOTOS=LOTOS-EUROS, MOCA=MOCAGE, EURAD=EURAD, MATCH=MATCH. (Units: $\mu\text{g}/\text{m}^3$).




Figure 5.5.c: Ozone profiles at Frankfurt during the period DJF 2020. Each colour/Line style corresponds to one of the 7 models for the ensemble. In the legend, the models are CHIM=CHIMERE, EMEP=EMEP, SILAM=SILAM, LOTOS=LOTOS-EUROS, MOCA=MOCAGE, EURAD=EURAD, MATCH=MATCH. (Units: $\mu\text{g}\cdot\text{m}^{-3}$).




Figure 5.6: Daily time series of CO at Frankfurt for the period December 2019 - February 2020. The regional ENSEMBLE forecast and associated analysis is shown in red and green respectively, and CAMS-global is shown in blue. The black dotted line is the monthly mean of the observations (IAGOS/MOZAIC) over the period 2003-2016 (Level 2 data), the black dashed line shows 1 standard deviation from the monthly mean and the black dotted-dashed line shows 3 standard deviations from the monthly mean. (Units: $\mu\text{g}\cdot\text{m}^{-3}$).

Some profiles obtained with all individual models of the ensemble are presented in Fig. 5.5.a-c (chronological order). For most profiles of the period the results from all models are similar with the exception of the EURAD model which sometimes present much larger overestimations than the other models in the free troposphere (e.g. 1201 10:23, 1216 13:59, 1222 16:19, 1223 09:48, 0102 10:28, 0103 07:54, 0103 13:23, 0204 09:29, 0205 06:19, 0205 10:20, 0211 09). MATCH is also showing some larger bias than the other individual models in the free troposphere for two profiles on 0104 04:59 and 0216 14:03, with respectively overestimations and underestimations. Moreover, the MOCAGE model also presents a different behaviour than the other models on profiles of 0226 at 09:30 and 13:37, with large positive bias at all levels while the other models present good agreement.

5.3 IAGOS Carbon Monoxide

Like for the ozone section, the daily time series of CO and associated averaged profile at Frankfurt are presented in Fig. 5.6 and 5.7. The forecast and analysis of the regional ENSEMBLE is shown in red and green respectively, and CAMS-global is shown in blue. As it can be seen on the time series, for CO sampling is poorer than for ozone especially for the month of December. As shown on the time series, available observations of CO are always below one standard deviation from the climatology with a few exceptions near one standard deviation (Fig. 5.6). Moreover, it can be seen that the average profile for this period is much below the climatological profile for all levels (Fig. 5.7). This may be explained by a global impact of the decrease in China's pollution due to the coronavirus shutdown which took place during this period.




Figure 5.7: Profile of CO at Frankfurt for the period DJF2020. The regional ENSEMBLE and associated analysis are shown in red and green respectively, and the global o-suite is shown in blue. The shaded area indicates the range of the mean climatology of the observations plus/minus one standard deviation during the same period for all years between 2003 and 2016 (level 2 data).

In the surface and boundary layers CO is always underestimated by both regional and global models (Fig 5.6) with on average slightly better performance from the regional ENSEMBLE (Fig. 5.7). In the free troposphere, the agreement is better than in the lowest layers with smaller underestimations (Fig. 5.6 and 5.7), and from the time series CAMS-global is often performing better than the regional ENSEMBLE (Fig. 5.6).

Some of the individual profiles at Frankfurt of the ENSEMBLE are presented in Fig. 5.4. These profiles present the largest CO concentrations observed during DJF2020 in the lowest layers. On the corresponding dates, daily CO values are in general slightly above one standard deviation as shown on Fig. 5.6. Results from the regional ENSEMBLE in the surface and boundary layers present in general large underestimations of these profiles with no notable difference between the different forecast times (Fig. 5.4). The highest CO concentrations are found in profiles for 19 and 20 December, 27 January and 6 February with values above $500 \mu\text{g}\cdot\text{m}^3$ near the surface with a maximum of nearly $600 \mu\text{g}\cdot\text{m}^3$ on the profile of 19 December at 09:43.




Figure 5.8: Selection of CO profiles at Frankfurt during the period DJF 2020. The ENSEMBLE is shown at 4 forecast times (blue: 1-day; cyan: 2-day; green: 3-day; red: 4-day). (Units: $\mu\text{g}\cdot\text{m}^3$).




Figure 5.9.a: CO profiles at Frankfurt during the period DJF 2020. Each color/ line style corresponds to one of the 7 models for the ensemble. In the legend, the models are CHIM=CHIMERE, EMEP=EMEP, SILAM=SILAM, LOTOS=LOTOS-EUROS, MOCA=MOCAGE, EURAD=EURAD, MATCH=MATCH. (Units: $\mu\text{g}\cdot\text{m}^3$).




Figure 5.9.b: CO profiles at Frankfurt during the period DJF 2020. Each color/ line style corresponds to one of the 7 models for the ensemble. In the legend, the models are CHIM=CHIMERE, EMEP=EMEP, SILAM=SILAM, LOTOS=LOTOS-EUROS, MOCA=MOCA, EURAD=EURAD, MATCH=MATCH. (Units: $\mu\text{g}\cdot\text{m}^3$).

As regards the results of the individual models (Fig. 5.9.a-b), there are in general very similar for all profiles of DJF with sometimes, but rarely, some differences in the boundary layer (20200205, 20200206, 20200208). The profiles that are presented in Fig. 5.8 are also shown in Fig. 5.9 for individual models. As it can be seen, high CO concentrations in the lowest layers are not always well reproduced with large underestimations from all the individual models which behave similarly.



6 Validation of regional model tropospheric NO₂ using MAX-DOAS

6.1 Summary

MAX-DOAS surface remote sensing observations provide tropospheric columns of NO₂, with the largest sensitivity in the boundary layer. The magnitude of VCDs derived from the measurements for the three urban stations De Bilt, Uccle and Bremen are reproduced rather well by the models. Although many of the model simulated values probably fall within the uncertainty range of MAX-DOAS retrievals, the latter alone cannot explain differences between retrievals and simulations, especially those found for variations in time. Moderate correlations on the order of 30-70 % are found for each station. The regional ENSEMBLE performs significantly better than the global model in terms of correlation at each station. However, some of the larger NO₂ values inside individual pollution plumes are underestimated by the models.

6.2 Introduction

MAX-DOAS observations of atmospheric composition are performed by taking measurements of the scattered sunlight at different elevation (and sometimes also azimuthal) angles. Depending on the viewing angle and solar position, the light path through the atmosphere is different, with the observation in the zenith direction usually providing the shortest light path through the lower troposphere. Therefore, using the zenith measurement as intensity of incident radiation and the observations in other angles as intensity of transmitted radiation, the total amount of molecules of a certain species along the light path difference, the so-called slant column densities, can be determined using Lambert Beer's law. Using radiative transfer modelling and Optimal Estimation techniques, this can be inverted to tropospheric columns and even lower altitude tropospheric profiles.

The advantage of MAX-DOAS measurements is their ability to observe several pollution related species at the same time (e.g. NO₂, HCHO, CHOCHO, SO₂, aerosols, potentially also O₃) and to provide data which is virtually free of interferences from other species such as PAN or NO_y for NO₂. Also, the fact that the observations integrate over a comparatively large volume can be an advantage for satellite and model validation as the observed quantity is relatively close to the modelled one. On the other hand, the uncertainty of the retrievals is considerable (on the order of 30% for NO₂ tropospheric columns and larger for individual layers) and depends on cloud occurrence and aerosol loading.

In this report, regional air quality model forecasts of tropospheric NO₂ columns are compared to MAX-DOAS retrievals from 3 urban stations (De Bilt– KNMI, Bremen – IUP-UB, Uccle - BIRA-IASB). The reader is referred to previous reports for comparisons from the rural station OHP (BIRA-IASB) (which showed in general an underestimation by the model ensemble and an overall better performance for the o-suite here) as the instrument at this site stopped working in March 2017. Profile retrieval and averaging kernel data are available for Bremen and Uccle, but not for De Bilt, where a block profile is assumed in the MAX-DOAS retrieval and where column averaging kernels are estimated based on the box air mass factor for each observation layer. An overview of the station data is given by Table 6.1.



6.3 Inter-comparison method

Model VCDs (vertical column densities) have been calculated based on regional model data interpolated to MAX-DOAS output altitudes. Column averaging kernels (AVKs) from the measurements were applied to model NO₂ partial columns before summing up NO₂ values in the vertical:

$$VCD_{method2}^{model} = \sum_{i=1}^{Nobs} AVK_i \cdot VCD_i^{model}$$

The averaging kernels are part of the profiling output and represent the sensitivity of the retrieved column to the NO₂ amount at different altitudes. As the sensitivity of MAX-DOAS retrievals is largest in the boundary layer, the application of averaging kernels from the measurements to model simulations can have a crucial influence on validation results.

Only those model values closest to the measurement time are used below. As the model output is given in hourly time steps, the maximum possible time difference between measurements and simulations shown here is 30 minutes.

6.4 Results

Figure 6.1 shows time series of tropospheric NO₂ VCDs derived from MAX-DOAS for the model ENSEMBLE. The magnitude of VCDs from the measurements for the three urban stations is reproduced by the models, retrievals show a larger variability of values compared to the models. Measurements and simulations don't agree very well for some of the time steps investigated. The models underestimate some of the larger NO₂ values inside individual pollution plumes. Models may fail to reproduce these peaks due to errors in NO_x emissions, transport of NO₂ towards the stations and chemistry.

Vertical profiles of NO₂ partial columns are shown in Figure 6.2. The largest disagreements between regional models and retrievals occurs at lower observation altitudes, but the overall shape shows higher concentrations near the surface compared to higher altitudes for both, models and retrievals. CAMS-global shows an underestimation compared to MAX-DOAS at all levels for Uccle, the ENSEMBLE overestimates values at the lowest observation levels at this station. The Ensemble underestimates values at the lowest observation layers at Bremen but agrees well at other altitudes. Note that the number of vertical levels shown in Figure 6.2 corresponds to the number of MAX-DOAS output levels and that the real number of independent vertical levels of the retrievals is generally lower (about 2 to 3 degrees of freedom). MAX-DOAS observations are most sensitive in the boundary layer, whereas for uppermost output levels retrievals do not differ much from the a-priori assumed within the retrievals.




Figure 6.1. Time series of tropospheric NO₂ VCDs [10^{15} molec. cm^{-2}] from (black circles) MAX-DOAS and (red circles) the ENSEMBLE forecasts for June 2016 - February 2020 (De Bilt, Uccle), June 2016 - November 2019 (Bremen). Results for Bremen Dec 2019 - Feb 2020 will be added in the next report. Model results were calculated by multiplying NO₂ partial columns with averaging kernels for each observation layer followed by summing up resulting values in the vertical. Model data was interpolated to observation layer altitudes prior to calculation of VCDs.



Table 6.1. Overview of MAX-DOAS station data used for validation of regional air quality model simulations. The time period covered in this report is July 2016 to February 2020. MAX-DOAS retrievals for De Bilt and Bremen are not available for August to October as well as large parts of February 2016 due to the CINDI-2 measurement campaign. Data from the instrument at Uccle is available since January 2017.

Station	Latitude, longitude	Altitude above sea level	Institution	Quantity	Character
Bremen (Germany)	53.106°N, 8.86°E	21 m	IUP-UB	column/ profile	urban
De Bilt (Netherlands)	52.1° N, 5.18° E	23 m	KNMI	column	urban
Uccle (Belgium)	50.8° N, 4.32° E	120 m	BIRA-IASB	column/ profile	urban

Figure 6.3 shows comparisons of diurnal cycles. Mean column amounts are comparable for urban stations. Regional models perform significantly better than CAMS-global for De Bilt and Uccle, where CAMS-global is negatively biased. Some regional models show different variations from one hour to another especially during the morning for Bremen, where rush hour peaks simulated by the models are not confirmed by the retrievals. This may be related to different photochemistry, scaling of emissions in time or vertical distribution of NO₂ and errors in simulating pollution transport towards the station. With respect to the ENSEMBLE, the simulation of diurnal cycles of tropospheric NO₂ columns depends on the location, with good to moderate results at urban stations.

Comparisons of weekly cycles are shown in Figure 6.4. Weekly cycles are underestimated by all regional models, with a stronger decrease of NO₂ columns from workdays towards the weekend retrieved by MAX-DOAS for all urban stations. CAMS-global fails to reproduce weekly cycles, with hardly any reduction in the NO₂ columns visible towards the weekend. Note that some variations of values from one day to another may just be coincidence due to data sampling.




Figure 6.2 Vertical profiles of tropospheric NO₂ partial columns [10¹⁵ molec. cm⁻²] for (left) De Bilt and (right) Uccle for the time period after the recent upgrade of CAMS-global 10 July - 29 February 2020 (De Bilt, Uccle), 10 July -30 November (Bremen). The black and light grey lines show the MAX-DOAS retrievals and a priori used for MAX-DOAS retrieval, respectively. All other lines refer to model data: (red) ENSEMBLE, (yellow) CHIMERE, (brown) EMEP, (orange) SILAM, (purple) LOTOS-EUROS, (cyan) EURAD-IM, (pink) MOCAGE, (grey) MATCH and (blue) CAMS-global. Model data was interpolated to observation layer altitudes. MAX-DOAS vertical profiles are not available for De Bilt for the investigated time period and only model results are shown for these stations accordingly.




Figure 6.3. Weekly cycles (averages over daily bins divided by mean over whole week) of tropospheric NO₂ VCDs [10^{15} molec cm⁻²] for (top left) De Bilt, (top right) Uccle and (lower left) Bremen averaged over SON-2019. The black lines show the MAX-DOAS retrievals. All other lines refer to model data: (red) ENSEMBLE, (yellow) CHIMERE, (brown) EMEP, (orange) SILAM, (purple) LOTOS-EUROS, (cyan) EURAD-IM, (pink) MOCAGE, (grey) MATCH and (blue) CAMS-global. Period: June 2016 – February 2020 (De Bilt, Uccle), June 2016 - November 2019 (Bremen).

Scatter density plots or heat maps of tropospheric NO₂ VCDs from MAX-DOAS against model ENSEMBLE values corresponding to the time series displayed in Figure 6.1 as well as statistical values (root mean squared error, bias, correlation) are given in Figure 6.5. Corresponding statistical values for all individual models are given in Table 6.2. Moderate correlations on the order of 30-70 % are found for each station for all models, with the ENSEMBLE reaching the highest correlation of about 68 % at Uccle. Models tend to overestimate lower and underestimate higher NO₂ VCDs for the three urban stations. While the spread of values is quite large for individual data points, there is a good agreement between models and retrievals for the majority of measurements for urban stations (as shown by the high percentage of values close to the reference line). The ENSEMBLE performs significantly better than CAMS-global in terms of correlation for all stations. In contrast to the ENSEMBLE, CAMS-global has a strong negative bias at De Bilt and Uccle.




Figure 6.4. Weekly cycles (averages over daily bins divided by mean over whole week) of tropospheric NO₂ VCDs [10¹⁵ molec cm⁻²] for (top left) De Bilt, (top right) Uccle and (lower left) Bremen averaged over SON-2019. The black lines show the MAX-DOAS retrievals. All other lines refer to model data: (red) ENSEMBLE, (yellow) CHIMERE, (brown) EMEP, (orange) SILAM, (purple) LOTOS-EUROS, (cyan) EURAD-IM, (pink) MOCAGE, (grey) MATCH and (blue) CAMS-global. Period: June 2016 – February 2020 (De Bilt, Uccle), June 2016 - November 2019 (Bremen).

	De Bilt	Uccle	Bremen
ENS	6.159/1.577/0.441	5.467/1.186/0.686	4.792/0.459/0.550
CHIMERE	7.154/3.140/0.404	7.324/3.364/0.568	4.850/0.726/0.521
EMEP	7.100/2.580/0.403	6.534/2.539/0.629	6.210/1.910/0.504
SILAM	8.287/4.067/0.400	6.757/2.637/0.645	8.796/3.811/0.508
LOTOS-EUROS	6.601/0.407/0.305	6.139/0.028/0.563	5.852/-0.043/0.401
EURAD-IM	7.471/3.022/0.368	7.529/3.337/0.566	6.777/2.137/0.483
MOCAGE	5.896/-0.604/0.418	5.552/-0.876/0.662	4.632/-0.840/0.515
MATCH	6.467/1.469/0.422	5.666/0.824/0.656	5.064/-0.344/0.523
OSUITE	7.020/-3.468/0.294	7.967/-4.350/0.422	8.556/3.178/0.366

Table 6.2: Statistics on how tropospheric NO₂ VCDs [10¹⁵ molec. cm⁻²] from models compare to MAX-DOAS retrievals at the four stations. Each column entry shows from left to right: root mean squared error [10¹⁵ molec. cm⁻²], bias [10¹⁵ molec. cm⁻²] and correlation coefficient (cor). Period: June 2016 - February 2020 (De Bilt, Uccle), June 2016 - November 2019 (Bremen).




Figure 6.5. Scatter density plots of tropospheric NO₂ VCDs [10¹⁵ molec. cm⁻²] from MAX-DOAS against model ENSEMBLE hourly data for (top left) De Bilt, (top right) Uccle and (lower left) Bremen for DJF2020. The data is shown with a bin size of 1 x 10¹⁵ molec. cm⁻² and colour according to the number of data points per bin [%]. The dashed line is the reference line ($f(x) = x$). The solid line is the regression line (see top left of each plot for $f(x)$ of this line). The root mean squared error (rms) [10¹⁵ molec. cm⁻²], bias [10¹⁵ molec. cm⁻²], pearson correlation coefficient (cor) as well as the number of data points N are given at the top left of each plot. Period: June 2016 - February 2020 (De Bilt, Uccle), June 2016 – November 2019 (Bremen).



7 Validation of tropospheric NO₂ columns against satellite retrievals

7.1 Summary

Regional air quality model columns of tropospheric NO₂, derived from the output provided on 8 levels with a top at 5km, are compared to 9:30 local time GOME-2/MetOp-A NO₂ satellite retrievals (IUP-UB v1.0 product). The overall spatial distribution of tropospheric NO₂ is reproduced by the ENSEMBLE, but values over central European emission hotspots are significantly underestimated by the majority of the models during winter, which results in a strong underestimation over these regions and of the seasonal cycle for the ENSEMBLE. There are stronger shipping signals compared to the satellite data. As a result of a major regional model upgrade in June (2019), which includes the use of an updated European emissions inventory with improved estimates for North African and Middle Eastern anthropogenic emissions, enhanced tropospheric columns of NO₂ are reproduced over these regions by all models. Differences between models and satellite observations may result from errors in anthropogenic emissions, photochemistry during winter months and from chemical processing inside ship plumes. In contrast to the analysis, the regional ENSEMBLE forecast shows a negative bias compared to the retrievals which is most pronounced for winter ($\sim 2\text{-}3 \times 10^{15}$ molec/cm²) but smaller during the rest of the year (overall $\sim 0.5 \times 10^{15}$ molec/cm²). The negative bias is even larger for CAMS-global, which is in agreement with the stronger underestimation of values for European emission hotspots compared to regional models, demonstrating the benefit of running models at higher horizontal resolution. A systematic negative bias is however not present in the analysis for seasons other than winter.

7.2 Comparison with GOME-2 NO₂

In this section, regional air quality model columns of tropospheric NO₂ are compared to GOME-2/MetOp-A NO₂ satellite retrievals (IUP-UB v1.0) [Richter et al., 2011]. This satellite data provides excellent coverage in space and time and very good statistics. However, only integrated tropospheric columns are available, and the satellite data is always taken at 09:30 LT for GOME-2 and at clear sky only. Therefore, model data are vertically integrated, interpolated in time and then sampled to match the satellite data. Uncertainties in NO₂ satellite retrievals are large and depend on the region and season. Winter values in mid and high latitudes are usually associated with larger error margins. As a rough estimate, systematic uncertainties in regions with significant pollution are on the order of 20% – 30%. Conclusions may differ for comparisons to other satellite NO₂ products (e.g. TEMIS GOME-2, <http://www.temis.nl> shows lower retrieved NO₂ values for January). It should be noted here that model data is only available for altitudes up to 5000 m, meaning that (depending on tropopause height) tropospheric model columns may not be representative of the total amount of NO₂ in the troposphere. Note that since the CAMS-global upgrade of 26 June 2018, GOME-2 observations are assimilated by the global system. This is, however, a different retrieval product than what is used in the validation reported here (University of Bremen retrieval).




Figure 7.1. Maps of satellite-retrieved and model-simulated tropospheric NO_2 columns [molec cm^{-2}] for February 2020 for, from left to right: (top row) GOME-2, CAMS-global and CHIMERE; (middle row) EMEP, SILAM and LOTOS-EUROS; (lower row) EURAD-IM, MOCAGE and MATCH. GOME-2 data were gridded to regional model resolution (i.e. $0.1^\circ \times 0.1^\circ$). Model data were treated with the same reference sector ($25^\circ\text{W} - 20^\circ\text{E}$) subtraction approach as the satellite data and linearly interpolated to the satellite overpass time (9:30 LT).

Figure 7.1 shows maps of monthly mean tropospheric NO_2 columns from GOME-2, regional models and CAMS-global for February 2020. The overall spatial distribution and magnitude of tropospheric NO_2 is reproduced by the regional models in principle. There are stronger shipping signals in all models compared to the satellite data, which may result from errors in anthropogenic emissions or from chemical processing inside the ship exhaust plumes (see e.g. Vinken et al., 2014).

Compared to CAMS-global, regional models perform better for Central European emission hotspots, showing the benefit of higher horizontal resolution runs. As a result of a major regional model upgrade in June (2019), which includes the use of an updated European emissions inventory with improved estimates for North African and Middle Eastern anthropogenic emissions, enhanced tropospheric columns of NO_2 are now reproduced over these regions (e.g. Lebanon, Israel) by all models.




Figure 7.2. Time series of monthly averaged tropospheric NO₂ columns [10^{15} molec cm^{-2}] retrieved by (black) GOME-2 and simulated by (red) ENSEMBLE forecast, (green) ENSEMBLE analyses, (yellow) CHIMERE, (brown) EMEP, (orange) SILAM, (purple) LOTOS-EUROS, (cyan) EURAD-IM, (pink) MOCAGE, (grey) MATCH and (blue) CAMS-global. GOME-2 data were gridded to regional model resolution (i.e. $0.1^\circ \times 0.1^\circ$). Model data were treated with the same reference sector ($25^\circ\text{W} - 20^\circ\text{E}$) subtraction approach as the satellite data and linearly interpolated to the satellite overpass time (9:30 LT). Period: September 2016 – February 2020.

Figure 7.2 shows time series of monthly mean tropospheric NO₂ columns for GOME-2 and the models. The seasonal variation is better reproduced by SILAM, MOCAGE and EMEP than by the other models. The latter clearly underestimate the seasonal cycle over Europe due to the strong underestimation of values in winter described above. The regional ENSEMBLE forecast shows a negative bias compared to the retrievals which is most pronounced during winter ($\sim 2\text{--}3 \times 10^{15}$ molec/ cm^2) but smaller during the rest of the year (overall $\sim 0.5 \times 10^{15}$ molec/ cm^2). One of the reasons for this may be that the regional model output is limited to 5 km altitude. Compared to the ENSEMBLE forecast, the negative bias of CAMS-global is a bit larger, which is in agreement with the stronger underestimation over European emission hotspots for CAMS-global especially during winter. A systematic negative bias is not present in the regional model ENSEMBLE analysis for seasons other than winter. The decrease in retrieved wintertime maxima from 2017 to 2019, and increase in 2020 is not reproduced by the majority of regional models and CAMS-global.



8 Comparison with high-altitude EEA Air Quality e-reporting surface stations

8.1 Summary

European ozone EA Air Quality e-reporting measurements from high-altitude stations (above 1km) have been used to evaluate the regional models. Differences between the regional model orography and the true altitude of the station were used to select the model altitude level to compare with. The ensemble median mostly overestimates ozone levels during the period December 2019 - February 2020. More specific, depending on the station the observed ozone levels are reproduced to within -5% and 35% by the ensemble median D+0 forecast (1h-24h). Correlations observed were between 0.35 and 0.80 and the ensemble median D+0 forecast has a performance better than any of the individual seven models. CHIMERE and MOCAGE model was deviating significantly from the ensemble median in terms of MNMBs and LOTOS EUROS, MOCAGE and EURAD is deviating significantly from the ensemble median in terms of correlations with observations. Validation metrics are also given for the ENS analysis. The ENS analysis has almost equivalent performance with ENS D+0 forecast in terms of MNMBs but performs better than ENS D+0 forecast in terms of correlations (significantly higher correlations).

8.2 Introduction

The seven models and their ENSEMBLE median (D+0 forecast as well as the analysis) have been compared against Background-Rural EA Air Quality e-reporting measurements for surface stations at elevation greater than 1000 m above mean sea level (<http://www.eea.europa.eu/data-and-maps/data/airbase-the-european-air-quality-database-7>). Elevated stations were selected to fall within classes 1-2 in the O₃ Joly-Peuch (2012) classification for EA Air Quality e-reporting NRT stations. Table 8.1 shows the stations altitude above mean sea level together with the LOTOS-EUROS model altitude (i.e. from model's topography) pertaining to the nearest to the station grid point. Modelled gas mixing ratios were extracted at the model level, which is closest to the stations altitude as defined from the orography (see column 7 in Table 8.1).




Figure 8.1. Amplitude of the diurnal cycle as measured (ppb) from stations observations (black diamonds) and as calculated from the ENSEMBLE forecast D+0 (red circles). With asterisks are denoted stations used in the assimilation process.

Table.8.1: Background-Rural EEA Air Quality e-reporting Stations (with NRT data) with Elevation higher than 1000 m.

Station Name	Stat_id	Longitude	Latitude	real Altitude	model Altitude	nearest Level	use in CAMS50
Lario	ES1989A	-5.09	43.04	1140	1199	0	validation
Capmisabalos	ES0009R	-3.14	41.27	1360	1124	2	assimilation
Vilafranca	ES1435A	-0.25	40.42	1125	907	2	validation
Torrelisa	ES1883A	0.18	42.46	1005	1282	0	validation
Ak- Pardines	ES1310A	2.21	42.31	1226	1117	1	assimilation
Chaumont	CH0004R	6.98	47.05	1136	727	3	-
Rageade	FR07031	3.28	45.11	1040	944	2	validation
Schlucht	FR30028	7.01	48.05	1200	520	3	assimilation
Schauinsland	DEUB004	7.91	47.91	1205	554	3	assimilation
Rigi-Seebodenalp	CH0005R	8.46	47.07	1031	997	1	validation
Sulzberg im Bregenzerwald	AT80503	9.93	47.53	1020	961	1	assimilation
Bad Hindelang/Oberjoch	DEBY122	10.40	47.52	1169	1150	0	assimilation
Brocken	DEST039	10.62	51.80	1130	302	4	assimilation
Fichtelberg	DESN053	12.95	50.43	1214	555	3	assimilation
Vorhegg bei Kötschach-Mauthen	ATOVVOR1	12.97	46.68	1020	1427	0	validation
Churanov	CZ0CCHU	13.62	49.07	1118	739	4	assimilation
Krkonose-Rychory	CZ0HKRY	15.85	50.66	1001	530	4	assimilation
Bratislava - Jeséniova	SK0042A	20.99	48.78	1244	445	5	validation
Vitosha mountain	BG0070A	23.24	42.64	1321	863	4	-




Figure 8.2. MNMBs [%] (left) and temporal correlations (right) calculated for the ENSEMBLE model for during daytime (orange) night-time (dark blue) as well as for the whole day (red) for the DJF2020 period. With asterisks are denoted stations used in the assimilation process.

For the validation, hourly O₃ concentration values (µg/m³) are extracted from the seven models as well as for the Ensemble Mean. It should be noted that, in the EEA Air Quality e-reporting network the O₃ measurements that were made by the instrument in ppb were converted from ppb to µg/m³, following the EU directive 2008/50, i.e. by multiplying by 2. This conversion is approximately correct for low altitude stations. However, at high altitude stations pressure and temperature should be taken into account when converting from ppb to µg/m³ and vice versa. As hourly pressure and temperature data were not available for all EEA Air Quality stations the comparison between observed and modelled ozone was done by re-converting both modelled and observed hourly O₃ concentration in ppb. For modelled ozone values the conversion was done by applying the following ideal gas equation with the model’s estimates of temperature (T) and pressure (P) (from CAMS-global):

$$O_3 \text{ (in ppb)} = O_3 \text{ (in mg/m}^3\text{)} * \left(\frac{R * T}{P_m * M_{O_3}} \right)$$

8.3 Regional ensemble results

In the previous report it was shown that comparing the observed and modelled amplitude of the diurnal variation of ozone at each high-altitude station could provide a criterion concerning the exposure suitability of the stations. We found out in this report that an additional criterion is needed to differentiate stations as to their suitability in exposure. The additional criterion is the correlation coefficients between the amplitude of the diurnal cycle as observed and modelled to be statistically significant roughly higher than 0.3. Figure 8.1 shows the observed and modelled diurnal amplitude of ozone at each station, moving from Spain to Cyprus. Figure 8.2 shows the MNMBs and the correlation coefficients calculated for the ENSEMBLE model during daytime, at night-time as well as for the whole day. We can see that the 2 criteria of diurnal amplitude and day and night MNMBs and correlation coefficients differentiate the 2 stations in Spain from all other stations. For the above-mentioned findings these 2 high altitude stations will be excluded from our analysis.




Figure 8.3. O₃ MNMBs [%] (top) and correlation coefficient (bottom) for the Ensemble mean (forecast D+0; red circles and analysis; green triangles) as well as for CAMS-global (forecast D+0; blue diamonds) for the period DJF2020. With asterisks are denoted stations used in the assimilation process.




Fig. 8.4. Ozone MNMBs [%] for each one of the 7 models and CAMS-global (December 2019 to February 2020) for stations above 1000m altitude. With asterisks are denoted stations used in the assimilation process.

Figure 8.3 shows the Modified Normalized Mean biases (top) and correlation coefficients (bottom) at each of the remaining stations, moving from Spain to Bulgaria (i.e. from West to East) pertaining to the median of the Ensemble forecast (D+0) and analysis (D+0) as well as CAMS-global (D+0). The ENSEMBLE overestimates ozone levels during the period December 2019 to February 2020. Depending on the station the range of MNMB for the ENSEMBLE D+0 forecast was found to be between -5% and 35%. From Figure 8.3 (bottom panel) it is obvious that the ENSEMBLE reproduces well the ozone variability. As it appears from Figs 8.3 (bottom panel) the correlation coefficients are highly significant ($0.35 < r < 0.85$). It should be noted that for the period December to February 2020 the ENSEMBLE analysis performs better than the ENSEMBLE D+0 forecast in terms of correlations (significantly higher correlations).




Fig. 8.5: Correlation coefficients (from hourly values) between observed and modelled O₃ from all models of the ensemble and CAMS-global (o-suite) for December 2019 to February 2020. With asterisks are denoted stations used in the assimilation process.

8.4 Results for the seven regional models

Figure 8.4 shows Modified Normalized Mean biases at each station with elevation greater than 1000 m above mean sea level, moving from Spain to Bulgaria (from West to East) pertaining to each one of the 7 model calculations, the ENSEMBLE as well as CAMS-global. All results are based on the forecast D+0 elevated ozone values. On top of the graph shown is the elevation of the station. Depending on the station the observed ozone levels are reproduced to within -5% and 35% from the ENSEMBLE. Figure 8.4 shows that CHIMERE and MOCAGE models deviate significantly from the ensemble median (strong positive offset up to 50% and 45% respectively) while the remaining models show scores closer to the ENSEMBLE.

Finally, Figure 8.5 shows the correlations between observations and each model. It is clear that the Ensemble Mean reproduces well the ozone variability and has a better score than any of the individual models. The EURAD, MOCAGE and LOTOS EUROS models show lower correlation while the remaining models show scores closer to the ENSEMBLE.



9 Comparison with ozone sonde observations

9.1 Summary

Free tropospheric ozone (<850 hPa) could be reproduced by the ENSEMBLE forecast and analysis with MNMBs between 4.1% and 15% during DJF2020. The other models show MNMBs between -2% and 20% (forecasts and analysis). An exception is the MATCH model (SMH), which shows larger negative MNMBs (up to -7%) in the analysis.

9.2 Comparison approach

For the validation, the sonde profiles are compared to the model data closest in time. The model data is provided at the geographical coordinates of the sonde stations, the horizontal drift during the ascend of the sonde is considered negligible.

The model concentrations at the different height levels (0, 50, 250, 500, 1000, 2000, and 5000m above the ground) are matched to the respective sonde observations and are converted to mass mixing ratios. Pressure and temperature values needed for the conversion are taken from the sonde observations. For each station and all individual launches, the differences between observation and model are calculated. In order to be able to compare the profiles of different stations, this is done for fixed altitude levels between 0 and 6000m (interval for the surface 50m, above 100m, interval 100m). The sonde and model values are then aggregated to monthly means for each station and altitude level. For each month mean modified normalized biases (MNMB) are then calculated over all European stations for the free troposphere (<850 hPa).

9.3 Results for the ENSEMBLE

For the period February 2019 to February 2020, the ENSEMBLE forecast shows MNMBs between -3% and 15%. The ENSEMBLE analysis shows a similar behaviour with MNMBs between -7% and 14%, see Fig. 9.1.



Table 9.1: Sonde stations used in the validation for DJF2020

Station/location	Lat	lon	alt [m]
De Bilt	52.1	5.2	4
Hohenpeissenberg	47.8	11.2	976
Jokioinen	60.8	23.5	103
Legionow	52.4	20.97	96
Lerwik	60.14	-1.19	84
Madrid Spain	40.5	-3.8	631
Prag	50	14.4	302
Sodankyla	67	27	180
Uccle Belgium	51	4	100




Figure 9.1 - MNMBs for the regional ENSEMBLE and CAMS-global between February 2019 and February 2020 for the free troposphere region (<850 hPa).




Figure 9.2. MNMBs for the individual regional models between February 2019 and February 2020 for the free troposphere region (pressure < 850 hPa).




Figure 9.3. Vertical ENSEMBLE forecast and sonde comparison profiles for European sonde stations for December 2019, January and February 2020.




Figure 9.4. MNMBs for the regional models between December 2019 and February 2020 for the free troposphere region (pressure < 850 hPa).

9.4 Results for individual regional models

Between February 2019 and February 2020 regional model forecasts show MNMBs in the range of -11 % and 19%, see Fig. 9.2.

Results for the regional model analyses

Similar to the results of the individual models’ forecasts, the analyses show MNMBs between -12% and 20% (Fig. 9.4). Only the MACH model shows larger negative MNMBs up to -62%, which is in contrast to the forecast. This issue is attributed to not properly processed background error statistics. Correction was implemented in the first week of January 2020.




Figure 9.5. Vertical ENSEMBLE analysis and sonde comparison profiles for European sonde stations for December 2019, January and February 2020.



10 Comparison with GAW stations

10.1 Summary

The model concentrations of O₃ and CO at higher model levels were compared with five GAW high-altitude stations in mountainous terrain. As for the EEA air quality e-reporting stations, differences between the regional model orography and the true altitude of the station were used for this model level selection. Good results were obtained for the ENSEMBLE for ozone with small biases and good correlations. The CHIMERE model shows slightly larger MNMBs in both the analysis and the forecast. For CO, especially the ENSEMBLE corresponds well to the observations, however, small underestimates are found. The MATCH model behaves differently in the analysis than in the forecast, showing larger overestimations for CO than the rest of the models. The time series and correlation coefficients for CO and O₃ show that the ENSEMBLE reproduces for a large part the variability observed.

10.2 Comparison method

Hourly O₃ and CO concentration values in µg/m³ are extracted from the seven models and are compared to the GAW measurements, which were converted from volume mixing ratios (ppb) into concentrations by using pressure and temperature values at the respective pressure levels from the IFS model.

The altitude of the stations Hohenpeissenberg (HPB), Jungfraujoch (JFJ), Monte Cimone (CMN), Sonnblick (SNB) and Zugspitze (ZUG) in the model has been extracted from the orography as used in the LOTOS-EUROS model, see Table 10.1. For the level choice, the GAW stations' altitudes together with the best correlation of the corresponding levels were taken into account. Uncertainties due to the choice of level (calculated as mean differences between the chosen level and one up/down for the period DJF2020 for the ENSEMBLE) are up to ±30 µg/m³ for CO and up to ±4 µg/m³ for O₃.

Table 10.1 - Validation set-up for December 2019 - February 2020.

station	altitude station [m]	altitude model [m]	level choice (range 0-7)	altitude at level [m]
HPB	985	813	2	1063
JFJ	3580	1837	5	3837
CMN	2165	602	4	1602
SNB	3105	1687	3	2187
ZUG	2670	1348	4	2348




Figure 10.1. MNMBs [%] (left) and correlation coefficients (right) for ozone (red: ENSEMBLE forecast, green: ENSEMBLE analysis, blue: CAMS-global) for the period December 2019 to February 2020.




Figure 10.2. Time series plots for the ENSEMBLE forecast (red) and ENSEMBLE analysis (green) for surface O₃ in comparison with high altitude stations for the period December 2019 to February 2020.




Figure 10.3. Mean diurnal cycle of O₃ for the ENSEMBLE forecast (red) and ENSEMBLE analysis (green) compared to the observations (black) for the period DJF2020.




Figure 10.4. MNMBs [%] (left) and correlation coefficients (right) for all model forecasts for ozone (red: ENSEMBLE, blue: CAMS-global, yellow: CHIMERE, brown: EMEP, orange: SILAM, purple: LOTOS-EUROS, cyan: EURAD-IM, pink: MOCAGE, grey: MATCH). Altitudes are listed in Table 10.1.

10.3 Ozone

The ENSEMBLE forecast shows positive MNMBs between 5% and 18% and correlation coefficients ranging between 0.24 and 0.71 for the period December 2019 to February 2020. The ENSEMBLE analysis shows the lower MNMBs (between 4% and 16%) and very similar correlation coefficients ranging from 0.27 and 0.71 (see Fig. 10.1). In February, the overestimation of the ENSEMBLE runs increases compared to the observations.

The time series plots show a good correspondence between model and observations (Fig. 10.2). Diurnal cycles (Fig. 10.3) are not very pronounced at the GAW station locations.




Figure 10.5. MNMBs [%] (left) and correlation coefficients (right) for all models analyses for ozone (red: ENSEMBLE, blue: CAM5-global, yellow: CHIMERE, brown: EMEP, orange: SILAM, purple: LOTOS-EUROS, cyan: EURAD-IM, pink: MOCAGE, grey: MATCH). Altitudes are listed in Table 10.1.




Figure 10.6. MNMBs [%] (left) and correlation coefficients (right) for the ENSEMBLE for CO (red: ENSEMBLE forecast, green: ENSEMBLE analysis, blue: CAM5-global).

Results for individual model forecasts:

The models show MNMBs in the range of 34% and 0% for the period December 2019 to February 2020 (Fig. 10.4). The CHIMERE model shows larger positive MNMBs than the other models. During DJF2020, correlation coefficients vary greatly between the individual models and stations (0 to 0.7). The EURAD model shows low correlation coefficients.

Results for the individual model analyses:

For the individual model analyses (Fig. 10.5), MNMBs range between 34% and -2% and are very similar to the forecast. Same as for the forecast, correlation coefficients vary greatly between the different models and stations.




Figure 10.7. Time series plots for the ENSEMBLE forecast (red) and ENSEMBLE analysis (green) for surface CO in comparison with high altitude stations.




Figure 10.8. Mean diurnal cycle of CO for the ENSEMBLE forecast (red) and ENSEMBLE analysis (green) compared to the observations (black) for the period DJF2020.




Figure 10.9. CO MNMBs [%] (left) and correlation coefficients (right) for all regional model forecasts, the ENSEMBLE and CAMS-global (red: ENSEMBLE forecast, blue: CAMS-global, yellow: CHIMERE, brown: EMEP, orange: SILAM, purple: LOTOS-EUROS, cyan: EURAD-IM, pink: MOCAGE, grey: MATCH). Altitude ranges are listed in Table 10.1.




Figure 10.10. CO MNMBs [%] (left) and correlation coefficients (right) for all regional model analyses, the ENSEMBLE and CAMS-global (green: ENSEMBLE analysis, blue: CAMS-global, yellow: CHIMERE, brown: EMEP, orange: SILAM, purple: LOTOS-EUROS, cyan: EURAD-IM, pink: MOCAGE, grey: MATCH). Altitude ranges are listed in Table 10.1.

10.4 Carbon monoxide

For CO, the ENSEMBLE forecast and analysis MNMBs range between 8% and -9% for the analysis and between 1% and -13% for the forecast during December 2019 - February 2020. Correlation coefficients are between 0.25 and 0.74 for the analysis and between 0.31 and 0.86 for the forecast (Fig. 10.6). The time series plots (Fig. 10.7) show mostly a very good agreement between model and observations, except for a small vertical offset for some stations.

The comparison between mean diurnal observations and mean model concentrations show that the forecast partly has a negative offset for the high-altitude stations (Fig. 10.8). The analysis partly overestimates CO (SNB, ZUG, CMN).

CO results for individual model forecasts:

CO show MNMBs between 22% and -17% (Fig. 10.9). The CHIMERE model shows the largest positive MNMBs of all models, see Fig. 10.9. Correlation coefficients are between 0 and 0.86 for the forecasts. The EURAD model here shows a weak performance.

Results for the individual model analyses:



CO mixing ratios range between 23% and -17%, except for the MATCH model which shows large positive MNMBs (up to 53%) (Fig. 10.10). The MATCH model thus behaves differently in analysis and forecast, see Figs. 10.9 and 10.10. This was due to not properly processed background error statistics. A correction was implemented in MATCH in the first week of January 2020.

Correlation coefficients between the models vary significantly and range between 0 and 0.86.



11 Comparisons with MOPITT CO

11.1 Summary

In comparison with MOPITT v8 satellite data, the ENSEMBLE forecast data in general, are in good agreement with the observations, showing slightly negative bias within 10% (with some regional and temporal exceptions). The modelling data captured well both, relatively high CO values over the northern part of domain and relatively low values over the southern part. All the models are in good agreement with the observations. Analysis data have similar patterns compared to the forecast data, but with slightly stronger biases. Exception is the MATCH model, showing higher positive bias over the southern part of domain (up to 30%).

11.2 Method


CO total column forecasts over Europe from seven regional models and the model ENSEMBLE are compared with CO total column retrievals from MOPITT Version 7 (thermal infrared radiances) (Emmons et. al., 2009). Modelled CO data were converted from $\mu\text{g}/\text{m}^3$ to VMR by using temperature obtained from CAMS-global (o-suite) model. Pressure at the middle of the layers was also interpolated from the global model. Regional model data are available from the surface up to altitude of 5 km. For the comparison with satellite retrievals, the averaging kernels were applied to the modelled data.

Regional model data up to 5 km were merged with CAMS-global data above 5 km in order to minimize uncertainty error. We performed several confidence tests to establish the method. To check the error due to coarse sampling of the profiles up to 5 km as provided by the regional models, CAMS-global data were sampled at the height levels of the regional models up to 5 km and merged with the CAMS-global original levels above 5 km. Comparison of this results with the original CAMS-global data showed that the errors due to coarse sampling of the profiles up to 5 km were very small. Both results showed slight underestimation of the MOPITT data. CAMS-global values up to 5 km sampled at the height levels as the regional models and ENSEMBLE data without merging with the levels above 5 km show overestimation of the satellite data over almost entire region. From this we concluded that error due to missing values above 5 km is significant and merging the regional data with CAMS-global values above 5 km is necessary for the proper comparison.


MOPITT shows relatively high CO values over the northern part of domain and relatively low values over the southern part. The ENSEMBLE forecast data slightly overestimate/underestimate observations over the land/ocean part of domain (within 10%) in January. CHIMERE shows slightly smaller negative bias. In February, the ENSEMBLE data have negative bias over the almost entire domain (within 10% with some exceptions). Analysis data show slightly higher biases, both negative and positive. MATCH has bigger positive biases over the southern part of domain (up to 30%).




MOPITT




ENS-MOPITT
Forecast (0H24H)




EMEP-MOPITT
Forecast (0H24H)




EURAD-MOPITT
Forecast (0H24H)



CHIMERE-MOPITT
Forecast (0H24H)




LOTOS-EUROS-MOPITT
Forecast (0H24H)






SILAM-MOPITT
Forecast (0H24H)



MATCH-MOPITT
Forecast (0H24H)



MOCAGE-MOPITT
Forecast (0H24H)






Figure 11.1. CO total column for MOPITT V8 satellite retrievals (top row, in molecules/cm²), relative difference between the regional forecasts of the seven models and the ENSEMBLE and MOPITT (other rows) for December 2019 (left column), January (middle column) and February 2020 (right column). Grey colour indicates missing values.




MOPITT




ENS-MOPITT
Analysis (-24H-1H)




EMEP-MOPITT
Analysis (-24H-1H)




EURAD-MOPITT
Analysis (-24H-1H)



CHIMERE-MOPITT
Analysis (-24H-1H)




LOTOS-EUROS-MOPITT
Analysis (-24H-1H)






SILAM-MOPITT
Analysis (-24H-1H)



MATCH-MOPITT
Analysis (-24H-1H)



MOCAGE-MOPITT
Analysis (-24H-1H)




Figure 11.2. CO total column for MOPITT V8 satellite retrievals (top row, in molecules/cm²), relative difference between the regional analyses of the seven models and the ENSEMBLE and MOPITT (other rows) for December 2019 (left column) January (middle column) and February 2020 (right column). Grey colour indicates missing values.



12 Summary of findings for individual models

This section provides a short overview of the main, model specific findings of this report.

CHIMERE

Ozone and CO cross sections at the lateral boundaries show features that are not in line with what is seen in the other models (figs 3.11 and 3.12). This indicates a probable issue with the CAMS-global boundary condition implementation.

EURAD

Relatively large overestimations of ozone are spotted in the free troposphere compared to the other models and IAGOS measurements (fig 5.5). Ozone correlation coefficients at high altitude stations and GAW stations are much lower than for the other models in both forecasts and analyses (figs 8.5, 10.4 and 10.5). Similar results are found in the correlation coefficient for CO based on comparisons at the GAW stations (figs 10.9 and 10.10).

LOTOS-EUROS

The difference between the forecasts and the analysis is very high for PM10 over Africa. Similarly, the difference for ozone at 5000m also exhibits some unexpected features. Both differences are attributed to an erroneous initialisation setting in the analysis runs, which was corrected on February 21st. LOTOS-EUROS also appears to be overestimating some dust episodes, which is more evident for PM10 (fig. 3.17).

MATCH

The model seems to be over-correcting CO concentrations based on CO column comparisons with MOPITT V8 satellite retrievals (fig. 11.2). The reason in both cases has been identified and pin-pointed to not properly processed background error statistics. Correction was implemented in the first week of January 2020.

MOCAGE

High PM10 concentrations are spotted over the Atlantic, up to 250m above the surface (fig 3.7). The western boundary for PM10 is quite high, while the southern boundary is very low (fig 3.13), indicating issues with the implementation of the CAMS-global boundary conditions.



13 Acknowledgements

The authors acknowledge all EARLINET and European ACTRIS/Aeronet data providers for providing aerosol lidar profiles and sun photometer data available from the ACTRIS data portal (<http://actris.nilu.no>), and the Aeronet NRT data dissemination system respectively (<https://aeronet.gsfc.nasa.gov>). The ACTRIS 2 project (<http://www.actris.eu>) has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 654109.

We wish to acknowledge the provision of GAW hourly NRT station data by: the National Air Pollution Monitoring Network (NABEL) (Federal Office for the Environment FOEN and Swiss Federal Laboratories for Materials Testing and Research EMPA) for Jungfrauoch station, the Umweltbundesamt (UBA, Germany) for Zugspitze (Schneefernerhaus) station, the Umweltbundesamt (Austria) for Sonnblick station, the Observatory Hohenpeissenberg (Deutscher Wetter Dienst, DWD) for Hohenpeissenberg station, and the Institute of Atmospheric Sciences and Climate (ISAC) of the Italian National Research Council (CNR) for Monte Cimone station.

We wish to acknowledge the provision of ozone sonde data by the World Ozone and Ultraviolet Radiation Data Centre established at EC in Toronto (<http://woudc.org>), by the Data Host Facility of the Network for the Detection of Atmospheric Composition Change established at NOAA (<http://ndacc.org>), and by the Norwegian Institute for Air Research (<http://nilu.no>).

We acknowledge the EEA Air quality e-reporting Network (<https://www.eea.europa.eu/data-and-maps/data/aqereporting-8>) for the provision of hourly NRT station observations.

We wish to acknowledge the Department of Labour Inspection - Ministry of Labour and Social Insurance, of Cyprus (<http://www.airquality.dli.mlsi.gov.cy/>) for the provision of hourly NRT ozone data from Mountain Troodos station.

The programmes MOZAIC and CARIBIC, and the current Research Infrastructure IAGOS are operated with support from the European Commission, national agencies in Germany (BMBF), France (MESR), and the UK (NERC), and the IAGOS member institutions (<http://www.iagos.org/partners>). The participating airlines (Lufthansa, Air France, Austrian, China Airlines, Iberia, Cathay Pacific, Air Namibia, Sabena) supported IAGOS by carrying the measurement equipment free of charge since 1994. The data are available at <http://www.iagos.fr> thanks to additional support from AERIS (CNRS and CNES).

GOME2 lv1 radiances and irradiances were provided by EUMETSAT.

We acknowledge the NASA Langley Research Center Atmospheric Science Data Center for providing the MOPITT data.

AUTH acknowledges the AUTH Scientific Computing Centre (<https://it.auth.gr/en/services>) for providing technical and infrastructure support for data analysis performed in this WP.



14 References

Richter, A., Begoin, M., Hilboll, A., and Burrows, J. P.: An improved NO₂ retrieval for the GOME-2 satellite instrument, *Atmos. Meas. Tech.*, **4**, 1147-1159, doi:10.5194/amt-4-1147-2011, 2011.

Ackermann, J. (1998). The extinction-to-backscatter ratio of tropospheric aerosol: A numerical study. *Journal of atmospheric and oceanic technology*, **15**(4), 1043-1050.

ACTRIS Deliverable WP6/D6.21,

http://www.actris.net/Portals/97/deliverables/PU/WP6_D6.21_M45v2.pdf

Catrrall, C., Reagan, J., Thome, K., & Dubovik, O. (2005). Variability of aerosol and spectral lidar and backscatter and extinction ratios of key aerosol types derived from selected Aerosol Robotic Network locations. *Journal of Geophysical Research: Atmospheres*, **110**(D10).

Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., ... & Nakajima, T. (2002). Tropospheric aerosol optical thickness from the GOCART model and comparisons with satellite and Sun photometer measurements. *Journal of the atmospheric sciences*, **59**(3), 461-483.

Eskes, H.J., S. Basart, A. Benedictow, Y. Bennouna, A.-M. Blechschmidt, S. Chabrillat, Y. Christophe, E. Cuevas, H. Flentje, K. M. Hansen, J. Kapsomenakis, B. Langerock, M. Ramonet, A. Richter, M. Schulz, N. Sudarchikova, A. Wagner, T. Warneke, C. Zerefos, Observation characterisation and validation methods document, Copernicus Atmosphere Monitoring Service (CAMS) report, December 2019, doi: 10.24380/Onsd- wb26.

Eskes, H., Huijnen, V., Arola, A., Benedictow, A., Blechschmidt, A.-M., Botek, E., Boucher, O., Bouarar, I., Chabrillat, S., Cuevas, E., Engelen, R., Flentje, H., Gaudel, A., Griesfeller, J., Jones, L., Kapsomenakis, J., Katragkou, E., Kinne, S., Langerock, B., Razinger, M., Richter, A., Schultz, M., Schulz, M., Sudarchikova, N., Thouret, V., Vrekoussis, M., Wagner, A., and Zerefos, C.: Validation of reactive gases and aerosols in the MACC global analysis and forecast system, *Geosci. Model Dev.*, **8**, 3523-3543, doi:10.5194/gmd-8-3523-2015, 2015.

Flemming, J., Huijnen, V., Arteta, J., Bechtold, P., Beljaars, A., Blechschmidt, A.-M., Diamantakis, M., Engelen, R. J., Gaudel, A., Inness, A., Jones, L., Josse, B., Katragkou, E., Marecal, V., Peuch, V.-H., Richter, A., Schultz, M. G., Stein, O., and Tsikerdekis, A.: Tropospheric chemistry in the Integrated Forecasting System of ECMWF, *Geosci. Model Dev.*, **8**, 975-1003, doi:10.5194/gmd-8-975-2015, 2015.

Joly, Mathieu, and Vincent-Henri Peuch, Objective classification of air quality monitoring sites over Europe, *Atmospheric Environment* **47**, 111-123, 2012.

Katragkou, E., Zanis, P., Tsikerdekis, A., Kapsomenakis, J., Melas, D., Eskes, H., Flemming, J., Huijnen, V., xsxInness, A., Schultz, M. G., Stein, O., and Zerefos, C. S.: Evaluation of near-surface ozone over Europe from the MACC reanalysis, *Geosci. Model Dev.*, **8**, 2299-2314, doi:10.5194/gmd-8-2299-2015, 2015.

Marécal, V., Peuch, V.-H., Andersson, C., Andersson, S., Arteta, J., Beekmann, M., Benedictow, A., Bergström, R., Bessagnet, B., Cansado, A., Chéroux, F., Colette, A., Coman, A., Curier, R. L., Denier van der Gon, H. A. C., Drouin, A., Elbern, H., Emili, E., Engelen, R. J., Eskes, H. J., Foret, G., Friese, E., Gauss, M., Giannaros, C., Guth, J., Joly, M., Jaumouillé, E., Josse, B., Kadygrov, N., Kaiser, J. W., Krajsek, K., Kuenen, J., Kumar, U., Liora, N., Lopez, E., Malherbe, L., Martinez, I., Melas, D., Meleux, F., Menut, L., Moinat, P., Morales, T., Parmentier, J., Piacentini, A., Plu, M., Poupkou, A., Queguiner, S., Robertson, L., Rouil, L., Schaap, M., Segers, A., Sofiev, M., Tarasson, L., Thomas, M., Timmermans, R., Valdebenito, Á., van Velthoven, P., van Versendaal, R., Vira, J.,



and Ung, A.: A regional air quality forecasting system over Europe: the MACC-II daily ensemble production, *Geosci. Model Dev.*, **8**, 2777-2813, doi:10.5194/gmd-8-2777-2015, 2015.

Morcrette, J.-J., O. Boucher, L. Jones, D. Salmond, P. Bechtold, A. Beljaars, A. Benedetti, A. Bonet, J. W. Kaiser, M. Razinger, M. Schulz, S. Serrar, A. J. Simmons, M. Sofiev, M. Suttie, A. M. Tompkins, and A. Untch: Aerosol analysis and forecast in the ECMWF Integrated Forecast System. Part I: Forward modelling, *J. Geophys. Res.*, **114**, D06206, doi:10.1029/2008JD011235, 2009.

Mortier, A., Goloub, P., Derimian, Y., Tanné, D., Podvin, T., Blarel, L., ... & Ndiaye, T. (2016). Climatology of aerosol properties and clear-sky shortwave radiative effects using Lidar and Sun photometer observations in the Dakar site. *Journal of Geophysical Research: Atmospheres*.

Müller, D., Ansmann, A., Mattis, I., Tesche, M., Wandinger, U., Althausen, D., & Pisani, G. (2007). Aerosol-type-dependent lidar ratios observed with Raman lidar. *Journal of Geophysical Research: Atmospheres*, **112**(D16).

Omar, A. H., Winker, D. M., Vaughan, M. A., Hu, Y., Trepte, C. R., Ferrare, R. A., ... & Kuehn, R. E. (2009). The CALIPSO automated aerosol classification and lidar ratio selection algorithm. *Journal of Atmospheric and Oceanic Technology*, **26**(10), 1994-2014.

Pappalardo, G., A. Amodeo, A. Apituley, A. Comeron, V. Freudenthaler, H. Linne, A. Ansmann, J. Bösenberg, G. D'Amico, I. Mattis, L. Mona, U. Wandinger, V. Amiridis, L. Alados-Arboledas, D. Nicolae, and Wiegner, M.: EARLINET: towards an advanced sustainable European aerosol lidar network, *Atmos. Meas. Tech.*, **7**, 2389–2409, www.atmos-meas-tech.net/7/2389/2014/, doi:10.5194/amt-7-2389-2014, 2014.

Richter, A., Begoin, M., Hilboll, A., and Burrows, J. P.: An improved NO₂ retrieval for the GOME-2 satellite instrument, *Atmos. Meas. Tech.*, **4**, 1147-1159, doi:10.5194/amt-4-1147-2011, 2011.

Vinken, G. C. M., Boersma, K. F., van Donkelaar, A., and Zhang, L.: Constraints on ship NO_x emissions in Europe using GEOS-Chem and OMI satellite NO₂ observations, *Atmos. Chem. Phys.*, **14**, 1353-1369, doi:10.5194/acp-14-1353-2014, 2014.

Wagner, A., M. Schulz, Y. Christophe, M. Ramonet, H. J. Eskes, S. Basart, A. Benedictow, Y. Bennouna, A.-M. Blechschmidt, S. Chabrillat, E. Cuevas, A. El-Yazidi, H. Flentje, K.M. Hansen, U. Im, J. Kapsomenakis, B. Langerock, A. Richter, N. Sudarchikova, V. Thouret, T. Warneke, C. Zerefos, Validation report of the CAMS near-real-time global atmospheric composition service: Period September - November 2019, Copernicus Atmosphere Monitoring Service (CAMS) report, CAMS84_2018SC2_D1.1.1_SON2019_v1.pdf, February 2020, doi:10.24380/xzkk-bz05.

